

Table 24. Summary of ACR 20 Response at Week 24 (Sensitivity Analysis 4)

ACR 20 Response at Week 24 (Sensitivity Analysis 4)			
	Placebo	Ustekinumab	
		45 mg	90 mg
Study CNTO1275PSA3001			
Subjects randomized	206	205	204
ACR 20			
N	206	205	204
Subjects in response	65 (31.6%)	96 (46.8%)	104 (51.0%)
p-value		<0.001	<0.001
Study CNTO1275PSA3002			
Subjects randomized	104	103	105
ACR 20			
N	104	103	105
Subjects in response	21 (20.2%)	45 (45.7%)	46 (43.8%)
p-value		<0.001	<0.001

Source: 24-wk CNTO1275PSA3001 Clinical Study Report, Att.3.4, page 266; 24-wk CNTO1275PSA3002 Clinical Study Report, Att.3.4, page 276.

Robert Abugov, PhD, (primary statistical reviewer) further explores these different sensitivity analyses in his review. Please see his review for detailed analysis.

ACR 20 Response Over Time

Figure 6 and Figure 7 plot out the proportion of ACR 20 responders over time (documented at each visit through Week 24). Both studies have a steady increase in response over time with a maximum response around Weeks 20 and 24. For both studies, there appears to be a decline in ACR 20 responders at Week 16 for subjects receiving ustekinumab, which would coincide with the first dose of maintenance therapy (after loading doses at Weeks 0 and 4 and no dosing for 12 weeks).