

prefilled syringe is first removed from the refrigerator on the carton in the space provided. When a syringe has been stored at room temperature, it should not be returned to the refrigerator. Throw away (discard) the syringe if not used within 15 days at room temperature storage. Do not use STEQEYMA after the expiration date on the carton or the prefilled syringe.

- Keep STEQEYMA and all medicines out of the reach of children.

Parts of the Prefilled Syringe (see Figure B)




Figure B

Preparing to Inject STEQEYMA

Carton containing prefilled syringe Cotton ball or gauze and alcohol swab Adhesive bandage Sharps disposal container	1. Gather the supplies for the injection 1.a. Prepare a clean, flat surface, such as a table or counter top, in a well-lit area. 1.b. Take the carton(s) containing the prefilled syringe(s) needed to administer your prescribed dose out of the refrigerator. Each STEQEYMA carton contains 1 prefilled syringe. Note: Depending on the dose prescribed to you by your doctor you may need to prepare 1 or more prefilled syringes and inject the contents of them all. 1.c. Make sure you have the following supplies (see Figure C) - Carton containing prefilled syringe Not included in the carton: - Cotton ball or gauze - Adhesive bandage - FDA-cleared sharps disposal container - Alcohol swab
EXP: YYYY/MM EXP:	2. Check the expiration date on the carton (see Figure D). • Do not use it if the expiration date has passed. If the expiration date has passed, safely throw away (dispose of) the carton in an FDA-cleared sharps disposal container (see Step 13. Throw away (dispose of) STEQEYMA).

Figure C

Figure D