




Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965



Blinding and Salting in SuperCDMS

BEN LOER

Pacific Northwest National Laboratory

Blind Analysis in High-Stakes Survey Science: When, Why, and How?

On behalf of the SuperCDMS Collaboration





Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Outline

- ▶ Dark matter direct detection signals
- ▶ SuperCDMS detectors
- ▶ Traditional box blinding
- ▶ Salting in current analysis
- ▶ Looking ahead






Pacific Northwest
NATIONAL LABORATORY


Proudly Operated by Battelle Since 1965

Direct detection of dark matter

- ▶ Dark matter is non-rotating, non-interacting gas with thermal distribution
- ▶ Interaction velocity ~ 250 km/s
- ▶ Billiard ball physics
- ▶ Exponential spectrum of low energy nuclear recoils (NRs)
- ▶ Current limits: few counts per ton*year for ~ 100 GeV mass
- ▶ Lower mass requires lower threshold





WIMP-induced Nuclear Recoil Spectrum



SuperCDMS iZIPs: *interleaved Z-sensitive Ionization and Phonon sensors*

- ▶ Ultra pure Ge and Si crystals operated at ~50 mK
- ▶ Simultaneously measure ionization and phonon signals




SuperCDMS iZIPs: interleaved Z-sensitive Ionization and Phonon sensors




Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

- ▶ Ultra pure Ge and Si crystals operated at ~50 mK
- ▶ Simultaneously measure ionization and phonon signals
- ▶ Outer guard ring and interleaved charge sensors reject surface events




Surface events: only detect charge on one face



SuperCDMS iZIPs: interleaved Z-sensitive Ionization and Phonon sensors


- ▶ Ultra pure Ge and Si crystals operated at ~50 mK
- ▶ Simultaneously measure ionization and phonon signals
- ▶ Outer guard ring and interleaved charge sensors reject surface events
- ▶ Ionization/phonon signal less for nuclear recoil (signal-like) events than electron-recoil (Compton backgrounds)





SuperCDMS Traditional box blinding

- ▶ Take signal and background calibration data
- ▶ Identify signal region of interest (ROI)
- ▶ Remove events in a slightly larger region
- ▶ Tune cuts
- ▶ Open the box
- ▶ No attempt to subtract backgrounds remaining in the box






Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

SuperCDMS Traditional box blinding


- ▶ Take signal and background calibration data
- ▶ Identify signal region of interest (ROI)
- ▶ Remove events in a slightly larger region
- ▶ Tune cuts
- ▶ Open the box
- ▶ No attempt to subtract backgrounds remaining in the box





SuperCDMS Traditional box blinding

- ▶ Take signal and background calibration data
- ▶ Identify signal region of interest (ROI)
- ▶ Remove events in a slightly larger region
- ▶ Tune cuts
- ▶ Open the box
- ▶ No attempt to subtract backgrounds remaining in the box





Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Problems with box blinding

- ▶ Delayed access to data: first must acquire and analyze calibration data to determine the box, then apply blinding cut
- ▶ Unanticipated backgrounds
- ▶ Anomalous events






Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

That's what dark matter looks like, right?




No cell phones near the detectors!



Case Study: SuperCDMS Low Threshold iZIP analysis

- ▶ Boosted Decision Tree (BDT) machine learning used to optimize signal/background identification
- ▶ Predicted 6 events in signal region, found 11 at unblinding
- ▶ Most significant leakage at high energy from single detector with shorted ionization channel
- ▶ Would earlier inspection have led to better modeling?
- ▶ Even on post-inspection, high energy events appear good quality, so not clear that non-blind analysis would have changed result






Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965



Pushing to lower mass



CDMSlite (low ionization threshold experiment) => SuperCDMS HV

- ▶ Electrons/holes gain energy from voltage bias, release as phonons
- ▶ High bias => high gain charge signal
- ▶ BUT lose “true” phonon signal and yield discrimination variable
- ▶ If “box” blinding, what is second parameter?
- ▶ If just blind entire energy range, no sidebands left!

2% resolution at 1 keV







Case Study: CDMSlite Run 2

- ▶ Source of clustered events in signal region spontaneously appeared
- ▶ Would not have been visible in box blinding analysis

y position



Radial Parameter [arb. unit]





Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

CDMSlite Run 2 vs Run 3

- ▶ CDMSlite at Soudan primarily R&D; thus far was non-blind analysis
- ▶ Run 3 data has been acquired
- ▶ Different detector with lower threshold, slightly less exposure
- ▶ Opportunity to test new analysis techniques for next generation SuperCDMS SNOLAB
- ▶ E.g., partial background modeling/subtraction
- ▶ With perils of box blinding recently shown, collaboration decided to try salting our data with fake signal events as blind analysis technique





Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Attempting data salting for SuperCDMS


- ▶ Principle: salt not **accidentally** identifiable
 - A sufficiently motivated malicious actor can always find a way
- ▶ The more signal-like the better, but does not need to follow signal distributions exactly
 - E.g. can not use salt to measure signal acceptance. Don't expect to recover "right" answer if a WIMP signal input
 - Salt tails are OK so long as small fraction of total salt
 - But presence of salt should not lead to unnecessary cuts
- ▶ Can't insert events at will with sequential event IDs
- ▶ Therefore method is to select events with correct metadata (e.g. trigger on correct detector) and replace waveform portion of data only





Salting method

- ▶ Pick a WIMP mass, estimate our 90% CL predicted sensitivity, calculate how many events would be in the final spectrum, call that N
- ▶ Randomly select between N and 3N events in raw data to replace with salt
- ▶ Weight selection to be uniform in time






Salt Spectrum

- ▶ Assign energy from exponential+constant spectrum
- ▶ Slope varies from 0.5-2 keV, expo/constant varies from 1/3 to 3

$3 \cdot \exp(-x/0.5) + \text{const}$

$\exp(-x) + \text{const}$


$1/3 \cdot \exp(-x/2) + \text{const}$





Creating fake pulses

- ▶ Select calibration events with similar energy
 - Want to avoid modeling distributions as much as possible
 - Lots of energy dependence, so tight energy selection
 - Hard to use real calibration events due to pileup, falling baselines (i.e. near pileup), etc
- ▶ Fit each channel of calibration events to 2 templates (average + residual)
- ▶ Scale template amplitudes to new target energy (keeping same trigger delay values)
- ▶ Add to empty trigger to sample noise






Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Creating fake pulses

- ▶ Select calibration events with similar energy
 - Want to avoid modeling distributions as much as possible
 - Lots of energy dependence, so tight energy selection
 - Hard to use real calibration events due to pileup, falling baselines (i.e. near pileup), etc
- ▶ Fit each channel of calibration events to 2 templates (average + residual)
- ▶ Scale template amplitudes to new target energy (keeping same trigger delay values)
- ▶ Add to empty trigger to sample noise







Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Can you spot the difference?








Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

First attempt at salting

- ▶ Was supposed to be 2-5% of final dataset
- ▶ Supposed to be exponential + constant
- ▶ Clustering from using same calibration events multiple times







Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

First attempt at salting

- ▶ Was supposed to be 2-5% of final dataset
- ▶ Supposed to be exponential + constant
- ▶ Clustering from calibration ev






Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Current status



- ▶ We now have better understanding of effect of cuts on salt events
- ▶ Unclear what factors should drive min/max amount of salt to insert
 - Is it all in our heads?
 - 90%CL * rand(1,3) arbitrary
- ▶ Plan to generate salted data for CDMSlite Run 3 analysis next week!
- ▶ Before release, one analyzer will perform quality control inspection and be removed from further analysis



The future: SuperCDMS



- ▶ 2 km under Sudbury, Ontario
- ▶ Entire lab is class 2000 clean room
- ▶ Bigger, cleaner shield
- ▶ Bigger, more advanced detectors
- ▶ Plan for salting as main blinding strategy






Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Salting in SuperCDMS SNOLAB

- ▶ Just beginning to think about this now
- ▶ Faking signal events with brand new detector types will not be easy
- ▶ Bandwidth, disk are cheap => DAQ stores **piles** of metadata
- ▶ Salting and DAQ efforts will be loosely integrated
- ▶ Once calibration preselection is in place, hope to have data available to analyzers within days instead of months





Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965

Thank you!



[California Inst. of Tech.](#)



[CNRS-LPN*](#)



[Durham University](#)



[FNAL](#)



[NISER](#)



[NIST*](#)



[Northwestern](#)



[PNNL](#)



[Queen's University](#) [Santa Clara University](#)



[SLAC](#)



[South Dakota SM&T](#)



[SMU](#)



[SNOLAB](#)



[Stanford University](#)



[Texas A&M University](#)



[U. British Columbia](#)



[U. California, Berkeley](#)



[U. Colorado Denver](#)



[U. Evansville](#)



[U. Florida](#)



[U. Minnesota](#)



[U. South Dakota](#)



[U. Toronto](#)

* Associate members



Pacific Northwest
NATIONAL LABORATORY


Proudly Operated by Battelle Since 1965

Backup slides





SuperCDMS SNOLAB trigger






Pacific Northwest
NATIONAL LABORATORY


Proudly Operated by Battelle Since 1965

Bridging the technological divide






Bridging the technological divide






Bridging the technological divide





Bridging the technological divide




Yield and charge fiducial volume background discrimination




Pacific Northwest
NATIONAL LABORATORY

Proudly Operated by Battelle Since 1965




- Surface leakage < 1.26E-5 (90% CL)
 - for ~50% signal acceptance
- Not using phonon position info





Problems with box blinding


- ▶ Delayed access to data: first must acquire and analyze calibration data to determine the box, then apply blinding cut
- ▶ Hidden trends →
- ▶ Unanticipated backgrounds
- ▶ Anomalous events





Problems with box blinding

- ▶ Delayed access to data: first must acquire and analyze calibration data to determine the box, then apply blinding cut
- ▶ Hidden trends →
- ▶ Unanticipated backgrounds
- ▶ Anomalous events





Problems with box blinding

- ▶ Delayed access to data: first must acquire and analyze calibration data to determine the box, then apply blinding cut
- ▶ Hidden trends →
- ▶ Unanticipated backgrounds
- ▶ Anomalous events

