

Pass the salt: LUX's alternative to blinding

Aaron Manalaysay
UCDAVIS
UNIVERSITY OF CALIFORNIA

for the *LUX* collaboration



Blind Analysis in High-Stakes Survey Science
SLAC, Palo Alto, CA, 13 March 2017

The LUX collaboration



Berkeley Lab / UC Berkeley

Bob Jacobsen	PI, Professor
Murdock Gilchriese	Senior Scientist
Kevin Lesko	Senior Scientist
Michael Witherell	Lab Director
Peter Sorensen	Scientist
Simon Fiorucci	Project Scientist
Attila Dobi	Postdoc
Daniel Hogan	Graduate Student
Kate Kamdin	Graduate Student
Kelsey Oliver-Mallory	Graduate Student



Brown University

Richard Gaitskell	PI, Professor
Samuel Chung	Graduate Student
Dongqing Huang	Graduate Student
Casey Rhyne	Graduate Student
Will Taylor	Graduate Student
James Verbus	Postdoc



University of Edinburgh, UK

Alex Murphy	PI, Professor
Paolo Beltrame	Research Fellow
Tom Davison	Graduate Student
Maria F. Marzoni	Graduate Student



Imperial College London

Henrique Araujo	PI, Reader
Tim Sumner	Professor
Alastair Currie	Postdoc
Adam Bailey	Graduate Student
Khadeeja Yazdani	Graduate Student



Lawrence Livermore

Adam Bernstein	PI, Leader of Adv. Detectors Grp.
Kareem Kazkaz	Staff Physicist
Jingke Xu	Postdoc
Brian Lenardo	Graduate Student



LIP Coimbra, Portugal

Isabel Lopes	PI, Professor
Jose Pinto da	Assistant Professor
Vladimir Solovov	Senior Researcher
Francisco Neves	Auxiliary Researcher
Alexander Lindote	Postdoc
Claudio Silva	Postdoc
Paulo Bras	Graduate Student



SLAC Stanford (CWRU)

Dan Akerib	PI, Professor
Thomas Shutt	PI, Professor
Tomasz Biesiadzinski	Research Associate
Christina Ignarra	Research Associate
Wing To	Research Associate
Rosie Bramante	Graduate Student
Wei Ji	Graduate Student
T.J. Whitis	Graduate Student



SD Mines

Xinhua Bai	PI, Professor
Doug Tiedt	Graduate Student



SDSTA / Sanford Lab

David Taylor	Project Engineer
Markus Horn	Research Scientist
Dana Byram	Support Scientist



University at Albany

Matthew Szydagis	PI, Professor
Jeremy Mock	Postdoc
Sean Fallon	Graduate Student
Jack Genovesi	Graduate Student
Steven Young	Graduate Student



Texas A&M University

James White †	PI, Professor
Robert Webb	PI, Professor
Rachel Mannino	Graduate Student
Paul Terman	Graduate Student

Berkeley / UC Berkeley (Yale)

Daniel McKinsey	PI, Professor
Ethan Bernard	Project Scientist
Scott Hertel	Postdoc
Kevin O'Sullivan	Postdoc
Elizabeth Boulton	Graduate Student
Evan Pease	Graduate Student
Brian Tennyson	Graduate Student
Lucie Tvrznikova	Graduate Student
Nicole Larsen	Graduate Student



UC Davis

Mani Tripathi	PI, Professor
Britt Hollbrook	Senior Engineer
John Thomson	Development
Dave Hemer	Senior Machinist
Ray Gerhard	Electronics Engineer
Aaron Manalaysay	Project Scientist
Jacob Cutter	Graduate Student
James Morad	Graduate Student
Sergey Uvarov	Graduate Student



UC Santa Barbara

Harry Nelson	PI, Professor
Susanne Kyre	Engineer
Dean White	Engineer
Carmen Carmona	Postdoc
Scott Haselschwardt	Graduate Student
Curt Nehrkorn	Graduate Student
Melih Solmaz	Graduate Student



University College London, UK	University College London, UK
Chamkaur Ghag	PI, Lecturer
James Dobson	Postdoc
Sally Shaw	Graduate Student



University of Maryland

Carter Hall	PI, Professor
Jon Balajthy	Graduate Student
Richard Knoche	Graduate Student
Frank Wolfs	PI, Professor
Wojtek Skutski	Senior Scientist
Eryk Druszkiewicz	Graduate Student
Dev Ashish Khaitan	Graduate Student
Diktat Koyuncu	Graduate Student
M. Moongweluwan	Graduate Student
Jun Yin	Graduate Student



University of South Dakota


Dongming Mei	PI, Professor
Chao Zhang	Postdoc
Kimberly Palladino	PI, Asst Professor
Shaun Alsum	Graduate Student




University of Wisconsin

Kimberly Palladino	PI, Asst Professor
Shaun Alsum	Graduate Student

Dark matter direct detection in a nutshell




Sanford Underground Research Facility




Detection technique

- Dual-phase Xenon Time Projection Chamber (TPC)
- E-dep:
 - ▶ primary scintillation (S1)
 - ▶ ionization (S2)(both measured by PMTs)
- PMT hit-pattern gives x - y position.
- S1-S2 delay time gives z position
- Signal: nuclear recoils
- Background: [mostly] electronic recoils.



Data parameter space


LUX's first science data (95 live days, non-blind)



- The two energy variables, scintillation (S1) and ionization (S2), permit distinction between Electronic Recoils and Nuclear Recoils.
- Analysis involves also three position variables.

Data parameter space

LUX's first science data (95 live days, non-blind)



- Traditional blinding: one masks the signal region from view until analysis is finalized.
- Electronic Recoils (BG)
- Nuclear Recoils (Signal)

Two important calibrations

- **CH₃T** (tritiated methane): Used to define our ER band (i.e. background). After injection, mixes uniformly in the detector in minutes. Removed on a ~8 hr timescale by our purification system.
- **Deuterium-Deuterium** neutron generator: Used to define our NR band (i.e. signal). Monoenergetic, collimated neutron beam.

Blinding problems

Dark matter direct detection searches are:

- low-background
- rare signal event
- **Blinding in this type of search means one is also blind to rare backgrounds.**
- Un-modeled rare BGs are not a showstopper for a 1-sided search (i.e. limit-setting only), but they are quite problematic for a 2-sided search.

∃ plenty of examples in this field of blind searches “discovering” un-modeled, pathological backgrounds. E.g.:

- XENON10: PRL **100** (2008) 021303 (arXiv:0706.0039)
- XENON100: PRL **109** (2012) 181301 (arXiv:1207.5988)
- ZEPLINIII: PRD **80** (2009) 052010 (arXiv:0812.1150)
- SuperCDMS: PRL **112** (2014) 241302 (arXiv:1402.7137)

Blinding 1

Dark matter direct detection

- low-background
- rare signal event

• Blinding in this type of experiment is very difficult because there is no simple filter that can tell the number of events.

• Unblinding is often done by opening a "signal box" (i.e. searching for a signal in a subset of the data).

• Example: unblinding a dark matter search.

BLIND ANALYSIS IN NUCLEAR AND PARTICLE PHYSICS

Joshua R. Klein

*Department of Physics, University of Texas, Austin, Texas 78712;
email: jrk@physics.utexas.edu*

Aaron Roodman

*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309;
email: roodman@slac.stanford.edu*


cover there is no simple filter that can tell the number of events. Although a hidden signal box approach like that described in Section 3.1 can be used for many

of the [traditional blinding] prevents the experimenter from being able to examine the characteristics of the signal, and hence carries the kind of risk discussed in Sec-

In addition, a hidden signal box approach assumes that the characteristics of the backgrounds are known well enough that nothing unexpected will be discovered when the signal box is opened.

A very general approach to blind analysis which is appropriate for counting experiments is to split the event count itself in an unknown way. The splitting can be done by adding an unknown set of false signal events, by removing a small unknown number of all events from the data set, or by doing both.


Blinding alternative



- Blinding is a means, not an end. The end is to mitigate subjective bias.
- Instead, one can leave the data open, but **add fake signal events**. In this case, it might look like sprinkling salt on the plot.
- Other fields sometimes use a similar technique (often called “spiking”), e.g.
 - ▶ neutrino searches
 - ▶ searches for fractional charge
 - ▶ gravitational wave detection




Alternative




- Blinding is a means, not an end. The end is to mitigate subjective bias.
- Instead, one can leave the data open, but **add fake signal events**. In this case, it might look like sprinkling salt on the plot.
- Other fields sometimes use a similar technique (often called “spiking”), e.g.
 - ▶ neutrino searches
 - ▶ searches for fractional charge
 - ▶ gravitational wave detection

Blinding alternative






- Blinding is a means, not an end. The end is to mitigate subjective bias.
- Instead, one can leave the data open, but **add fake signal events**. In this case, it might look like sprinkling salt on the plot.
- Other fields sometimes use a similar technique (often called “spiking”), e.g.
 - ▶ neutrino searches
 - ▶ searches for fractional charge
 - ▶ gravitational wave detection

Salty details




Salty details



Salty details

At which stage
do we add salt?


Raw
waveforms
(digitized samples)



data
processing

Reduced
quantities
(used in analysis)


pulse areas, timing, widths, etc.



Salty details

At which stage
do we add salt?

Raw
waveforms
(digitized samples)




data
processing

Easy,
insufficient

Reduced
quantities
(used in analysis)

pulse areas, timing, widths, etc.




Salty details

At which stage
do we add salt?

Difficult,
necessary

Raw
waveforms
(digitized samples)




data
processing

Easy,
insufficient

Reduced
quantities
(used in analysis)

pulse areas, timing, widths, etc.





Salty details

At which stage
do we add salt?

Difficult,
necessary

Raw
waveforms
(digitized samples)




data
processing

Easy,
insufficient

Reduced
quantities
(used in analysis)

pulse areas, timing, widths, etc.



How to construct salted waveforms?

Three options for generating salt

- **Monte Carlo**

- ▶ Pro: One is flexible to choose pulse parameters
- ▶ Con: will probably miss some features of a real-world waveform, allowing identification of salt

- **DD Neutron calibration data**

- ▶ Pro: WIMP-like S2/S1 ratio
- ▶ Pro: actual detector data
- ▶ Con: Not super clean (may contain multiple scatters)
- ▶ Con: Events only exist at a fixed z position

- **Tritium calibration data**

- ▶ Pro: Clean events (single scatter, low-energy)
- ▶ Pro: actual detector data
- ▶ Pro: Spatially uniform (physically)
- ▶ Con: Wrong S2/S1 ratio

How to construct salted waveforms?

Three options for generating salt

- **Monte Carlo**

- ▶ Pro: One is flexible to choose pulse parameters
- ▶ Con: will probably miss some features of a real-world waveform, allowing identification of salt


- **DD Neutron calibration data**

- ▶ Pro: WIMP-like S2/S1 ratio
- ▶ Pro: actual detector data
- ▶ Con: Not super clean (may contain multiple scatters)
- ▶ Con: Events only exist at a fixed z position

- **Tritium calibration data**

- ▶ Pro: Clean events (single scatter, low-energy)
- ▶ Pro: actual detector data
- ▶ Pro: Spatially uniform (physically)
- ▶ Con: Wrong S2/S1 ratio


Turning tritium into salt




- By splicing together two ${}^3\text{H}$ events, we can create a chimera event whose S2-S1 proportion is consistent with a WIMP event.

Turning tritium into salt

- The LUX DAQ records waveforms of pulses only (no baseline recorded between)
 - simple to splice together bits from different pulses



Turning tritium into salt





- Generate targeted S1 and S2 pulse areas with MC.
- Collect sequestered, blind ${}^3\text{H}$ data set.
- Find a ${}^3\text{H}$ event with the targeted S2 size (Event 1)
- Find a ${}^3\text{H}$ event with the targeted S1 size (Event 2), from a similar position as Event 1.
- Rip out the S1 from Event 1, implant the S1 from Event 2.
- New event contains all the detector quirks / features from Event 1.

Salt development

Before implementing the salt in the real data stream, we created several sets of “known” salt, and presented these data to the data-analysis group in a series of “salt Turing tests” to see if analyzers could identify which events are real and which are salt.

Implementation



SaltDB


- Each generated salt waveform is given a random **TIME STAMP** and loaded into a database (“SaltDB”).
- SaltDB is locked; it injects salt waveforms into our data stream.
- A sysadmin (not on LUX) at one of our data centers has access to SaltDB. This person is “salt keeper”
- Un-salting: we give a timestamp list to the salt keeper, they identify which on the list are salt.

•
•

the results...


LUX's second science run (with salt)

LUX's second science data (332 live days, with salt)




- These are the data with salt still un-tagged.
- Salting was quite useful: it enabled us to identify a number of rare pathologies that contributed events to the signal band.

LUX's second science run (with salt)




- Salt events identified in blue.
- These lie in the signal band, as expected.

Scrutiny after the salt removal



- After un-salting, events outside the ER band were scrutinized again.
- Three events were identified away from the BG region that were found to be clearly pathological.


Scrutiny after the salt removal



- After un-salting, events outside the ER band were scrutinized again.
- Three events were identified away from the BG region that were found to be clearly pathological.

Pathological events

- Events **A** and **B** have ~80% of the light in a single top-edge PMT. Consistent with energy deposited outside the TPC, and light leaking through a gap near the edge of the PMT array. p of $O(10^{-9})$ and $O(10^{-10})$ for A and B, respectively.
- Event **C** is highly concentrated under a few top PMTs and has a time structure consistent with gas scintillation emission. Event came after high rate in preceding 1 second.
- Since these events do not correspond to interactions in the TPC, we develop cuts [post-un-salting] to target them.




Scrutiny after the salt removal

Two important points:

1. These events are clearly pathological: they do not correspond to interactions within the TPC.
2. These events have a very low background likelihood.




To be clear, these events COULD HAVE BEEN identified prior to un-salting (it was an oversight on our part).

As a collaboration, we decided these two facts warranted us to take steps to remove these pathologies.




Removal of new pathologies

- Post-un-salting cuts: (pre-existing cuts left untouched)
 - ▶ Loose cuts (high signal acceptance), defined on calibration data.
 - ▶ Flat signal acceptance.




Pathological events



- Red indicates events that are removed by these additional post-unsalting cuts. There are a few more that we missed, living in the background band.

Results



- Sets world's-best limits on spin-independent WIMP-nucleon scattering over wide range of masses!

PRL 118, 021303 (2017)

Summary


- Traditional blinding is problematic in a low-background, rare-event search.
- LUX implemented a method of adding fake “salt” signal events to the data stream.
- This method allowed a number of pathological events to be identified (which would have been missed in a traditional blind search). A few rare undetected pathologies were not caught before unsalting the data, but could have been.
- LUX has had an overall positive experience with this technique. Implementation details are very experiment specific.

Backup slides

Salt energy distribution


- Exponential + uniform distribution
- 4 free parameters:
- Similar to a low-mass and high-mass WIMP recoil spectrum.

$$\begin{aligned}\mu_E \\ \max(E_{\text{flat}}) \\ f_{\text{exp}} = A/(A + B) \\ N_{\text{tot}} = A + B\end{aligned}$$



- Exact values for these four distributions are chosen at [uniform] random, and are kept blind to analyzers.

Salt injection



- Each fake waveform is assigned a random timestamp and saved in a database (“saltDB”). The event builder queries the saltDB for any appropriate salt timestamps and adds them to raw1 if they do not conflict with the timestamps of existing. Our waveform viewer works on raw1 files.