

M4P57 Complex Manifolds

Lectured by Prof Paolo Cascini
Typed by David Kurniadi Angdinata

Spring 2020

Syllabus

Contents

1	Introduction	3
2	Local theory	5
2.1	Holomorphic functions in several variables	5
2.2	Cauchy formula in one variable	6
2.3	Rank theorem	7
2.4	Holomorphic differential forms	7
3	Complex manifolds	10
3.1	Complex manifolds	10
3.2	Holomorphic functions on complex manifolds	11
3.3	Complex submanifolds	12
3.4	Examples of complex manifolds	13
3.5	Tangent spaces of complex manifolds	14
3.6	Holomorphic differential forms on complex manifolds	15
4	Vector bundles	16
4.1	Holomorphic vector bundles	16
4.2	Examples of holomorphic vector bundles	17
4.3	Complexification of tangent bundles	19
4.4	Differential forms on complex tangent bundles	22
4.5	Dolbeault cohomology	23
4.6	Connections	24

1 Introduction

Lecture 1
Thursday
09/01/20

The following are references.

- O Biquard and A Höring, Kähler geometry and Hodge theory, 2008.
- J P Demailly, Complex analytic and differential geometry, 2012.
- C Voisin, Hodge theory and complex algebraic geometry, 2002.
- R O Wells, Differential analysis on complex manifolds, 1973.
- A Gathmann, Algebraic geometry, 2002
- P Griffiths and J Harris, Principles of algebraic geometry, 1978.

Complex manifolds are manifolds over \mathbb{C}^n .

Example 1.1. \mathbb{C}^1 is a complex manifold. Any open $U \subset \mathbb{C}^n$ is a complex manifold.

Example 1.2. The sphere $S^2 \subset \mathbb{R}^3$ is a complex manifold by

$$S^2 \cong \mathbb{C} \cup \{\infty\} = \mathbb{P}_{\mathbb{C}}^1 = \mathbb{CP}^1.$$

More in general $\mathbb{P}_{\mathbb{C}}^n$ is a complex manifold for all n .

Example 1.3. The torus

$$S^1 \times S^1 = \mathbb{R}^2 / \mathbb{Z}^2 = \mathbb{C} / \mathbb{Z}^2$$

is a complex manifold. More in general a $2n$ -dimensional torus \mathbb{C}^n / Λ for a lattice $\Lambda \cong \mathbb{Z}^{2n}$ is a complex manifold.

Example 1.4. Compact Riemannian surfaces of genus $g > 1$, called **hyperbolics**, are all complex manifolds.

Example 1.5. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be holomorphic. The graph of f ,

$$\Gamma_f = \{(x, f(x)) \mid x \in \mathbb{C}\} \subset \mathbb{C} \times \mathbb{C},$$

is a complex manifold. From Γ_f we can recover f , by

$$f(x) = q(p^{-1}(x) \cap \Gamma_f),$$

where $p, q : \mathbb{C}^2 \rightarrow \mathbb{C}$ are the projections to the first and second factors. This allows us to define f^{-1} . Assume f is bijective. Define

$$\tau : \begin{array}{ccc} \mathbb{C}^2 & \longrightarrow & \mathbb{C}^2 \\ (x, y) & \longmapsto & (y, x) \end{array}.$$

Define

$$\Gamma_{f^{-1}} = \tau(\Gamma_f).$$

Then f^{-1} is the function induced by $\Gamma_{f^{-1}}$. This makes sense even if f is not bijective. Then we get a multivalued function, such as $\log z$ as the inverse of $\exp z$.

Example 1.6. Generalising Example 1.5, we can consider two complex manifolds M and N and we can consider holomorphisms $f : M \rightarrow N$. Given M ,

$$\text{Aut } M = \{f : M \rightarrow M \text{ holomorphic bijective and } f^{-1} \text{ holomorphic}\}.$$

If $M = \mathbb{C}$, there are lots of C^∞ -functions $\mathbb{C} \rightarrow \mathbb{C}$ but the automorphisms of \mathbb{C} are just affine linear maps. If $M = \mathbb{C}/\mathbb{Z}^2$, then $\text{Aut } M$ is interesting.

Example 1.7. Algebraic geometry is the zeroes of polynomials. That is, fix m , and take polynomials f_1, \dots, f_k in m variables. Define

$$M = \{(x_1, \dots, x_m) \in \mathbb{C}^m \mid f_1(x_1, \dots, x_m) = \dots = f_k(x_1, \dots, x_m) = 0\}.$$

Then M is called an **algebraic variety**. If M is smooth then M is a complex manifold. Fix m , take homogeneous polynomials F_1, \dots, F_k in $m + 1$ variables, where F is **homogeneous** if it is the sum of monomials of the same degree. Consider

$$N = \{(x_0, \dots, x_m) \in \mathbb{P}_{\mathbb{C}}^m \mid F_1(x_0, \dots, x_m) = \dots = F_k(x_0, \dots, x_m) = 0\}.$$

Then N is called a **projective variety**. If N is smooth then N is a complex manifold.

The idea is if M is a differentiable manifold, then M contains lots of submanifolds N . This is not true for complex manifolds. There exist complex manifolds without any proper complex submanifolds, which is not true for projective varieties. The following are questions.

- What can we say about the topology of complex manifolds? For example, what is $\pi_1(M)$? What is the cohomology of M ?
- Assume that M and N are complex manifolds which are diffeomorphic. Are they also isomorphic, so there exists a biholomorphism $M \rightarrow N$?

What is next?

- Hodge decomposition theorem. Understand the cohomology of M by using the complex structure.
- Kodaira embedding theorem. Understand when a compact complex manifold is projective.

Note. If $M \subset \mathbb{P}_{\mathbb{C}}^m$ is a compact complex manifold then M is projective.

Example. Let $M = \Gamma_{\exp}$ for $\exp : \mathbb{C} \rightarrow \mathbb{C}$. Then $M \subset \mathbb{C}^2$ but it is not algebraic.

2 Local theory

2.1 Holomorphic functions in several variables

Notation 2.1. Given $z_0 \in \mathbb{C}$ and $r > 0$, the **disc** is

$$D(z_0, r) = \{z \in \mathbb{C} \mid |z - z_0| < r\},$$

and $\partial D(z_0, r)$ is the boundary of $D(z_0, r)$.

Definition 2.2. Let $U \subset \mathbb{C}$, and let $f : U \rightarrow \mathbb{C}$ be a function. Then f is **holomorphic at** $z_0 \in U$ if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists.

Theorem 2.3 (Cauchy). *Let $U \subset \mathbb{C}$ be open, let f be holomorphic on U , and let $z_0 \in U$. Assume that if $D = D(z_0, r) \subset U$ then $\overline{D} \subset U$. Then*

$$f(z_0) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(z)}{z - z_0} dz.$$

Notation 2.4. Fix $z_0 = (z_{01}, \dots, z_{0n}) \in \mathbb{C}^n$ and $R = (r_1, \dots, r_n) \in \mathbb{R}_{>0}^n$. Then the **polydisc** is

$$D(z_0, R) = \{z = (z_1, \dots, z_n) \in \mathbb{C}^n \mid |z_i - z_{0i}| < r_i \text{ for each } i\},$$

where R is the **polyradius**.

Definition 2.5. Let $U \subset \mathbb{C}^n$ be open, let $f : U \rightarrow \mathbb{C}$ be a continuous function, and let $z = (z_1, \dots, z_n) \in \mathbb{C}^n$. Then f is **holomorphic at** z , if assuming that $D = D(z, R) \subset U$ for some $R = (r_1, \dots, r_n)$ then

$$f(z_1, \dots, z_{i-1}, \cdot, z_{i+1}, \dots, z_n) : D(z_i, r_i) \rightarrow \mathbb{C}$$

is holomorphic for all i .

Example 2.6. Any convergent power series in n -variables is holomorphic.

The opposite is also true.

Theorem 2.7 (Cauchy). *Let $U \subset \mathbb{C}^n$ be an open set, let $f : U \rightarrow \mathbb{C}$ be holomorphic, and let $z = (z_1, \dots, z_n) \in U$. Assume that if $D = D(z, R)$ for some $R = (r_1, \dots, r_n)$ then $\overline{D} \subset U$. If $z' = (z'_1, \dots, z'_n) \in D$ then*

$$f(z') = \frac{1}{(2\pi i)^n} \int_{\partial D(z_1, r_1)} \cdots \int_{\partial D(z_n, r_n)} \frac{f(z)}{(z - z'_1) \cdots (z - z'_n)} dz_n \cdots dz_1.$$

Proof. Use induction on n and Cauchy theorem at each step. □

Corollary 2.8. *Let $U \subset \mathbb{C}^n$ be open, let $f : U \rightarrow \mathbb{C}$ be holomorphic, and let $z = (z_1, \dots, z_n) \in U$. Then there exists $D = D(z, R) \subset U$ for some $R = (r_1, \dots, r_n)$ and there exists*

$$p(w) = \sum_{m_1, \dots, m_n \geq 0} a_{m_1, \dots, m_n} (w_1 - z_1)^{m_1} \cdots (w_n - z_n)^{m_n},$$

such that p is convergent on D and $f(w) = p(w)$ inside D .

Proof. The idea is to use Theorem 2.7 and $1/(1-w) = \sum_{k \geq 0} w^k$. □

Definition 2.9. Let $U \subset \mathbb{C}^n$ be open. Then $f : U \rightarrow \mathbb{C}^m$ is **holomorphic** if $f_i = p_i \circ f$ is holomorphic for any $i = 1, \dots, m$ where $p_i : \mathbb{C}^m \rightarrow \mathbb{C}$ is the i -th projection, so $f = (f_1, \dots, f_m)$.

Fact. If $f : U \rightarrow \mathbb{C}^m$ is holomorphic and $g : V \rightarrow \mathbb{C}^p$ is holomorphic where $V \supset f(U)$ then $g \circ f$ is holomorphic.

Definition 2.10. Let $U \subset \mathbb{C}^n$ be open. A holomorphic function $f : U \rightarrow \mathbb{C}^m$ is **biholomorphic at** $z_0 \in U$ if there exists an open neighbourhood $V \subset U$ of z_0 such that $f : V \rightarrow f(V)$ is bijective and $f^{-1} : f(V) \rightarrow V$ is holomorphic. Then f is **biholomorphic** if f is bijective and f is biholomorphic at any point.

Note. $f(V)$ is automatically open in \mathbb{C}^m if $m = n$.

Example 2.11. Let $\Phi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be linear such that $\det \Phi \neq 0$. Then Φ is a biholomorphism.

Example 2.12. Let $U = \mathbb{C} \setminus \{0\}$ and

$$\begin{array}{ccc} f & : & U \longrightarrow U \\ z & \longmapsto & z^2 \end{array}.$$

Check that f is biholomorphic at any point of U but f is not biholomorphic.

Remark. $\mathbb{C}^n \cong \mathbb{R}^{2n}$ and $\mathbb{C}^m \cong \mathbb{R}^{2m}$. Then a holomorphic $f : U \subset \mathbb{C}^n \rightarrow \mathbb{C}^m$ is also a diffeomorphism $U \subset \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2m}$.

Theorem 2.13 (Hartogs). *Let $n \geq 2$, let $\epsilon = (\epsilon_1, \dots, \epsilon_n)$ and $\delta = (\delta_1, \dots, \delta_n)$ such that $\epsilon_i > \delta_i > 0$, let $U = D(0, \epsilon) \setminus \overline{D(0, \delta)}$, and let $f : U \rightarrow \mathbb{C}^m$ be holomorphic. Then there exists a holomorphic $\bar{f} : D(0, \epsilon) \rightarrow \mathbb{C}^m$ such that $\bar{f}|_U = f$.*

Example. Hartogs theorem is false for $n = 1$. If $f(z) = 1/z$, for all $\epsilon > \delta > 0$, then f cannot be extended.

2.2 Cauchy formula in one variable

Let $\omega = x + iy \in \mathbb{C}$ for $x, y \in \mathbb{R}$, and let $f : U \rightarrow \mathbb{C}$ be C^∞ for some $U \subset \mathbb{C}$. Recall that

$$\frac{\partial f}{\partial \omega} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) f, \quad \frac{\partial f}{\partial \bar{\omega}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) f.$$

Recall that f is holomorphic if and only if $\frac{\partial f}{\partial \bar{\omega}} = 0$ on U . More in general, let $U \subset \mathbb{C}^n$ be open, let $z_i = x_i + iy_i$, and let $f : U \rightarrow \mathbb{C}$ be a C^∞ -function. Then f is holomorphic if and only if $\frac{\partial f}{\partial \bar{z}_i} = 0$ for all $i = 1, \dots, n$. Let $\omega \in \mathbb{C}$. Since $dx \wedge dy = -dy \wedge dx$, let

$$dA = \frac{i}{2} d\omega \wedge d\bar{\omega} = \frac{i}{2} (dx + idy) \wedge (dx - idy) = dx \wedge dy,$$

which is the Lebesgue measure on $\mathbb{R}^2 \cong \mathbb{C}$.

Proposition 2.14. *Let $f : U \rightarrow \mathbb{C}$ for $U \subset \mathbb{C}$ be a C^∞ -function, and let $D = D(z, r)$ such that $\bar{D} \subset U$. Then*

$$f(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f}{\omega - z} d\omega - \frac{1}{\pi} \int_D \frac{1}{\omega - z} \frac{\partial f}{\partial \bar{\omega}} dA.$$

Proof. Assume $z = 0$. Recall that $f(\omega) = 1/\omega$ is locally integrable around zero, so

$$\int_D \frac{1}{\omega} \frac{\partial f}{\partial \bar{\omega}} dA = \lim_{\epsilon \rightarrow 0} \int_{D \setminus D(0, \epsilon)} \frac{1}{\omega} \frac{\partial f}{\partial \bar{\omega}} dA.$$

Away from zero

$$\begin{aligned} d \left(\frac{f}{\omega} d\omega \right) &= \frac{1}{\omega} df \wedge d\omega + f d \left(\frac{1}{\omega} \right) \wedge d\omega = \frac{1}{\omega} \left(\frac{\partial f}{\partial \omega} d\omega + \frac{\partial f}{\partial \bar{\omega}} d\bar{\omega} \right) \wedge d\omega + f \frac{\partial}{\partial \omega} \left(\frac{1}{\omega} \right) d\omega \wedge d\omega \\ &= \frac{1}{\omega} \frac{\partial f}{\partial \bar{\omega}} d\bar{\omega} \wedge d\omega = \frac{2i}{\omega} \frac{\partial f}{\partial \bar{\omega}} dA. \end{aligned}$$

Then

$$\begin{aligned} \frac{1}{\pi} \int_D \frac{1}{\omega} \frac{\partial f}{\partial \bar{\omega}} dA &= \frac{1}{\pi} \lim_{\epsilon \rightarrow 0} \int_{D \setminus D(0, \epsilon)} \frac{1}{\omega} \frac{\partial f}{\partial \bar{\omega}} dA \\ &= \frac{1}{2\pi i} \lim_{\epsilon \rightarrow 0} \int_{D \setminus D(0, \epsilon)} d \left(\frac{f}{\omega} d\omega \right) & \frac{1}{\omega} \frac{\partial f}{\partial \bar{\omega}} dA &= \frac{1}{2i} d \left(\frac{f}{\omega} d\omega \right) \\ &= \frac{1}{2\pi i} \lim_{\epsilon \rightarrow 0} \left(\int_{\partial D} \frac{f}{\omega} d\omega - \int_{\partial D(0, \epsilon)} \frac{f}{\omega} d\omega \right) & \text{Stokes' theorem} \\ &= \frac{1}{2\pi i} \left(\int_{\partial D} \frac{f}{\omega} d\omega - 2\pi i f(0) \right) & \lim_{\epsilon \rightarrow 0} \int_{\partial D(0, \epsilon)} \frac{1}{\omega} d\omega &= 2\pi i. \end{aligned}$$

□

If f is holomorphic, then $\frac{\partial f}{\partial \bar{\omega}} = 0$, which implies Theorem 2.3.

2.3 Rank theorem

Let $U \subset \mathbb{C}^n$ be open, and let $f : U \rightarrow \mathbb{C}^m$ be holomorphic. Then the **Jacobian** is

$$J_f = \left(\frac{\partial f_j}{\partial z_i}(z) \right),$$

where $f_j = p_j \circ f$ and $p_j : \mathbb{C}^m \rightarrow \mathbb{C}$ is the j -th projection.

Exercise. Show that the real Jacobian, which is $2n \times 2n$, has non-negative determinants.

Theorem 2.15 (Rank theorem). *Let $z \in U$ such that $r = \text{rk } J_f(z')$ is constant around z . Then there exist open $z \in V \subset U \subset \mathbb{C}^n$ and $f(z) \in W \subset \mathbb{C}^m$ such that $\phi : D(0, 1)^n \rightarrow V$ and $\psi : D(0, 1)^m \rightarrow W$ are biholomorphisms such that*

$$\eta = \psi^{-1} \circ f \circ \phi : \begin{array}{ccc} D(0, 1)^n & \longrightarrow & D(0, 1)^m \\ (z_1, \dots, z_n) & \longmapsto & (z_1, \dots, z_r, 0, \dots, 0) \end{array},$$

so

$$\begin{array}{ccccc} \mathbb{C}^n \supset U & \supset & V \ni z & \xrightarrow{f} & f(z) \in W \subset f(U) \subset \mathbb{C}^m \\ & & \uparrow \phi & & \uparrow \psi \\ & & D(0, 1)^n & \xrightarrow{\eta} & D(0, 1)^m \end{array}.$$

Corollary 2.16 (Inverse function theorem). *Let $f : U \rightarrow \mathbb{C}^n$ be holomorphic for $U \subset \mathbb{C}^n$, and let $z \in U$ such that $\det J_f(z) \neq 0$. Then f is a biholomorphism at z .*

Proof. $\det J_f(z) \neq 0$ if and only if $\text{rk } J_f(z) = n$, so $\text{rk } J_f(z') = n$ around z , since $\det J_f(z)$ is a continuous function. Let ϕ and ψ as in the theorem. Then $\eta = \psi^{-1} \circ f \circ \phi = \text{id}$, so on V , $f = \psi \circ \phi^{-1}$ is a composition of biholomorphisms, which is a biholomorphism. \square

Remark 2.17. Let $f : U \rightarrow \mathbb{C}^n$ for $U \subset \mathbb{C}^n$. Then $\det J_f(z)$ is a holomorphism, so

$$Z = \{z \in U \mid \det J_f(z) = 0\}$$

is closed.

2.4 Holomorphic differential forms

Let $U \subset \mathbb{C}^n$ be open.

Definition 2.18. A **holomorphic vector field** on U is the expression

$$X = \sum_i a_i \frac{\partial}{\partial z_i},$$

where a_i are holomorphic functions on U .

For all $x \in U$, the **tangent space** is

$$T_x U = \left\langle \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right\rangle \cong \mathbb{C}^n.$$

If $x \in U$, then $X(x) \in T_x U$.

Notation 2.19.

$$H^0(U, \mathcal{O}_U) = \{\text{holomorphic functions } f : U \rightarrow \mathbb{C}\}, \quad H^0(U, T_U) = \{\text{holomorphic vector fields on } U\}.$$

Remark. $R = H^0(U, \mathcal{O}_U)$ is a ring and $M = H^0(U, T_U)$ is a module over R . That is, if $X \in H^0(U, T_U)$ and $f \in H^0(U, \mathcal{O}_U)$, then $fX \in H^0(U, T_U)$.

Definition 2.20. Let R be a ring and M be an R -module for $p \geq 1$. The p -th exterior power $\Lambda^p M$ of M is the R -module $M^{\otimes p}$ with the relations

$$m_1 \otimes \cdots \otimes m_p - \epsilon(\sigma) m_{\sigma(1)} \otimes \cdots \otimes m_{\sigma(p)}, \quad m_1, \dots, m_p \in M, \quad \sigma \in \mathcal{S}_p,$$

where $\epsilon(\sigma) = (-1)^m$ is the signature of σ and m is the number of transpositions defining σ . Then $M^* = \text{Hom}_R(M, R)$ is the **dual** of M as an R -module.

Let $R = H^0(U, \mathcal{O}_U)$ and $M = H^0(U, T_U)$.

Definition 2.21. Let $p > 0$. We define a **holomorphic p -form**, as an element of

$$H^0(U, \Omega_U^p) = \Lambda^p M^*.$$

If $p = 0$, by convention a **holomorphic 0-form** is just an element in R .

Let (z_1, \dots, z_n) be coordinates for U . Recall $\eta \in M$ is given by $\eta = \sum_i a_i \frac{\partial}{\partial z_i}$ for holomorphic functions $a_i \in R$. Then $\omega \in M^*$ is given by the expression

$$\sum_i b_i dz_i, \quad b_i \in R, \quad dz_i \left(\frac{\partial}{\partial z_j} \right) = \delta_{ij}.$$

More in general $\omega \in H^0(U, \Omega_U^p)$ is given by

$$\omega = \sum_{|I|=p} f_I dz_{i_1} \wedge \cdots \wedge dz_{i_p}, \quad f_I \in R, \quad I = (i_1, \dots, i_p), \quad i_1 < \cdots < i_p,$$

where $dz_{i_1}, \dots, dz_{i_p}$ is an R -basis of $H^0(U, \Omega_U^p)$.

Example.

$$H^0(U, \Omega_U^p) \cong \Lambda^p H^0(U, \Omega_U^1)$$

is an isomorphism as R -modules. This is not true for complex manifolds in general.

The **exterior product** is

$$\begin{aligned} H^0(U, \Omega_U^p) \otimes H^0(U, \Omega_U^q) &\longrightarrow H^0(U, \Omega_U^{p+q}) \\ \omega_1 \otimes \omega_2 &\longmapsto \omega_1 \wedge \omega_2 \end{aligned},$$

where we just need to define

$$\omega_1 \wedge \omega_2 = f dz_{i_1} \wedge \cdots \wedge dz_{i_p} \otimes g dz_{j_1} \wedge \cdots \wedge dz_{j_q} = fg dz_{i_1} \wedge \cdots \wedge dz_{i_p} \wedge dz_{j_1} \wedge \cdots \wedge dz_{j_q},$$

by linearity. Then $\omega_1 \wedge \omega_2 = 0$ if $\{i_1, \dots, i_p\} \cap \{j_1, \dots, j_q\} \neq \emptyset$, since $dz_i \wedge dz_i = 0$.

Exercise. Check that this definition coincides with the definition in M4P54.

The **exterior derivative** is

$$\begin{aligned} d : H^0(U, \Omega_U^p) &\longrightarrow H^0(U, \Omega_U^{p+1}) \\ f dz_{i_1} \wedge \cdots \wedge dz_{i_p} &\longmapsto \sum_{j=1}^n \frac{\partial f}{\partial z_j} dz_j \wedge dz_{i_1} \wedge \cdots \wedge dz_{i_p}. \end{aligned}$$

By definition d is \mathbb{C} -linear, but not R -linear. That is,

$$d(a\omega_1 + b\omega_2) = ad\omega_1 + bd\omega_2, \quad \omega_1, \omega_2 \in H^0(U, \Omega_U^p), \quad a, b \in \mathbb{C}.$$

Proposition 2.22. Let $U \subset \mathbb{C}^n$ be open. Then

- the Leibnitz rule

$$d(\omega_1 \wedge \omega_2) = d\omega_1 \wedge \omega_2 + (-1)^p \omega_1 \wedge d\omega_2, \quad \omega_1 \in H^0(U, \Omega_U^p), \quad \omega_2 \in H^0(U, \Omega_U^q),$$

- $d^2 = 0$, that is

$$d(d\omega) = 0, \quad \omega \in H^0(U, \Omega_U^p).$$

Lecture 4
Thursday
16/01/20

Definition 2.23. Let $f : U \subset \mathbb{C}^n \rightarrow \mathbb{C}^m$ be holomorphic, let $f_i = p_i \circ f : U \rightarrow \mathbb{C}$ where $p_i : \mathbb{C}^m \rightarrow \mathbb{C}$ is the i -th projection, and let $f(U) \subset V \subset \mathbb{C}^m$ be open. Then if

$$\omega = h dz_{i_1} \wedge \cdots \wedge dz_{i_p} \in H^0(V, \Omega_V^p), \quad h \in H^0(U, \mathcal{O}_U),$$

then we can define the **pull-back** of ω ,

$$f^*\omega = h \circ f df_{i_1} \wedge \cdots \wedge df_{i_p} \in H^0(U, \Omega_U^p),$$

since $f_i \in H^0(V, \mathcal{O}_V) = H^0(V, \Omega_V^0)$ implies that $df_i \in H^0(V, \Omega_V^1)$, so

$$\begin{array}{ccc} U & \xrightarrow{f} & f(U) \subset V \\ & \searrow h \circ f \in H^0(U, \mathcal{O}_U) & \downarrow h \\ & & \mathbb{C} \end{array} .$$

This is linear over \mathbb{C} and over $H^0(U, \mathcal{O}_U)$.

Proposition 2.24. Let $U \subset \mathbb{C}^n$, $V \subset \mathbb{C}^m$, and $W \subset \mathbb{C}^{m'}$ be open, let $f : U \rightarrow \mathbb{C}^m$ and $g : V \rightarrow \mathbb{C}^{m'}$ be holomorphic such that $V \supset f(U)$ and $W \supset g(V)$, and let $\omega \in H^0(V, \Omega_V^p)$ and $\eta \in H^0(W, \Omega_W^q)$. Then

- $f^*(\omega + \eta) = f^*\omega + f^*\eta$ if $p = q$,
- $f^*(\omega \wedge \eta) = f^*\omega \wedge f^*\eta$,
- $df^*\omega = f^*d\omega$, and
- $f^*g^*\omega = (g \circ f)^*\omega$.

Let $U \subset \mathbb{C}^n \cong \mathbb{R}^{2n}$, and let $z_i = x_i + iy_i$ for $i = 1, \dots, n$ and $x_i, y_i \in \mathbb{R}$. Then

$$dz_i = dx_i + idy_i,$$

so any holomorphic form is a differentiable form on \mathbb{R}^{2n} . A (p, q) -**form** is a differentiable $(p + q)$ -form of the expression

$$\omega = \sum_{|I|=p, |J|=q} f_{I,J} dz_{i_1} \wedge \cdots \wedge dz_{i_p} \wedge d\bar{z}_{j_1} \wedge \cdots \wedge d\bar{z}_{j_q}, \quad f_{I,J} : U \rightarrow \mathbb{C} \cong \mathbb{R}^2 \in C^\infty,$$

where $d\bar{z}_j = dx_j - idy_j$. We denote

$$C^\infty(U, \Omega_U^{p,q}) = \{\text{differentiable } (p + q)\text{-forms on } U\}.$$

If ω is a (p, q) -form, then the **conjugate** $\bar{\omega}$ of ω is the (q, p) -form defined by

$$\bar{\omega} = \sum_{|I|=p, |J|=q} \overline{f_{I,J}} d\bar{z}_{i_1} \wedge \cdots \wedge d\bar{z}_{i_p} \wedge dz_{j_1} \wedge \cdots \wedge dz_{j_q}.$$

3 Complex manifolds

3.1 Complex manifolds

Definition 3.1. A **complex manifold** of dimension n is a connected Hausdorff topological space X , with a countable open cover $\{U_\alpha\}$ of X such that for all α , there exists $\phi_\alpha : U_\alpha \rightarrow \mathbb{C}^n$ such that $\phi_\alpha : U_\alpha \rightarrow \phi_\alpha(U_\alpha)$ is a homeomorphism and

$$\phi_\alpha \circ \phi_\beta^{-1} : \phi_\beta(U_\alpha \cap U_\beta) \rightarrow \phi_\alpha(U_\alpha \cap U_\beta)$$

is a biholomorphism for each α and β , so

$$\begin{array}{ccc} & U_\alpha \cap U_\beta & \\ \phi_\alpha \swarrow & & \searrow \phi_\beta \\ \mathbb{C}^n \supset \phi_\alpha(U_\alpha \cap U_\beta) & \xrightarrow{\phi_\alpha \circ \phi_\beta^{-1}} & \phi_\beta(U_\alpha \cap U_\beta) \subset \mathbb{C}^n \end{array} .$$

The pair (U_α, ϕ_α) is called a **holomorphic chart**. The set $\{(U_\alpha, \phi_\alpha)\}$ is called a **holomorphic atlas** or a **complex structure**.

Recall X is Hausdorff if for all $x, y \in X$ there exist U and V open in X such that $U \cap V = \emptyset$ and $x \in U$ and $y \in V$.

Example 3.2.

- If $U \subset \mathbb{C}^n$ is an open set then U is a complex manifold. More in general if X is a complex manifold and $U \subset X$ is open then U is a complex manifold. Let $\{(U_\alpha, \phi_\alpha)\}$ be a complex structure on M . Then

$$\{(\overline{U_\alpha}, \overline{\phi_\alpha})\} = \{(U_\alpha \cap U, \phi_\alpha|_{\overline{U_\alpha}})\}$$

is a complex structure of M .

- If X and Y are complex manifolds, then $X \times Y$ is a complex manifold.

Example 3.3. The projective space $\mathbb{P}_{\mathbb{C}}^n$ or \mathbb{CP}^n . Let $V^* = \mathbb{C}^{n+1} \setminus \{0\}$, with coordinates (z_0, \dots, z_n) . Define an equivalence on V^* as

$$v_1 \sim v_2 \iff \exists \lambda \in \mathbb{C}, v_1 = \lambda v_2.$$

Check that \sim is an equivalence. Consider the Euclidean topology on V^* . Then there exists an induced topology on $X = V^*/\sim = \{[v] \mid v \in V^*\}$, with quotient map

$$\begin{array}{ccc} q & : & V^* \longrightarrow X \\ & & v \longmapsto [v] \end{array} .$$

Given $v = (z_0, \dots, z_n) \in V^*$ we denote $[v] = [z_0, \dots, z_n]$ such that $z_i \neq 0$ for some i . Two elements $[x_0, \dots, x_n]$ and $[y_0, \dots, y_n]$ of X define the same point if and only if there exists λ such that $x_i = \lambda y_i$ for all i . Let

$$V_i = \{(z_0, \dots, z_n) \in V^* \mid z_i \neq 0\},$$

which is open in V^* , and let $U_i = q(V_i)$, which is open in X , such that $\{U_i\}$ is a cover of X , that is $\bigcup_i U_i = X$. Let

$$H_i = \{(z_0, \dots, z_n) \in V^* \mid z_i = 1\}.$$

Then there exists a homeomorphism

$$\begin{array}{ccc} r_i & : & H_i \longrightarrow \mathbb{C}^n \\ & & (z_0, \dots, z_n) \longmapsto [z_0, \dots, z_{i-1}, z_{i+1}, \dots, z_n] \end{array} ,$$

and let

$$\begin{array}{ccc} q_i = q|_{H_i} & : & H_i \subset V^* \longrightarrow U_i \subset X \\ & & (z_0, \dots, z_n) \longmapsto [z_0, \dots, z_n] \end{array}$$

be also a homeomorphism.

Lecture 5
Thursday
16/01/20

- q_i is surjective. Take $[x_0, \dots, x_n] \in U_i$. Then $x_i \neq 0$, so choose $\lambda = 1/x_i$. Then

$$[x_0, \dots, x_n] = \left[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i} \right] = q(z_0, \dots, z_n), \quad z_j = \frac{x_j}{x_i},$$

and in particular $z_i = 1$, so there exists $(z_0, \dots, z_n) \in H_i$ such that $q_i(z_0, \dots, z_n) = [x_0, \dots, x_n]$.

- q_i is injective.¹

For all i , $q_i^{-1} : U_i \rightarrow H_i$ and $r_i : H_i \rightarrow \mathbb{C}^n$ are homeomorphisms, so $\phi_i = r_i \circ q_i^{-1} : U_i \rightarrow \mathbb{C}^n$ is also a homeomorphism. We want to show that (U_i, ϕ_i) define a holomorphic atlas, so

$$\phi_i \circ \phi_j^{-1} : \phi_j(U_i \cap U_j) \rightarrow \phi_i(U_i \cap U_j)$$

is a biholomorphism. Consider the case $j = 0$ and $i = 1$. Then $\phi_0(U_0 \cap U_1) = \{(x_1, \dots, x_n) \mid x_1 \neq 0\}$, so

$$\begin{aligned} \phi_1 \circ \phi_0^{-1} : \phi_0(U_0 \cap U_1) &\longrightarrow \phi_1(U_0 \cap U_1) \\ (x_1, \dots, x_n) &\longmapsto \left(1, \frac{x_2}{x_1}, \dots, \frac{x_n}{x_1}\right) \end{aligned}$$

is a biholomorphism. Thus X is a compact complex manifold. If $n = 1$, then $\mathbb{P}_{\mathbb{C}}^1 \cong \mathbb{S}^2$.

Example 3.4. The complex torus. Let

$$\begin{aligned} \Lambda = \mathbb{Z}^{2n} &\longrightarrow \mathbb{C}^n \\ (a_1, \dots, a_n, b_1, \dots, b_n) &\longmapsto (a_1 + ib_1, \dots, a_n + ib_n) \end{aligned}$$

Define an equivalence on \mathbb{C}^n by

$$v_1 \sim v_2 \iff v_1 - v_2 \in \Lambda.$$

Then $X = \mathbb{C}^n / \sim$ with quotient map $q : \mathbb{C}^n \rightarrow X$ is Hausdorff and compact. Topologically $X \cong [0, 1]^{2n} / \sim$. For each $x \in \mathbb{C}^n$, we can find an open set $x \in U \subset \mathbb{C}^n$ such that $q|_U : U \rightarrow X$ is a homeomorphism. The idea is if $x \in (0, 1)^{2n}$ then we can take $U = (0, 1)^{2n}$. If not, there exists a translation of $\mathbb{C}^n \rightarrow \mathbb{C}^n$ such that the property holds. We define

$$\phi_V = q|_U^{-1} : V \subset \mathbb{C}^n / \Lambda \rightarrow U \subset \mathbb{C}^n, \quad V = q(U).$$

Show that (V, ϕ_V) define a complex structure on X .² This is also a compact complex manifold. More in general \mathbb{C}^n / Λ where $\Lambda \cong \mathbb{Z}^{2n}$ is a lattice is a compact complex manifold.

3.2 Holomorphic functions on complex manifolds

Definition 3.5. Let $f : X \rightarrow Y$ be a continuous morphism between complex manifolds. Then f is **holomorphic** if there exists a complex structure $\{(U_\alpha, \phi_\alpha)\}$ on Y and for all $y \in Y$ there exists a holomorphic chart (V_α, ψ_α) such that $x \in V_\alpha$ and $f(V_\alpha) \subset U_\alpha$ around any point x of $f^{-1}(y)$ and $\phi_\alpha \circ f \circ \psi_\alpha^{-1}$ is holomorphic, so

$$\begin{array}{ccc} X \supset V_\alpha & \xrightarrow{f} & U_\alpha \subset Y \\ \psi_\alpha \downarrow & & \downarrow \phi_\alpha \\ \psi_\alpha(V_\alpha) & \xrightarrow{\tilde{f}} & \phi_\alpha(U_\alpha) \end{array}$$

Then $J_f = J_{\tilde{f}}$, and a **holomorphic function on X** is a holomorphic function $f : X \rightarrow \mathbb{C}$.

Exercise 3.6. If X is a compact complex manifold then any holomorphic function $f : X \rightarrow \mathbb{C}$ is constant.

¹Exercise

²Exercise

Definition 3.7. Let $f : X \rightarrow Y$ be a holomorphic function between complex manifolds. Then f is

- a **submersion** if $\dim Y \geq \dim X = r$ and $\text{rk } J_f = r$ at any point,
- an **immersion** if $r = \dim X \leq \dim Y$ and $\text{rk } J_f = r$ at any point, and
- an **embedding** if it is an immersion and $f : X \rightarrow f(X)$ is a homeomorphism.

Example 3.8. Let $f_2, \dots, f_n : \mathbb{C} \rightarrow \mathbb{C}$ be holomorphic, and let

$$\begin{aligned} f &: \mathbb{C} \longrightarrow \mathbb{C}^n \\ z &\longmapsto (z, f_2(z), \dots, f_n(z)) \end{aligned}$$

Then f is an embedding.

Example 3.9. Let $X = \mathbb{C}^2 / \Lambda$ for $\Lambda = \mathbb{Z}^4 \subset \mathbb{C}^2$, and let $q : \mathbb{C}^2 \rightarrow X$. Fix $\lambda \in \mathbb{C}$. Let

$$\begin{aligned} f &: \mathbb{C} \longrightarrow \mathbb{C}^2 \\ z &\longmapsto (z, \lambda z) \end{aligned}$$

Then $\tilde{f} = q \circ f : \mathbb{C} \rightarrow X$ is an immersion.

- If $\lambda = 0$ or $\lambda = \frac{1}{2}$, then $\tilde{f}(\mathbb{C})$ is a closed submanifold.
- If λ is general then $\tilde{f}(\mathbb{C})$ is dense inside X , so it is not closed. Thus it is not a complex submanifold of X .

3.3 Complex submanifolds

Definition 3.10. Let $i : X \rightarrow Y$ be an embedding of complex manifolds. If $i(X) \subset Y$ is closed then $i(X)$ is called a **complex submanifold** of Y . The **codimension** of X in Y is $\dim Y - \dim X$.

Theorem 3.11.

1. Let $i : X \rightarrow Y$ be a submanifold of codimension k , and let $n = \dim X$. Then for all $x \in X$, there exists an open neighbourhood $x \in U \subset Y$ and a submersion $f : U \rightarrow D(0, 1)^k \subset \mathbb{C}^k$ such that $X \cap U = f^{-1}(0)$.
2. If $X \subset Y$ is a closed subset such that for all $x \in X$ there exists $U \ni x$ open in Y and a submersion $f : U \rightarrow D(0, 1)^k$ such that $X \cap U = f^{-1}(0)$, then X is a complex submanifold.

Proof.

1. We can assume that if there exists a holomorphic chart (U, ψ) on Y such that $x \in U$ and if $V = i^{-1}(U)$ then there exists $\phi : V \rightarrow \mathbb{C}^n$ such that (V, ϕ) is a holomorphic chart on X . After possibly shrinking U smaller, by the rank theorem, there exist biholomorphic $a : \psi(U) \rightarrow D(0, 1)^{n+k}$ and $b : \phi(U) \rightarrow D(0, 1)^n$ such that the induced morphism is given by

$$\begin{aligned} D(0, 1)^n &\longrightarrow D(0, 1)^{n+k} \\ (z_1, \dots, z_n) &\longmapsto (z_1, \dots, z_n, 0, \dots, 0) \end{aligned}$$

Let

$$\begin{aligned} c &: D(0, 1)^{n+k} \longrightarrow D(0, 1)^k \\ (z_1, \dots, z_{n+k}) &\longmapsto (z_{n+1}, \dots, z_{n+k}) \end{aligned}$$

so

$$\begin{array}{ccccc} Y & \supset & U & \xrightarrow{\phi} & \phi(U) & \xrightarrow{b} & D(0, 1)^n \subset \mathbb{C}^n & \xleftarrow{c} & \\ \uparrow i & & \uparrow i & & \downarrow & & \downarrow & & \\ X & \supset & V & \xrightarrow{\psi} & \psi(U) & \xrightarrow{a} & D(0, 1)^{n+k} \subset \mathbb{C}^{n+k} & & \end{array}$$

Then f is the composition $c \circ a \circ \psi : U \rightarrow D(0, 1)^n$.

2. Let $\{(U_\alpha, \phi_\alpha)\}$ be a complex structure on Y , and let $V_\alpha = X \cap U_\alpha$ and $\psi_\alpha = \phi_\alpha|_{V_\alpha}$. The goal is to show that $\{(V_\alpha, \psi_\alpha)\}$ defines a complex structure on X . By assumption,

$$\phi_\alpha \circ \phi_\beta^{-1} : \phi_\beta(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n+k} \rightarrow \phi_\alpha(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n+k}$$

is biholomorphic. Let $U' = \phi_\beta(U)$, let $X' = \phi_\beta(X \cap U)$, and let $f' = f \circ \phi_\beta^{-1}$, so

$$\begin{array}{ccccccc} & & & & \phi_\alpha(U) & \subset & \phi_\alpha(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n+k} \\ & & & & \nearrow \phi_\alpha & & \uparrow \phi_\alpha \circ \phi_\beta^{-1} \\ Y & \supset & U_\alpha \cap U_\beta & \supset & U & \xrightarrow{\phi_\beta} & U' \subset \phi_\beta(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n+k} \\ \uparrow i & & \cup & & \cup & & \cup \\ X & \supset & X \cap U_\alpha \cap U_\beta & \supset & X \cap U & \xrightarrow{f} & X' \subset D(0,1)^k \subset \mathbb{C}^k \end{array}$$

Then $f'^{-1}(0) = \phi_\beta(X \cap U_\alpha \cap U_\beta)$ and f' is also a submersion. By the rank theorem, we may assume that $U' = D(0,1)^{n+k}$ and $f'(z_1, \dots, z_{n+k}) = (z_1, \dots, z_k)$, so $\phi_\beta(X' \cap U_\alpha \cap U_\beta) = f'^{-1}(0)$. Thus

$$(\psi_\alpha \circ \psi_\beta^{-1})(z_1, \dots, z_n) = (\phi_\alpha \circ \phi_\beta^{-1})(z_1, \dots, z_n, 0, \dots, 0)$$

is also a biholomorphism. □

3.4 Examples of complex manifolds

Example 3.12. Let $U \subset \mathbb{C}^n$ be open, let $k \leq n$, let $f_1, \dots, f_k : U \rightarrow \mathbb{C}$, and let

$$V = \{x \in \mathbb{C}^n \mid f_1(x) = \dots = f_k(x) = 0\}.$$

Assume that $\left(\frac{\partial f_i}{\partial z_j}\right)$ has maximal rank k at any point of U . Then V is a complex submanifold of U . The idea is if $f = (f_1, \dots, f_k) : U \rightarrow \mathbb{C}^k$, then f is a submersion around any point of V , and use the previous Theorem 3.11.

Example 3.13. Let $f : X \rightarrow Y$ be a holomorphism between complex manifolds, and let $W \subset X$ be a submanifold. Then $f|_W : W \rightarrow Y$ is holomorphic.

Exercise 3.14. Let $X = \mathbb{C}^n$. Show that all the compact submanifolds of X are zero-dimensional, that is points.

Exercise 3.15. Let X and Y be compact manifolds. Recall that $X \times Y$ is also a complex manifold. Assume $f : X \rightarrow Y$, so

$$\Gamma_f = \{(x, f(x)) \mid x \in X\} \subset X \times Y.$$

Show that Γ_f is a complex submanifold.

Example 3.16. Let $n, m > 0$, and let

$$\text{Mat}_{n,m} \mathbb{C} = \{(n \times m)\text{-matrices}\} \cong \mathbb{C}^{n \cdot m}.$$

Then $\text{Mat}_{n,m} \mathbb{C}$ is a complex manifold. Let

$$\text{GL}_n \mathbb{C} = \{(n \times n)\text{-matrices } A \mid A \text{ invertible}\}.$$

Then $\text{GL}_n \mathbb{C}$ is a complex manifold, open in $\text{Mat}_{n,n} \mathbb{C}$.

Lecture 7
Thursday
23/01/20

Example 3.17. Projective manifolds. Let $R = \mathbb{C}[x_0, \dots, x_n]$ be the ring of polynomials, and let $X = \mathbb{P}_{\mathbb{C}}^n$ be the complex projective space. Then $f \in R$ is homogeneous of degree d if $f(\lambda x) = \lambda^d f(x)$. Let $q : \mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}_{\mathbb{C}}^n$, let F_1, \dots, F_k be homogeneous polynomials in R , and let

$$V = \{F_1 = \dots = F_k = 0\} \subset \mathbb{C}^{n+1} \setminus \{0\}, \quad W = q(V) \subset \mathbb{P}_{\mathbb{C}}^n,$$

so $q^{-1}(W) = V$, because F_i are homogeneous. Since V is closed in $\mathbb{C}^{n+1} \setminus \{0\}$, W is closed in $\mathbb{P}_{\mathbb{C}}^n$. Claim that if V is a submanifold of $\mathbb{C}^{n+1} \setminus \{0\}$ then W is a compact submanifold of $\mathbb{P}_{\mathbb{C}}^n$. If $\{U_i\}$ is the open covering given by

$$U_i = \{[x_0, \dots, x_n] \mid x_i \neq 0\},$$

then it is enough to show that $W \cap U_i$ is a complex submanifold of U_i for all i . Assume $i = n$. Let $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$. Then $q(x) = \mathbb{C}^*$ for all $x \in X$ but $\mathbb{P}_{\mathbb{C}}^n \times \mathbb{C}^* \neq \mathbb{C}^{n+1} \setminus \{0\}$. We want to show there exists a biholomorphism

$$\begin{aligned} \phi_n : \quad U_n \times \mathbb{C}^* &\longrightarrow q^{-1}(U_n) = \{(x_0, \dots, x_n) \in \mathbb{C}^{n+1} \mid x_n \neq 0\} \\ ([x_0, \dots, x_n], t) &\longmapsto \left(\frac{tx_0}{x_n}, \dots, \frac{tx_{n-1}}{x_n}, t \right), \end{aligned}$$

such that

$$\begin{aligned} \phi_n^{-1} : \quad q^{-1}(U_n) &\longrightarrow U_n \times \mathbb{C}^* \\ (y_0, \dots, y_n) &\longmapsto (q(y_0, \dots, y_n), y_n) = ([y_0, \dots, y_n], y_n). \end{aligned}$$

From this, it follows that $V \cap q^{-1}(U_n) \cong (W \cap U_n) \times \mathbb{C}^*$, so the claim follows.

Example 3.18. Plane curves. Let $X = \mathbb{P}_{\mathbb{C}}^2$, let $F \in R[x_0, x_1, x_2]$ be homogeneous of degree d , and let $W = \{F = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$. Then W is a compact complex submanifold if and only if for all $x \in \mathbb{P}_{\mathbb{C}}^2$, $\partial_{x_i} F(x) \neq 0$ for some i .

$d = 1$. W is the projective line, so $F = ax_0 + bx_1 + cx_2$ for a, b, c not all zero. Then W is a complex submanifold. There exists a biholomorphism $\mathbb{P}_{\mathbb{C}}^1 \rightarrow W$.

$d = 2$. W is a conic, so F is a degree two polynomial. Then $F = x_0x_1$ does not define a manifold. If $F = x_0x_1 - x_2^2$, then W is a complex submanifold of X . There exists

$$\begin{aligned} \mathbb{P}_{\mathbb{C}}^1 &\longrightarrow W \subset X \\ [t_0, t_1] &\longmapsto [t_0^2, t_1^2, t_0t_1]. \end{aligned}$$

Check that it is a biholomorphism. ³ This is true for any f of degree two such that W is a complex submanifold.

$d \geq 3$. If W is a complex submanifold then we will show that W is not biholomorphic to $\mathbb{P}_{\mathbb{C}}^1$.

3.5 Tangent spaces of complex manifolds

Definition 3.19. Let X be a complex manifold of dimension n , and let $x \in X$. Then there exists a chart (U, ϕ) around x such that $\phi(U) \subset \mathbb{C}^n$. The **holomorphic tangent space** $T_x X$ of X at x , is the vector space over \mathbb{C} generated by

$$\left(\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n} \right).$$

Let X be a real manifold. The **real tangent space** $T_x^{\mathbb{R}} X$ is the vector space over \mathbb{R} defined by

$$\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}, \frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial y_n} \right),$$

where $(x_1, \dots, x_n, y_1, \dots, y_n)$ are coordinates of \mathbb{R}^{2n} . The **complex tangent space** $T_x^{\mathbb{C}} X$ is the vector space over \mathbb{C} generated by

$$\left(\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}, \frac{\partial}{\partial \bar{z}_1}, \dots, \frac{\partial}{\partial \bar{z}_n} \right),$$

a $2n$ -dimensional vector space over \mathbb{C} . Then $T_x^{\mathbb{C}} X = T_x^{\mathbb{R}} X \otimes_{\mathbb{R}} \mathbb{C}$.

³Exercise

3.6 Holomorphic differential forms on complex manifolds

Definition 3.20. Let X be a complex manifold of dimension n . Let $\{(U_\alpha, \phi_\alpha)\}$ be a complex structure on X . A **holomorphic p -form** on X is the data ω_α , the p -forms on $\phi_\alpha(U_\alpha) \subset \mathbb{C}^n$ such that if

$$h_{\alpha\beta} = \phi_\alpha \circ \phi_\beta^{-1} : \phi_\beta(U_\alpha \cap U_\beta) \rightarrow \phi_\alpha(U_\alpha \cap U_\beta),$$

then $h_{\alpha\beta}^* \omega_\beta = \omega_\alpha$ for all α and β .

Notation 3.21.

$$\Omega_x^p(X) = H^0(X, \Omega_x^p) = \{\text{holomorphic } p\text{-forms on } X\},$$

$$\mathcal{O}_x(X) = H^0(X, \mathcal{O}_x) = \{\text{holomorphic functions on } X\}.$$

$R = \mathcal{O}_x(X)$ is a ring and $M = \Omega_x^p(X)$ is an R -module.

Lemma 3.22. Let $f : X \rightarrow Y$ be holomorphic. Then $f^* : \Omega^p(Y) \rightarrow \Omega^p(X)$.

Proof. Let $\{(U_\alpha, \phi_\alpha)\}$ be a complex structure on Y . We can write $f^{-1}(U_\alpha) = \bigcup_{\alpha,\beta} V_{\alpha,\beta}$ where $\{(V_{\alpha,\beta}, \psi_{\alpha,\beta})\}$ is a complex structure on X , so

$$\mathbb{C}^n \xleftarrow{\psi_{\alpha,\beta}} V_{\alpha,\beta} \xrightarrow{f|_{V_{\alpha,\beta}}} U_\alpha \xrightarrow{\phi_\alpha} \mathbb{C}^n.$$

Assume ω is defined by ω_α on $\phi_\alpha(U_\alpha)$. Let

$$\omega_{\alpha,\beta} = \left(\left(\psi_{\alpha,\beta}^{-1} \right)^* \circ f^* \circ \phi_\alpha^* \right) (\omega_\alpha)$$

be a p -form on $\psi_{\alpha,\beta}(V_{\alpha,\beta})$. Check that $\omega_{\alpha,\beta}$ are compatible with respect to the atlas on X .⁴ □

As in the local case, we can define

$$\begin{array}{ccc} \Omega_x^p(X) \otimes \Omega_x^q(X) & \longrightarrow & \Omega_x^{p+q}(X) \\ \omega_1 \otimes \omega_2 & \longmapsto & \omega_1 \wedge \omega_2 \end{array}.$$

Similarly there exists $d : \Omega_x^p(X) \rightarrow \Omega_x^{p+1}(X)$.

⁴Exercise

4 Vector bundles

4.1 Holomorphic vector bundles

Definition 4.1. Let X be a complex manifold. A **holomorphic vector bundle** E of rank r on X is a complex manifold E , a holomorphism $\pi : E \rightarrow X$, and an open covering U_α of X such that there exists a biholomorphism

$$\psi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{C}^r,$$

such that if $p_\alpha : U_\alpha \times \mathbb{C}^r \rightarrow U_\alpha$ is the projection then $\pi|_{\pi^{-1}(U_\alpha)} = p_\alpha \circ \psi_\alpha$, so

$$\begin{array}{ccc} E & \supset & \pi^{-1}(U_\alpha) \xrightarrow{\psi_\alpha} U_\alpha \times \mathbb{C}^r \\ \pi \downarrow & & \downarrow \pi \swarrow p_\alpha \\ X & \supset & U_\alpha \end{array} .$$

A vector bundle of rank one is called a **line bundle**.

For any $x \in X$, there exists α such that $x \in U_\alpha$, so

$$\begin{array}{ccc} \pi^{-1}(x) & \xrightarrow{\psi_\alpha} & \{x\} \times \mathbb{C}^r \\ \pi \downarrow & & \swarrow p_\alpha \\ x & & \end{array} .$$

Then $E(x) = \pi^{-1}(x)$ is a vector space of rank r over \mathbb{C} . Let $U_\alpha \ni x \in U_\beta$. There exists a biholomorphism

$$\mathbb{C}^r \cong p_\alpha^{-1}(x) \rightarrow p_\beta^{-1}(x) \cong \mathbb{C}^r,$$

because they are both biholomorphic to $\pi^{-1}(x)$, so $g_{\alpha\beta}(x) \in \mathrm{GL}_r \mathbb{C}$ because all the biholomorphisms from $\mathbb{C}^r \rightarrow \mathbb{C}^r$ are linear. The holomorphisms

$$g_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow \mathrm{GL}_r \mathbb{C}$$

are called **transition functions**. Then

$$\begin{array}{ccc} p_\alpha^{-1}(x) & \xrightarrow{\mathrm{id}} & p_\alpha^{-1}(x) \\ & \searrow & \swarrow \\ & p_\beta^{-1}(x) & \end{array} ,$$

so

$$(g_{\alpha\beta} \circ g_{\beta\alpha})(x) = x, \quad x \in U_\alpha \cap U_\beta,$$

and

$$\begin{array}{ccc} p_\alpha^{-1}(x) & \xrightarrow{g_{\alpha\gamma}} & p_\gamma^{-1}(x) \\ & \searrow & \swarrow \\ & p_\beta^{-1}(x) & \end{array} ,$$

so

$$(g_{\alpha\beta} \circ g_{\beta\gamma})(x) = g_{\alpha\gamma}(x), \quad x \in U_\alpha \cap U_\beta \cap U_\gamma.$$

Definition 4.2. Let X be a complex manifold, and let E and F be vector bundles on X of rank r and s respectively, with $\pi : E \rightarrow X$ and $\pi' : F \rightarrow X$. A **holomorphic map** $f : E \rightarrow F$ is a holomorphic function $E \rightarrow F$ such that $\pi = \pi' \circ f$ and such that the rank of the induced linear map $E(x) \rightarrow F(x)$ is independent of $x \in X$, so

$$\begin{array}{ccc} E & \xrightarrow{f} & F \\ \pi \searrow & & \swarrow \pi' \\ & X & \end{array} , \quad \begin{array}{ccc} E(x) = \pi^{-1}(x) & \xrightarrow{f} & \pi'^{-1}(x) = F(x) \\ \pi \searrow & & \swarrow \pi' \\ & x & \end{array} .$$

4.2 Examples of holomorphic vector bundles

Example 4.3. $\pi : E = X \times \mathbb{C}^r \rightarrow X$ is a vector bundle of rank r , called **trivial**.

Example 4.4. Algebra of vector bundles. Let $\pi : E \rightarrow X$ and $\pi'^{-1} : F \rightarrow X$ be vector bundles on X of rank r and s respectively.

- The **direct sum** $E \oplus F$ is the $(r + s)$ -vector bundle such that

$$(E \oplus F)(x) = E(x) \oplus F(x), \quad x \in X.$$

The idea is to take an open cover which trivialises both E and F . Find the transition function of $E \oplus F$.⁵

- The **tensor product** $E \otimes F$ is the $(r \cdot s)$ -vector bundle such that

$$(E \otimes F)(x) = E(x) \otimes F(x), \quad x \in X.$$

- The **p -th exterior power** of E is the vector bundle $\Lambda^p E$ such that

$$(\Lambda^p E)(x) = \Lambda^p(E(x)), \quad x \in X.$$

If $p = r = \text{rk } E$ then $\det E = \Lambda^r E$ is a line bundle on X .

- The **dual** of E is the rank r vector bundle E^* such that

$$E^*(x) = (E(x))^*, \quad x \in X,$$

the dual $\text{Hom}(E(x), \mathbb{C})$ of $E(x)$.

- Let $f : E \rightarrow F$ be a holomorphic map. Then the **kernel** $\text{Ker } f$ is a vector bundle such that

$$(\text{Ker } f)(x) = \text{Ker } f(x) \subset E(x), \quad x \in X.$$

The **cokernel** $\text{Coker } f$ is a vector bundle such that

$$(\text{Coker } f)(x) = \text{Coker } f(x) \subset F(x), \quad x \in X.$$

Example 4.5. Let $X = \mathbb{P}_{\mathbb{C}}^1$, and let

$$\mathcal{O}(-1) = \{(x, v) \mid x = [x_0, \dots, x_n] \in \mathbb{P}_{\mathbb{C}}^n, v = \mu(x_0, \dots, x_n), \mu \in \mathbb{C}\} \subset \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C}^{n+1}.$$

Then $\pi = p_1 : \mathcal{O}(-1) \rightarrow \mathbb{P}_{\mathbb{C}}^n$, so

$$\pi^{-1}([x_0, \dots, x_n]) = \{v = \mu(x_0, \dots, x_n) \mid \mu \in \mathbb{C}\} \cong \mathbb{C}^1.$$

Let $\{U_i\}$ be an open covering of X given by $U_i = \{[x_0, \dots, x_n] \mid x_i \neq 0\}$. We define

$$\begin{aligned} \psi_i : \quad & \pi^{-1}(U_i) \longrightarrow U_i \times \mathbb{C} \\ & ([x_0, \dots, x_n], (v_0, \dots, v_n)) \longmapsto ([x_0, \dots, x_n], v_i) \end{aligned}$$

which is a biholomorphism. Thus $\mathcal{O}(-1)$ is a complex manifold and $\mathcal{O}(-1)$ is a line bundle. The **tautological line bundle** $\mathcal{O}(1)$ is the dual of $\mathcal{O}(-1)$. Let

$$\mathcal{O}(k) = \begin{cases} X \times \mathbb{C} & k = 0 \\ \mathcal{O}(1)^{\otimes k} & k > 0 \\ \mathcal{O}(-1)^{\otimes k} & k < 0 \end{cases}.$$

Then $\mathcal{O}(k) = \mathcal{O}(-k)^*$.⁶ On $\mathbb{P}_{\mathbb{C}}^n$ these are the only line bundles. That is, if \mathcal{L} is a line bundle on $\mathbb{P}_{\mathbb{C}}^1$, there exists $k \in \mathbb{Z}$ such that $\mathcal{L} \cong \mathcal{O}(k)$. Let $X = \mathbb{P}_{\mathbb{C}}^1$, and let E be a line bundle of rank r on X . Then

$$E \cong \bigoplus_{i=1}^r \mathcal{O}(a_i), \quad a_1, \dots, a_r \in \mathbb{Z}.$$

This is false for $X = \mathbb{P}_{\mathbb{C}}^n$, with $n \geq 2$.

⁵Exercise

⁶Exercise

Definition 4.6. Let $f : Y \rightarrow X$ be a holomorphism between complex manifolds, and let E be a vector bundle of rank r on X . Then there exists a vector bundle f^*E of rank r on Y defined by

$$f^*E = \{(y, v) \in Y \times E \mid f(y) = \pi(v)\},$$

the **fibre product** of E and Y over X , such that

$$\begin{array}{ccc} f^*E & \xrightarrow{f'} & E \\ \pi' \downarrow & & \downarrow \pi \\ Y & \xrightarrow{f} & X \end{array}$$

Let $U = \{U_i\}$ be an open cover of X which trivialises E , so

$$\begin{array}{ccc} \pi^{-1}(U_i) & \xrightarrow{\psi_i} & U_i \times \mathbb{C}^r \\ & \searrow \pi \quad \swarrow p_1 & \\ & U_i & \end{array}$$

Then $U' = \{f^{-1}(U_i)\}$ is an open covering of Y , so

$$\begin{array}{ccccccc} \pi'^{-1}(f^{-1}(U_i)) & \xrightarrow{f'} & \pi^{-1}(U_i) & \xrightarrow{\psi_i} & U_i \times \mathbb{C}^r & \xrightarrow{p_2} & \mathbb{C}^r \\ \pi' \downarrow & & \downarrow \pi & & & & \\ f^{-1}(U_i) & \xrightarrow{f} & U_i & & & & \end{array},$$

and

$$\begin{aligned} \pi'^{-1}(f^{-1}(U_i)) &= \{(y, v) \in f^{-1}(U_i) \times \pi^{-1}(U_i) \mid f(y) = \pi(v)\} \longrightarrow f^{-1}(U_i) \times \mathbb{C}^r \\ (y, v) &\longmapsto (y, p_2(\psi_i(v))) \end{aligned}$$

is a biholomorphism. Thus f^*E is a vector bundle, where

$$f^*E(y) = \pi'^{-1}(y) = E(f(y)), \quad y \in Y.$$

Notation 4.7. Let $f : Y \rightarrow X$ be a morphism, and let E be a vector bundle on X . Then $f^*E = E|_Y$, mostly used if $f : Y \hookrightarrow X$.

Definition 4.8. Let E be a holomorphic vector bundle on a complex manifold X , and let $\pi : E \rightarrow X$. A **section** of E is a holomorphic function $s : X \rightarrow E$ such that $\pi \circ s = \text{id}_X$.

Example 4.9. Let $E = X \times \mathbb{C}^r$ be the trivial vector bundle of rank r . Fix $v \in \mathbb{C}^r$. Then

$$\begin{aligned} s_v &: X \longrightarrow E \\ x &\longmapsto (x, v) \end{aligned}$$

is a section of E . If v_1, \dots, v_r is a basis of \mathbb{C}^r then s_{v_1}, \dots, s_{v_r} have the property that $s_{v_1}(x), \dots, s_{v_r}(x)$ forms a basis of $E(x)$. Vice versa, assume E is a vector bundle on X of rank r such that there exist sections s_1, \dots, s_r of E such that for all $x \in X$, $s_1(x), \dots, s_r(x)$ is a basis of $E(x)$. Then $E \cong X \times \mathbb{C}^r$, since

$$\begin{aligned} X \times \mathbb{C}^r &\longrightarrow E \\ (x, (v_1, \dots, v_r)) &\longmapsto \sum_i v_i s_i(x) \end{aligned}$$

is a biholomorphism. Then s_1, \dots, s_r is called a **holomorphic frame** for E . Recall that for all $E \rightarrow X$ and for all $x \in X$ there exists open $U \ni x$ such that $E|_U$ is trivial, so there exists a frame on U for $E|_U$. This is called a **local frame** around x .

Example 4.10. Let X be a complex manifold of dimension n , and let (z_1, \dots, z_n) be coordinates on \mathbb{C}^n . There exists an atlas $\{(U_\alpha, \phi_\alpha)\}$ for $\phi_\alpha : U_\alpha \rightarrow V_\alpha \subset \mathbb{C}^n$. For all $x \in U_\alpha$, $T_x U_\alpha \rightarrow T_{\phi_\alpha(x)} V_\alpha$, and $T_{\phi_\alpha(x)} V_\alpha = \left\langle \frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n} \right\rangle$ is a frame of T_{V_α} . Let

$$T_X = \bigcup_{x \in X} T_x X,$$

and let $\pi^{-1} : T_X \rightarrow X$ such that $\pi^{-1}(x) = T_x X$. Then T_X is a holomorphic vector bundle of rank n called the **tangent bundle**, where $U = \{U_\alpha\}$ and

$$\psi_\alpha : \pi^{-1}(U_\alpha) = T_X|_{U_\alpha} \rightarrow T_{\mathbb{C}^n}|_{V_\alpha} \cong V_\alpha \times \mathbb{C}^r \rightarrow U_\alpha \times \mathbb{C}^r$$

defines the trivialisation. The **cotangent bundle** of X is

$$\Omega_X^1 = T_X^*,$$

and let

$$\Omega_X^p = \Lambda^p \Omega_X^1, \quad p \geq 1.$$

A holomorphic p -form on X is a section of Ω_X^p .⁷

4.3 Complexification of tangent bundles

Let X be a complex manifold. How to view X as a differentiable manifold? Let V be a vector space of dimension m over \mathbb{R} . An **almost complex structure** on V is a linear map $J : V \rightarrow V$ such that $J^2 = -\text{id}_V$. If V admits an almost complex structure, then V can be seen as a vector space over \mathbb{C} . Let $\lambda = a + ib$ for $a, b \in \mathbb{R}$, and let $v \in V$. Define

$$\lambda v = av + bJ(v).$$

If $\lambda_1, \lambda_2 \in \mathbb{C}$, then $\lambda_1(\lambda_2 v) = (\lambda_1 \lambda_2)v$.⁸ Let $v_1, \dots, v_n \in V$ be a basis over \mathbb{C} . Then

$$v_1, \dots, v_n, J(v_1), \dots, J(v_n)$$

is a basis of V over \mathbb{R} . The idea is to assume that $a_i, b_i \in \mathbb{R}$ such that $\sum_i a_i v_i + \sum_i b_i J(v_i) = 0$, then

$$0 = \sum_i a_i v_i + \sum_i b_i J(v_i) = \sum_i (a_i v_i + b_i J(v_i)) = \sum_i (a_i + ib_i) v_i,$$

so $a_i + ib_i = 0$ for all i . Thus $a_i = b_i = 0$, so $m = 2n$. On a vector space an almost complex structure is a complex structure. Let V be a vector space of dimension $2n$ over \mathbb{R} . Then the **complexification** $V_{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$ of V is a \mathbb{C} -vector space of dimension $2n$ over \mathbb{C} , where

$$\begin{aligned} \lambda & : V_{\mathbb{C}} \longrightarrow V_{\mathbb{C}} \\ v \otimes \mu & \longmapsto v \otimes \mu \lambda, \quad \lambda \in \mathbb{C}. \end{aligned}$$

Let J be an almost complex structure on V . Then we can extend J to a linear map

$$\begin{aligned} J & : V_{\mathbb{C}} \longrightarrow V_{\mathbb{C}} \\ v \otimes \mu & \longmapsto J(v) \otimes \mu, \end{aligned}$$

such that $J^2 = -\text{id}_{V_{\mathbb{C}}}$,⁹ so $J^2 + \text{id}_{V_{\mathbb{C}}} = 0$. Thus the eigenvalues of J on $V_{\mathbb{C}}$ are $\pm i$. Let $V^{1,0}$ be the eigenspace for i and $V^{0,1}$ be the eigenspace for $-i$, so

$$V_{\mathbb{C}} = V^{1,0} \oplus V^{0,1}.$$

The **conjugation**

$$\begin{aligned} \bar{\cdot} & : V_{\mathbb{C}} \longrightarrow V_{\mathbb{C}} \\ v \otimes \mu & \longmapsto v \otimes \bar{\mu} \end{aligned}$$

on $V_{\mathbb{C}}$ is linear over \mathbb{R} , such that $\overline{V^{1,0}} = V^{0,1}$ and $\overline{V^{0,1}} = V^{1,0}$,¹⁰ so $V^{1,0}$ and $V^{0,1}$ are \mathbb{C} -vector spaces of dimension n .

⁷Exercise

⁸Exercise

⁹Exercise

¹⁰Exercise

Example 4.11. Let $W = \mathbb{C}^n$ with coordinates (z_1, \dots, z_n) , and let $z_j = x_j + iy_j$ with coordinates $(x_1, y_1, \dots, x_n, y_n)$ for \mathbb{R}^{2n} . Define

$$J : \mathbb{R}^{2n} \longrightarrow \mathbb{R}^{2n} \\ (x_1, y_1, \dots, x_n, y_n) \longmapsto (-y_1, x_1, \dots, -y_n, x_n) .$$

Then $J^2 = \text{id}_{\mathbb{R}^{2n}}$, and J is the **standard almost complex structure** on \mathbb{R}^{2n} . Let $V = \mathbb{R}^{2n}$, so $V_{\mathbb{C}} \cong \mathbb{C}^{2n}$ with complex coordinates $(x_1, y_1, \dots, x_n, y_n)$. Then $V^{0,1}$ is spanned by $x_j - iy_j$ and $V^{1,0}$ is spanned by $x_j + iy_j$, where $\overline{x_j + iy_j} = x_j - iy_j$ for $j = 1, \dots, n$.

Definition 4.12. Let X be a differentiable manifold. A **real, or complex, vector bundle** of rank r is a differentiable manifold E with a smooth morphism $\pi : E \rightarrow X$ such that if $K = \mathbb{R}$, or $K = \mathbb{C}$, then there exists an open covering $U = \{U_i\}$ of X such that

- for all $x \in X$, the fibre of π , $E(x) = \pi^{-1}(x)$, is a vector space of rank r over K ,
- for all i there exists a diffeomorphism h_i such that

$$\begin{array}{ccc} \pi^{-1}(U_i) & \xrightarrow{h_i} & U_i \times K^r \xrightarrow{p_2} K^r \\ & \searrow \pi & \swarrow p_1 \\ & U_i & \end{array} ,$$

and for all x , $p_2 \circ h_i : E(x) \rightarrow K^r$ is an isomorphism of vector spaces.

Pull-backs, sections, exterior powers, tensors, direct sums, frames, etc are the same as holomorphic vector bundles, where holomorphic becomes smooth and biholomorphic becomes diffeomorphic, and for all X there exists a tangent bundle T_X . Assume X is a complex manifold of dimension n . Let T_X be the holomorphic tangent bundle of X . Then X is also a differentiable manifold of dimension $2n$, so let $T_X^{\mathbb{R}}$ be the **real tangent bundle** of X , which is a rank $2n$ vector bundle, and let $T_X^{\mathbb{C}} = T_X^{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ be the **complex tangent bundle** of X , which is a non-holomorphic complex vector bundle of rank $2n$. Smooth morphisms of real or complex vector bundles are defined similarly as holomorphisms between holomorphic vector bundles such that the rank of the image is constant, so

$$\begin{array}{ccc} E & \xrightarrow{f} & F \\ & \searrow \pi & \swarrow \pi' \\ & X & \end{array} .$$

Let X be a differentiable manifold of dimension $m = 2n$. Then an **almost complex structure** on X is a smooth morphism between the real tangent bundle $J : T_X^{\mathbb{R}} \rightarrow T_X^{\mathbb{R}}$ such that $J^2 = -\text{id}$. In particular, $J(x) : T_x^{\mathbb{R}} X \rightarrow T_x^{\mathbb{R}} X$ is an almost complex structure for all $x \in X$.

Proposition 4.13. *Let X be a complex manifold. Then the underlying differentiable manifold admits an almost complex structure $J : T_X^{\mathbb{R}} \rightarrow T_X^{\mathbb{R}}$ such that $J^2 = -\text{id}$.*

Proof. Let $x \in X$, and let (U, ϕ) be a complex chart around x such that

$$\begin{array}{ccc} \phi & : & U \longrightarrow V \\ & & x \longmapsto 0 \end{array} .$$

Fix holomorphic coordinates (z_1, \dots, z_n) on U . The tangent bundle of X on U is trivial, with a local frame $\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}$, so

$$T_X|_U \xrightarrow{\sim} T_V = V \times \mathbb{C}^n .$$

Define $x_i = \text{Re } z_i$ and $y_i = \text{Im } z_i$. Then $(x_1, y_1, \dots, x_n, y_n)$ are smooth coordinates $U \rightarrow \mathbb{R}$ around x , and $\frac{\partial}{\partial x_1}, \frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial x_n}, \frac{\partial}{\partial y_n}$ define a local smooth frame of $T_X^{\mathbb{R}}$ on U , so

$$T_X^{\mathbb{R}}|_U \xrightarrow{\sim} T_V = V \times \mathbb{R}^{2n} .$$

In particular, there exists an almost complex structure J_U for $T_V \cong T_X^{\mathbb{R}}|_U$, so

$$J_U : T_X^{\mathbb{R}}|_U \rightarrow T_X^{\mathbb{R}}|_U, \quad J_U^2 = -\text{id} .$$

Lecture 11
Thursday
30/01/20

Let $f : V \rightarrow V$ be a biholomorphism, so

$$\begin{array}{ccc} & U \cap U' & \\ \phi \swarrow & & \searrow \phi \\ V & \xrightarrow{f} & V \end{array},$$

and let z'_1, \dots, z'_n be local holomorphic coordinates given by

$$z'_i = f_i(z_1, \dots, z_n), \quad f_i = p_i \circ f,$$

where $p_i : \mathbb{C}^n \rightarrow \mathbb{C}$ is the i -th projection. Define

$$x'_i = \operatorname{Re} z'_i = \operatorname{Re} f_i(z_1, \dots, z_n) = u_i(z_1, \dots, z_n), \quad y'_i = \operatorname{Im} z'_i = \operatorname{Im} f_i(z_1, \dots, z_n) = v_i(z_1, \dots, z_n),$$

so $f_j = u_j + iv_j$. The real Jacobian J_f of f is given by the derivatives of u_j and v_j with respect to $x_1, y_1, \dots, x_n, y_n$, a $(2n \times 2n)$ -matrix of $n \times n$ blocks of 2×2 blocks of

$$\begin{pmatrix} \frac{\partial u_j}{\partial x_k} & \frac{\partial u_j}{\partial y_k} \\ \frac{\partial v_j}{\partial x_k} & \frac{\partial v_j}{\partial y_k} \end{pmatrix}.$$

These define the transition function of $T_X^{\mathbb{R}}$. To show that J extends to X , it is enough to show that J commutes with J_f at each point, so

$$\begin{array}{ccc} T_X^{\mathbb{R}}|_{U \cap U'} & \xrightarrow{J_f} & T_X^{\mathbb{R}}|_{U \cap U'} \\ J \downarrow & & \downarrow J \\ T_X^{\mathbb{R}}|_{U \cap U'} & \xrightarrow{J_f} & T_X^{\mathbb{R}}|_{U \cap U'} \end{array}.$$

Since f_j is holomorphic $\frac{\partial f_j}{\partial \bar{z}_k} = 0$ for all j and k , so the Cauchy-Riemann equations

$$\frac{\partial u_j}{\partial x_k} - \frac{\partial v_j}{\partial y_k} = 0, \quad \frac{\partial v_j}{\partial x_k} + \frac{\partial u_j}{\partial y_k} = 0,$$

or

$$\begin{pmatrix} \frac{\partial u_j}{\partial x_k} & \frac{\partial u_j}{\partial y_k} \\ \frac{\partial v_j}{\partial x_k} & \frac{\partial v_j}{\partial y_k} \end{pmatrix} = \begin{pmatrix} \frac{\partial v_j}{\partial y_k} & \frac{\partial u_j}{\partial y_k} \\ -\frac{\partial u_j}{\partial y_k} & \frac{\partial v_j}{\partial y_k} \end{pmatrix},$$

hold. Since J is the standard almost complex structure on \mathbb{R}^{2n} , where $x_j \mapsto y_j$ and $y_j \mapsto -x_j$,

$$J = \begin{pmatrix} 0 & 1 & & \\ -1 & 0 & & \\ & & \ddots & \\ & 0 & & 0 & 1 \\ & & & -1 & 0 \end{pmatrix}.$$

Check that J_f commutes with J .¹¹

□

Corollary 4.14. *Every complex manifold is orientable.*

Proof. We prove that if $T_X^{\mathbb{R}}$ admits an almost complex structure then X is an orientable manifold. For all $x \in X$ choose the orientation on $T_x^{\mathbb{R}}X$, a vector space of dimension $2n$ over \mathbb{R} , given by any ordered basis of the form

$$v_1, \dots, v_n, J(v_1), \dots, J(v_n).$$

Assume that $v_1, \dots, v_n, J(v_1), \dots, J(v_n)$ and $w_1, \dots, w_n, J(w_1), \dots, J(w_n)$ are ordered bases. Show that the determinant of the matrix given by the change of basis is positive.¹²

□

¹¹Exercise

¹²Exercise

4.4 Differential forms on complex tangent bundles

Let X be a complex manifold. Then there exists an almost complex structure $J : T_X^{\mathbb{R}} \rightarrow T_X^{\mathbb{R}}$ on X . Then J extends to

$$J : \begin{array}{ccc} T_X^{\mathbb{C}} & \longrightarrow & T_X^{\mathbb{C}} \\ v \otimes \mu & \longmapsto & J(v) \otimes \mu \end{array}.$$

For all x , $J(x)$ has two eigenvalues $\pm i$, so

$$T_X^{\mathbb{C}} = T_X^{1,0} \oplus T_X^{0,1},$$

which are complex vector bundles and **eigenbundles**. Locally $T_X^{1,0}$ and $T_X^{0,1}$ are spanned by the frames $\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}$ and $\frac{\partial}{\partial \bar{z}_1}, \dots, \frac{\partial}{\partial \bar{z}_n}$ respectively. Moreover there exists a conjugation

$$\begin{array}{ccc} T_X^{\mathbb{C}} & \longrightarrow & T_X^{\mathbb{C}} \\ v \otimes \mu & \longmapsto & v \otimes \bar{\mu} \end{array}$$

over \mathbb{R} , such that $\overline{T_X^{1,0}} = T_X^{0,1}$ and $\overline{T_X^{0,1}} = T_X^{1,0}$. Let

$$\Omega_{X,\mathbb{C}}^1 = (T_X^{\mathbb{C}})^*$$

be the dual of the complex vector bundle $T_X^{\mathbb{C}}$. Then

$$\Omega_{X,\mathbb{C}}^1 = \Omega_{X,\mathbb{R}}^1 \otimes_{\mathbb{R}} \mathbb{C} = \Omega_{X,\mathbb{C}}^{1,0} \oplus \Omega_{X,\mathbb{C}}^{0,1} = (T_X^{1,0})^* \oplus (T_X^{0,1})^*.$$

Exercise. Let V and W be vector spaces. Show that

$$\Lambda^k(V \oplus W) = \bigoplus_{p+q=k} \Lambda^p V \otimes \Lambda^q W$$

is a canonical isomorphism.

Thus, for all $k \geq 0$

$$\Omega_{X,\mathbb{C}}^k = \Lambda^k \Omega_{X,\mathbb{C}}^1 = \bigoplus_{p+q=k} \Omega_{X,\mathbb{C}}^{p,q}, \quad \Omega_{X,\mathbb{C}}^{p,q} = \Lambda^p \Omega_{X,\mathbb{C}}^{1,0} \otimes \Lambda^q \Omega_{X,\mathbb{C}}^{0,1},$$

where $\Omega_{X,\mathbb{C}}^{p,q}$ is a complex vector bundle for any p and q .

Definition 4.15. The sections of $\Omega_{X,\mathbb{C}}^{p,q}$ are called (p, q) -forms on X , or **forms of type (p, q)** .

Locally, let $x \in X$, and let $(U \ni x, \phi)$ be a holomorphic chart for $\phi : U \xrightarrow{\sim} V \subset \mathbb{C}^n$. A (p, q) -form on U can be locally written as

$$\omega = \sum_{I,J} \alpha_{I,J} dz_I \wedge d\bar{z}_J = \sum_{|I|=p, |J|=q} \alpha_{I,J} dz_{i_1} \wedge \dots \wedge dz_{i_p} \wedge d\bar{z}_{j_1} \wedge \dots \wedge d\bar{z}_{j_q},$$

where $\alpha_{I,J}$ are smooth functions on U . Let X be a manifold. If E is a complex vector bundle then

$$C^\infty(X, E) = \{\text{smooth sections of } E\}.$$

The **differential**

$$d : C^\infty(X, \Omega_{X,\mathbb{C}}^k) \rightarrow C^\infty(X, \Omega_{X,\mathbb{C}}^{k+1})$$

satisfies the Leibnitz rule and $d^2 = 0$, so $d(d\omega) = 0$. If $\omega \in C^\infty(X, \Omega_{X,\mathbb{C}}^{p,q})$, then $d\omega \in C^\infty(X, \Omega_{X,\mathbb{C}}^{p+q+1})$.

Assume that locally $\omega = \sum_{I,J} \alpha_{I,J} dz_I \wedge d\bar{z}_J$. Then

$$d\omega = \sum_{I,J} d\alpha_{I,J} dz_I \wedge d\bar{z}_J, \quad d\alpha_{I,J} = \sum_{i=1}^n \frac{\partial}{\partial z_i} \alpha_{I,J} dz_i + \sum_{i=1}^n \frac{\partial}{\partial \bar{z}_i} \alpha_{I,J} d\bar{z}_i.$$

Lecture 12
Tuesday
04/02/20

Let

$$\partial\alpha_{I,J} = \sum_{i=1}^n \frac{\partial}{\partial z_i} \alpha_{I,J} dz_i \in C^\infty(X, \Omega_X^{1,0}), \quad \bar{\partial}\alpha_{I,J} = \sum_{i=1}^n \frac{\partial}{\partial \bar{z}_i} \alpha_{I,J} d\bar{z}_i \in C^\infty(X, \Omega_X^{0,1}).$$

Then $d = \partial + \bar{\partial}$ for smooth functions. Back to $d\omega$. Then

$$d\omega = \sum_{I,J} d\alpha_{I,J} dz_I \wedge d\bar{z}_J = \sum_{I,J} \partial\alpha_{I,J} dz_I \wedge d\bar{z}_J + \sum_{I,J} \bar{\partial}\alpha_{I,J} dz_I \wedge d\bar{z}_J.$$

Let

$$\partial\omega = \sum_{I,J} \partial\alpha_{I,J} dz_I \wedge d\bar{z}_J, \quad \bar{\partial}\omega = \sum_{I,J} \bar{\partial}\alpha_{I,J} dz_I \wedge d\bar{z}_J.$$

Then $d = \partial + \bar{\partial}$ for ω .

Lemma 4.16. *The linear maps*

$$\partial : C^\infty(X, \Omega_X^{p,q}) \rightarrow C^\infty(X, \Omega_X^{p+1,q}), \quad \bar{\partial} : C^\infty(X, \Omega_X^{p,q}) \rightarrow C^\infty(X, \Omega_X^{p,q+1})$$

satisfy the Leibnitz rule. That is, if $\omega \in C^\infty(X, \Omega_X^{p,q})$ and $\eta \in C^\infty(X, \Omega_X^{p',q'})$, then

$$\partial(\omega \wedge \eta) = \partial\omega \wedge \eta + (-1)^{p+q} \omega \wedge \partial\eta, \quad \bar{\partial}(\omega \wedge \eta) = \bar{\partial}\omega \wedge \eta + (-1)^{p+q} \omega \wedge \bar{\partial}\eta.$$

Proof. d satisfies the Leibnitz rule

$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^{p+q} \omega \wedge d\eta,$$

since $\omega \in C^\infty(X, \Omega_{X,\mathbb{C}}^{p+q})$, so

$$\begin{aligned} \partial(\omega \wedge \eta) + \bar{\partial}(\omega \wedge \eta) &= (\partial\omega + \bar{\partial}\omega) \wedge \eta + (-1)^{p+q} \omega \wedge (\partial\eta + \bar{\partial}\eta) \\ &= (\partial\omega \wedge \eta + (-1)^{p+q} \omega \wedge \partial\eta) + (\bar{\partial}\omega \wedge \eta + (-1)^{p+q} \omega \wedge \bar{\partial}\eta). \end{aligned}$$

Then $\partial(\omega \wedge \eta)$ and $\partial\omega \wedge \eta + (-1)^{p+q} \omega \wedge \partial\eta$ are $(p+1, q)$ -forms, and $\bar{\partial}(\omega \wedge \eta)$ and $\bar{\partial}\omega \wedge \eta + (-1)^{p+q} \omega \wedge \bar{\partial}\eta$ are $(p, q+1)$ -forms. Forms of the same type in the decomposition of $d(\omega \wedge \eta)$ must coincide. \square

Lemma 4.17. $\partial^2 = \bar{\partial}^2 = \bar{\partial}\partial + \partial\bar{\partial} = 0$.

Proof. Let $\omega \in C^\infty(X, \Omega_X^{p,q})$. Because $d^2 = 0$,

$$0 = d^2\omega = (\partial + \bar{\partial})((\partial + \bar{\partial})\omega) = \partial^2\omega + \partial\bar{\partial}\omega + \bar{\partial}\partial\omega + \bar{\partial}^2\omega.$$

Then $d^2\omega$ is a $(p+q+2)$ -form, $\partial^2\omega$ is a $(p+2, q)$ -form, $\partial\bar{\partial}\omega + \bar{\partial}\partial\omega$ is a $(p+1, q+1)$ -form, and $\bar{\partial}^2\omega$ is a $(p, q+2)$ -form. Forms of the same type in the decomposition of $d^2\omega$ must coincide. \square

4.5 Dolbeault cohomology

Let X be a complex manifold. Fix $p, q \geq 0$. Let

$$\begin{aligned} \mathcal{Z}^{p,q}(X) &= \text{Ker} \left(\bar{\partial} : C^\infty(X, \Omega_X^{p,q}) \rightarrow C^\infty(X, \Omega_X^{p,q+1}) \right) \\ &= \{ \omega \in C^\infty(X, \Omega_X^{p,q}) \mid \bar{\partial}\omega = 0 \} \end{aligned}$$

and let

$$\begin{aligned} \mathcal{B}^{p,q}(X) &= \text{Im} \left(\bar{\partial} : C^\infty(X, \Omega_X^{p,q-1}) \rightarrow C^\infty(X, \Omega_X^{p,q}) \right) \\ &= \left\{ \omega \in C^\infty(X, \Omega_X^{p,q}) \mid \exists \eta \in C^\infty(X, \Omega_X^{p,q-1}), \omega = \bar{\partial}\eta \right\}. \end{aligned}$$

Since $\bar{\partial}^2 = 0$ we have $\mathcal{B}^{p,q}(X) \subset \mathcal{Z}^{p,q}(X)$ for all p and q . The **Dolbeault cohomology** is

$$H^{p,q}(X) = \mathcal{Z}^{p,q}(X) / \mathcal{B}^{p,q}(X).$$

Exercise. Assume X and Y are biholomorphic complex manifolds. Then $H^{p,q}(X) = H^{p,q}(Y)$.

If $H^{p,q}(X)$ is finite dimensional then we define the **Hodge numbers** of X as

$$h^{p,q} = \dim_{\mathbb{C}} H^{p,q}(X).$$

Our goal is if X is Kähler and compact

$$\bigoplus_{p+q=k} H^{p,q}(X) = H^{p+q}(X)$$

as the de Rham cohomology. In particular this is true if X is projective. How to compute $H^{p,q}(X)$? We need to use analysis.

Proposition 4.18. *Let X be a complex manifold. Then there exists an isomorphism*

$$H^{p,0}(X) \cong H^0(X, \Omega_X^p) = \{\text{holomorphic sections of } \Omega_X^p\} = \{\text{holomorphic } p\text{-forms on } X\}, \quad p \geq 0.$$

Remark 4.19. If X is compact then $H^{0,0}(X) = \mathbb{C}$ because $H^{0,0}(X) = H^0(X, \mathcal{O}_X)$ are constants.

Proof.

$$H^{p,0}(X) = \mathcal{Z}^{p,0}(X) / \mathcal{B}^{p,0}(X) = \mathcal{Z}^{p,0}(X) = \left\{ \omega \in C^\infty(X, \Omega_X^{p,0}) \mid \bar{\partial}\omega = 0 \right\}.$$

Locally $\omega = \sum_{|I|=p} \alpha_I dz_I$. Then

$$\bar{\partial}\omega = \sum_{|I|=p} \bar{\partial}\alpha_I dz_I = \sum_{|I|=p} \sum_{i=1}^n \frac{\partial}{\partial \bar{z}_j} \alpha_I d\bar{z}_j \wedge dz_I,$$

where $d\bar{z}_j \wedge dz_I$ are linearly independent. For all I and for all j , the Cauchy-Riemann equations $\frac{\partial}{\partial \bar{z}_j} \alpha_I = 0$ hold, so for all I , α_I is holomorphic. Then $\omega = \sum_{|I|=p} \alpha_I dz_I$ is a holomorphic p -form, so $\omega \in H^0(X, \Omega_X^p)$. \square

4.6 Connections

Let X be a differentiable manifold, and let E be a complex vector bundle. Then

$$C^\infty(X, E) = \{\text{C}^\infty\text{-sections of } E\}.$$

Is there a way to compute the derivatives of these sections?

Definition 4.20. Let X and E be as above. A **connection** of E is a \mathbb{C} -linear map

$$\nabla : C^\infty(X, E) \rightarrow C^\infty(X, \Omega_{X,\mathbb{C}}^1 \otimes E)$$

such that the Leibnitz rule holds, so

$$\nabla(f\omega) = f\nabla(\omega) + df \otimes \omega, \quad f \in C^\infty, \quad \omega \in C^\infty(X, E).$$

The following is the idea. Let $\omega \in C^\infty(X, E)$. Then

$$\nabla(\omega) = \sum_i \eta_i \otimes \omega_i,$$

where η_i are 1-forms on X and ω_i are sections of E . Let $x \in X$, and let $v \in T_x X$. Then

$$\nabla_v(\omega)_x = \sum_i \eta_i(v) \omega_i$$

is a section of E at x . The goal is to study connections locally. Let $x \in X$, and let (U, ϕ) be a chart around x that trivialises E , so $\pi^{-1}(U) = U \times \mathbb{C}^r$ for $\pi : E \rightarrow X$ and $r = \text{rk } E$. Then there exists a frame $s_1, \dots, s_r \in C^\infty(U, E)$ of E on U . Let $\sigma \in C^\infty(X, E)$ be any section. Locally on U we write

$$\sigma \stackrel{U}{=} f = (f_1, \dots, f_r), \quad \sigma = \sum_{i=1}^r f_i s_i, \quad f_1, \dots, f_r \in C^\infty(U).$$

By the Leibnitz rule, on U ,

$$\nabla(\sigma) = \sum_{i=1}^r \nabla(f_i s_i) = \sum_{i=1}^r (f_i \nabla(s_i) + df_i \otimes s_i) \in C^\infty(U, \Omega_{X,\mathbb{C}}^1 \otimes E).$$

Lecture 13
Thursday
06/02/20

Notation 4.21. $df = (df_1, \dots, df_r)$.

Then

$$\nabla(s_j) = \sum_{i=1}^r a_{ij} \otimes s_i, \quad a_{ij} \in C^\infty(U, \Omega_{X, \mathbb{C}}^1).$$

Notation 4.22. $A = (a_{ij})$ is an $(r \times r)$ -matrix with coefficients 1-forms.

With this notation, this becomes

$$\nabla(\sigma) \stackrel{U}{=} Af + df.$$

- A depends very much on the choice of the frame.
- Locally on U , ∇ is determined by A .

Consider another chart (U', ϕ') which also gives a trivialisation of E . So we can choose a corresponding frame s'_1, \dots, s'_r . Assume $\sigma \in C^\infty(U \cap U', E)$. Then $\sigma \stackrel{U}{=} (f_1, \dots, f_r)$ and

$$\sigma \stackrel{U'}{=} f' = (f'_1, \dots, f'_r), \quad \sigma = \sum_{j=1}^r f'_j s'_j, \quad f'_1, \dots, f'_r \in C^\infty(U).$$

Let A' be the matrix with respect to this frame. Then

$$\nabla(\sigma) \stackrel{U'}{=} A'f' + df'.$$