

# M3P8 Algebra III

Lectured by Dr David Helm  
Typeset by David Kurniadi Angdinata

Autumn 2018

## Contents

<b>0</b>	<b>Introduction</b>	<b>2</b>
<b>1</b>	<b>Basic definitions and examples</b>	<b>2</b>
1.1	Rings . . . . .	2
1.2	Polynomial rings . . . . .	3
1.3	Subrings and extensions . . . . .	4
1.4	Integral domains and rings of fractions . . . . .	4
<b>2</b>	<b>Homomorphisms, ideals, and quotients</b>	<b>5</b>
2.1	Homomorphisms . . . . .	5
2.2	Evaluation homomorphisms . . . . .	6
2.3	Images, kernels, and ideals . . . . .	6
2.4	Ideals: examples and basic operations . . . . .	7
2.5	Quotients . . . . .	8
2.6	Prime and maximal ideals . . . . .	8
<b>3</b>	<b>Factorisation</b>	<b>9</b>
3.1	Divisibility, units, associates, and irreducibles . . . . .	9
3.2	Unique factorisation domains . . . . .	9
3.3	Principal ideal domains . . . . .	10
3.4	Euclidean domains . . . . .	11
3.5	Examples . . . . .	11
<b>4</b>	<b>Chinese remainder theorem</b>	<b>12</b>
4.1	Products . . . . .	12
4.2	The Chinese remainder theorem . . . . .	12
4.3	Examples . . . . .	13
<b>5</b>	<b>Fields and field extensions</b>	<b>13</b>
5.1	Prime fields . . . . .	14
5.2	Field extensions . . . . .	14
5.3	Extensions generated by one element . . . . .	14

## 0 Introduction

This course is an introduction to ring theory. The topics covered will include ideals, factorisation, the theory of field extensions, finite fields, polynomial rings in several variables, and the theory of modules.

In addition to the lecture notes, the following will cover much of the material we will be studying.

1. M Artin, Algebra, 1991

Rings are contexts in which it makes sense to add and multiply. For example,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$ ,  $\mathbb{C}$ , polynomials, functions  $\{0, 1\} \rightarrow \mathbb{R}$ , and  $\mathbb{Z}/n\mathbb{Z}$  are rings. The goals of this course include

1. to unify arguments that apply in all of the above contexts, and
2. to study relationships between different rings.

The applications of rings include

1. number theory, by studying extensions of  $\mathbb{Z}$  in which particular Diophantine equations have solutions, for example  $n = x^2 + y^2 = (x + iy)(x - iy)$  to study solutions in  $\mathbb{Z}\{i\}$  and pass to result about  $\mathbb{Z}$ ,
2. algebraic geometry, by the study of zero sets of polynomials in several variables via rings of functions, and
3. topology, by cohomology classes of topological spaces.

## 1 Basic definitions and examples

### 1.1 Rings

Recall the definition of a commutative ring.

**Definition 1.1.** A **commutative ring with identity**  $R$  is a set together with two binary operations  $+, \cdot : R \times R \rightarrow R$ , addition and multiplication, and two distinguished elements  $0_R$  and  $1_R$  such that the following holds.

1. The operation  $+$  makes  $R$  into an abelian group with identity  $0_R$ .
  - (a) For all  $r \in R$ ,  $0_R + r = r + 0_R = r$  (additive identity).
  - (b) For all  $r, s, t \in R$ ,  $(r + s) + t = r + (s + t)$  (associativity of  $+$ ).
  - (c) For all  $r, s \in R$ ,  $r + s = s + r$  (commutativity of  $+$ ).
  - (d) For all  $r \in R$ , there exists  $-r \in R$  such that  $r + (-r) = (-r) + r = 0_R$  (additive inverses).
2. The operation  $\cdot$  is associative and commutative with identity  $1_R$ .
  - (a) For all  $r \in R$ ,  $1_R \cdot r = r \cdot 1_R = r$  (multiplicative identity).
  - (b) For all  $r, s, t \in R$ ,  $(r \cdot s) \cdot t = r \cdot (s \cdot t)$  (associativity of  $\cdot$ ).
  - (c) For all  $r, s \in R$ ,  $r \cdot s = s \cdot r$  (commutativity of  $\cdot$ ).
3. Multiplication distributes over addition.
  - (a) For all  $r, s, t \in R$ ,  $r \cdot (s + t) = r \cdot s + r \cdot t$  and  $(s + t) \cdot r = s \cdot r + t \cdot r$  (distributivity of  $\cdot$  over  $+$ ).

There is some redundancy here, of course. I have written things this way so that one obtains the definition of a noncommutative ring simply by removing the condition that multiplication is commutative. In this course, however, all rings will be commutative.

**Proposition 1.2.** Let  $R$  be a ring. Then for all  $r \in R$ ,  $r \cdot_R 0_R = 0_R$ .

*Proof.*  $r \cdot_R 0_R = r \cdot_R (0_R +_R 0_R) = r \cdot_R 0_R +_R r \cdot_R 0_R$ . Thus  $0_R = -(r \cdot_R 0_R) +_R (r \cdot_R 0_R) = -(r \cdot_R 0_R) +_R (r \cdot_R 0_R +_R r \cdot_R 0_R) = r \cdot_R 0_R$ .  $\square$

Some people require  $0_R \neq 1_R$  in  $R$ .

**Proposition 1.3.** If  $0_R = 1_R$ , then  $R = \{0_R\}$ .

*Proof.*  $0_R = r \cdot_R 0_R = r \cdot_R 1_R = r$ .  $\square$

When it is clear from the context what ring we are working with, we will write  $0_R$  and  $1_R$  as 0 and 1,  $a +_R b$  as  $a + b$  and  $a \cdot_R b$  as  $ab$ .

**Definition 1.4.** A ring  $R$  is a **field** if  $R \neq \{0_R\}$  and every nonzero element of  $R$  has a multiplicative inverse, that is for every  $r \in R \setminus \{0_R\}$  there exists  $r^{-1} \in R$  such that  $rr^{-1} = r^{-1}r = 1_R$ .

We do not consider the zero ring  $\{0_R\}$  to be a field. We have seen many examples of rings at this point. The sets  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$ ,  $\mathbb{C}$  are all rings with their usual notion of addition and multiplication. All of them but  $\mathbb{Z}$  are in fact fields. As another example, we have the ring  $\mathbb{Z}/n\mathbb{Z}$  of integers modulo  $n$ . Let  $n \in \mathbb{Z}_{>0}$ , and recall that  $a$  and  $b$  are said to be **congruent modulo  $n$**  if  $a - b$  is divisible by  $n$ . It is easy to check that this is an equivalence relation on  $\mathbb{Z}$ . Moreover, since any  $a \in \mathbb{Z}$  can uniquely be written as  $qn + r$  with  $q, r \in \mathbb{Z}$  and  $0 \leq r < n$ , the set  $\{[0]_n, \dots, [n-1]_n\}$  is a complete list of the equivalence classes under this relation, where  $[a]_n$  denotes the set of all integers congruent to  $a \pmod n$ . We denote this  $n$ -element set by  $\mathbb{Z}/n\mathbb{Z}$ , and we can define addition and multiplication in  $\mathbb{Z}/n\mathbb{Z}$  by setting  $[a]_n + [b]_n = [a + b]_n$  and  $[a]_n [b]_n = [ab]_n$ . This defines a ring structure on  $\mathbb{Z}/n\mathbb{Z}$  once one checks that it is well-defined. This is the first example of a general construction of the quotient of a ring by an ideal we will define later.

Lecture 2  
Monday  
08/10/18

## 1.2 Polynomial rings

A very important class of rings that we will study are the polynomial rings. Let  $R$  be any ring. Then we can form a new ring  $R[X]$ , called the **ring of polynomials in  $X$  with coefficients in  $R$** . Informally, a polynomial in  $R[X]$  is a finite sum of the form  $r_0 + \dots + r_n X^n$  for some  $n \in \mathbb{Z}_{\geq 0}$  and  $r_i \in R$ . If  $n > m$ , we consider  $r_0 + \dots + r_n X^n$  to represent the same polynomial of  $R[X]$  as  $s_0 + \dots + s_m X^m$  if  $r_i = s_i$  for  $i \leq m$  and  $r_i = 0_R$  for  $i > m$ . That is, you can pad out a polynomial with terms of the form  $0_R X^i$  without changing it. From a formal standpoint, it is better to define a polynomial to be an infinite sum  $\sum_{i=0}^{\infty} r_i X^i$  for  $r_i \in R$  in which all but finitely many  $r_i$  are zero. This makes it easier to define addition and multiplication. The **degree** of such an expression is the largest  $i$  such that  $r_i$  is nonzero. We add and multiply in  $R[X]$  just as we would any other polynomials, by

$$\left( \sum_{i=0}^{\infty} r_i X^i \right) +_{R[X]} \left( \sum_{i=0}^{\infty} s_i X^i \right) = \sum_{i=0}^{\infty} (r_i +_R s_i) X^i,$$

$$\left( \sum_{i=0}^{\infty} r_i X^i \right) \cdot_{R[X]} \left( \sum_{i=0}^{\infty} s_i X^i \right) = \sum_{i=0}^{\infty} \left( \sum_{j=0}^i (r_j \cdot_R s_{i-j}) \right) X^i.$$

What about polynomial rings in more than one variable? Since the construction of polynomial rings takes an arbitrary ring as input, one can iterate it. Start with a ring  $R$ , and consider first the ring  $R[X]$  and then the ring  $(R[X])[Y]$ . An polynomial of this has the form  $\sum_{i=0}^{\infty} \left( \sum_{j=0}^{\infty} r_{ij} X^j \right) Y^i$  for  $r_{ij} \in R$ . On the other hand, we can consider the ring  $(R[Y])[X]$ , whose polynomials have the form  $\sum_{i=0}^{\infty} \left( \sum_{j=0}^{\infty} s_{ij} Y^j \right) X^i$  for  $s_{ij} \in R$ . Alternatively, we could consider the ring  $R[X, Y]$  whose polynomials are formal expressions of the form  $\sum_{i=0}^{\infty} \left( \sum_{j=0}^{\infty} r_{ij} X^i \right) Y^j$  with only finitely many nonzero coefficients  $r_{ij}$  and define addition and multiplication in the usual way. It is not hard to see that all three approaches yield the same ring. There is a bijection between these expressions. We will therefore primarily use notation like  $R[X, Y]$  for polynomial rings in multiple variables, but we will occasionally need to know that this is the same as  $(R[X])[Y]$  or  $(R[Y])[X]$ . The identification we have made here is an example of an isomorphism of rings, a notion we will make precise later.

### 1.3 Subrings and extensions

**Definition 1.5.** Let  $R$  be a ring. A subset  $S$  of  $R$  is a **subring** of  $R$  if

1.  $0_R, 1_R, -1_R \in S$ .
2.  $S$  is closed under  $+_R$  and  $\cdot_R$ , so if  $r, s \in S$ , then so are  $r +_R s$  and  $r \cdot_R s$ .

Subrings inherit the additive and multiplicative structures from the ring that contains them, and are thus themselves rings.

**Example.**  $\mathbb{Z}$  is a subring of  $\mathbb{R}$ , which is itself a subring of  $\mathbb{C}$ .

It is easy to see that the intersection of two subrings of  $R$ , or even an arbitrary collection of subrings of  $R$ , is also a subring of  $R$ .

**Definition 1.6.** Let  $S \subseteq R$  be a subring of a ring  $R$ , and let  $\alpha$  be an element of  $R$ . We can then form a subring  $S[\alpha]$  of  $R$ , called the **subring of  $R$  generated by  $\alpha$  over  $S$** , consisting of all elements of  $R$  that can be expressed as  $r_0 + \cdots + r_n \alpha^n$  for some  $n \in \mathbb{Z}^*$ , and  $r_i \in S$ .

This operation is known as **adjoining** the element  $\alpha$  to the ring  $S$ . An alternative way of defining the ring  $S[\alpha]$  is to note that it is the smallest subring of  $R$  containing  $S$  and  $\alpha$ . In one direction, any such subring contains every expression of the form  $r_0 + \cdots + r_n \alpha^n$ , with  $r_i \in S$ , so any subring of  $R$  containing  $S$  and  $\alpha$  contains  $S[\alpha]$ . One can thus construct  $S[\alpha]$  as the intersection of every subring of  $R$  containing  $S$  and  $\alpha$ . Since the intersection of any collection of subrings of  $R$  is a subring of  $R$  it is clear that this intersection is equal to  $S[\alpha]$  as defined above.

**Example.** Let  $i$  denote a square root of  $-1$  in  $\mathbb{C}$ .  $\mathbb{Z} \subseteq \mathbb{C}$  and  $i$  form  $\mathbb{Z}[i]$ . Note  $-1 = i^2 = i^6 = i + i^3 + i^{10}$ .

**Proposition 1.7.** Every element of  $\mathbb{Z}[i]$  can be uniquely expressed as  $a + bi$  for  $a, b \in \mathbb{Z}$ .

**Example.** Given  $\sum_{n=0}^{\infty} a_n i^n$  with only finitely many  $a_n$  nonzero, set  $a = a_0 - a_2 + \dots$  and  $b = a_1 - a_3 + \dots$ . Then  $\sum_{n=0}^{\infty} a_n i^n = a + bi$ . For uniqueness, if  $a + bi = c + di$  in  $\mathbb{C}$  for  $a, b, c, d \in \mathbb{Z}$ , then  $a = c$  or  $b = d$ .

If  $\alpha$  is more complicated than the elements of  $\mathbb{Z}[\alpha]$  may well be harder to describe.

**Example.** If  $\alpha$  is the real cube root of 2, then every element of  $\mathbb{Z}[\alpha]$  can be uniquely expressed as  $a + b\alpha + c\alpha^2$  for  $a, b, c \in \mathbb{Z}$ .

**Example.** In  $\mathbb{Z}[\pi]$ , any element has a unique expression in the form  $\sum_{n=0}^{\infty} a_n \pi^n$  for all but finitely many  $a_n$  are zero. Suppose  $\sum_{n=0}^{\infty} a_n \pi^n = \sum_{n=0}^{\infty} b_n \pi^n$ , then  $0 = \sum_{n=0}^{\infty} (a_n - b_n) \pi^n$ . Since  $\pi$  is transcendental, this polynomial must be zero. Thus each  $a_n = b_n$ .

**Example.** The elements of  $\mathbb{Z}[\frac{1}{2}]$  can be expressed uniquely as  $a/b$ , where  $b$  is a power of 2 and  $a$  is odd unless  $b = 1$ .

**Example.** Let  $\alpha$  be a root of  $x^2 - \frac{1}{2}x + 1$ . Then  $\alpha^2 \in \mathbb{Z}[\alpha]$  and  $\alpha^2 = \alpha/2 - 1$ . Can show that every element of  $\mathbb{Z}[\alpha]$  can be expressed as  $a + b\alpha$  for  $a, b \in \mathbb{Z}[\frac{1}{2}]$ , but not every  $a + b\alpha$  arises  $a, b \in \mathbb{Z}[\frac{1}{2}]$ .

### 1.4 Integral domains and rings of fractions

**Definition 1.8.** A **zero divisor** in a ring  $R$  is a nonzero element  $r$  of  $R$  such that there exists a nonzero  $s \in R$  with  $rs = 0$ . A ring  $R$  in which there are no zero divisors is called an **integral domain**.

**Example.**  $\mathbb{Z}$  is an integral domain and any subring of a field is an integral domain, but  $\mathbb{Z}/6\mathbb{Z}$  is not an integral domain, as  $[2][3]$  is zero modulo 6 even though neither  $[2]$  nor  $[3]$  is zero modulo 6.

If  $R$  is an integral domain, then we can form the field of fractions of  $R$  in analogy to the way we build  $\mathbb{Q}$  from  $\mathbb{Z}$ .

Lecture 3  
Wednesday  
10/10/18

**Definition 1.9.** Let  $R$  be an integral domain. The **field of fractions**  $K(R)$  is the set of equivalence classes of expressions of the form  $a/b$  for  $a, b \in R$ ,  $b \neq 0$ , where  $a/b \sim a'/b'$  iff  $ab' = a'b$ . We add and multiply elements of  $K(R)$  just as we do for fractions, by

$$\frac{a}{b} + \frac{a'}{b'} = \frac{ab' + a'b}{bb'}, \quad \frac{a}{b} \cdot \frac{a'}{b'} = \frac{aa'}{bb'}, \quad 0_{K(R)} = \frac{0_R}{1_R}, \quad 1_{K(R)} = \frac{1_R}{1_R}.$$

If  $a \neq 0$  in  $R$ , then  $b/a \in K(R)$ , so  $(a/b) \cdot (b/a) = ab/ba \sim 1/1$ .

Then  $K(R)$  is a field, and it contains  $R$  in a natural way as a subring if we identify  $r$  with  $r/1_R \in K(R)$ . The field  $K(R)$  is in some sense the smallest field containing  $R$  as a subring. When we talk about homomorphisms and isomorphisms, we will be able to state this more precisely. More generally, a subset  $S$  of  $R$  is a **multiplicative system** if  $1 \in S$  and  $0 \notin S$ , and  $S$  is closed under multiplication, that is if  $a, b$  are in  $S$  then so is  $ab$ . For any integral domain  $R$  and any multiplicative system  $S$ , we can define  $S^{-1}R \subseteq K(R)$  consisting of all fractions of the form  $a/b$  with  $b \in S$ . It is easy to see that this is closed under addition and multiplication, and defines a ring in between  $R$  and  $K(R)$ .

**Example.** If  $R = \mathbb{Z}$  and  $S$  is the set of powers of 2, then  $S^{-1}R = \mathbb{Z}[\frac{1}{2}]$ . On the other hand, if  $S$  is the set of odd integers, then  $S^{-1}R$  is the set of all rational numbers of the form  $a/b$  with  $b$  odd.

In general  $S^{-1}R$  is the smallest subring of  $K(R)$  containing  $R$  in which every element of  $S$  has a multiplicative inverse, that is  $b^{-1} \in S$  for all  $b \in S$ . The process of obtaining  $S^{-1}R$  from  $R$  is called **localisation** and is an extremely powerful tool. One can even make sense of it when  $R$  is not an integral domain, but one has to be more careful. The equivalence relation on fractions is trickier, for example. We will not discuss this in this course but it will be quite useful in future courses.

## 2 Homomorphisms, ideals, and quotients

### 2.1 Homomorphisms

Let  $R$  and  $S$  be rings. A ring homomorphism from  $R$  to  $S$  is, roughly, a way of interpreting elements of  $R$  as elements of  $S$ , in a way that is compatible with the addition and multiplication laws on  $R$  and  $S$ . More precisely is the following.

**Definition 2.1.** A function  $f : R \rightarrow S$  is a **ring homomorphism** if

1.  $f(1_R) = 1_S$ ,
2. for all  $r, r' \in R$ ,  $f(r +_R r') = f(r) +_S f(r')$ , and
3. for all  $r, r' \in R$ ,  $f(r \cdot_R r') = f(r) \cdot_S f(r')$ .

**Note.** If  $f$  is a homomorphism then  $f(0_R) = f(0_R + 0_R) = f(0_R) +_S f(0_R)$  gives  $f(0_R) = 0_S$ . Thus we do not need to require this as an axiom. On the other hand we do need to require  $f(1_R) = 1_S$ . For certain  $R, S$  one can construct examples of maps  $f : R \rightarrow S$  that satisfy properties 2 and 3 of the definition without satisfying property 1.

**Example.** If  $R$  is a subring of  $S$ , then the inclusion of  $R$  into  $S$ , such as  $\mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$ , is a homomorphism. This is just a fancy way of saying that the addition and multiplication on  $R$  are induced from the corresponding operations on  $S$ .

**Example.** The composition of two homomorphisms is a homomorphism, as is easily checked from the definitions.

**Example.** The map  $\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$  that takes an integer  $m$  into its congruence class modulo  $n$  is also a homomorphism. In fact, this is a special case of the following construction.

**Proposition 2.2.** Let  $R$  be any ring. Then there is a unique ring homomorphism  $f : \mathbb{Z} \rightarrow R$  such that

$$f(n) = \begin{cases} 1_R + \cdots + 1_R & n > 0 \\ -(1_R + \cdots + 1_R) & n < 0 \\ 0_R & n = 0 \end{cases}.$$

*Proof.* Let  $f : \mathbb{Z} \rightarrow R$  be a homomorphism. Then, directly from the definition, we have  $f(0) = 0_R$  and  $f(1) = 1_R$ . In particular for all  $n > 0$ ,  $f(n) = f(1 + \cdots + 1) = 1_R + \cdots + 1_R$ , where there are  $n$  copies of  $1_R$  in the sum. Moreover,  $0_R = f(n + (-n)) = f(n) + f(-n)$ , so  $f(-n) = -(1_R + \cdots + 1_R)$ . Thus  $f(n)$  is determined, for all  $n$ , completely by the fact that  $f$  is a homomorphism. In the converse direction, it is not hard to check that the map defined above is in fact a homomorphism.  $\square$

Thus, for any ring  $R$ , we can regard an integer as an element of  $R$  via this homomorphism.

**Definition 2.3.** A bijective homomorphism  $f : R \rightarrow S$  is called an **isomorphism**. Write  $S \cong R$  for  $S$  is isomorphic to  $R$ .

In this case one verifies easily that the inverse map  $f^{-1} : S \rightarrow R$  is also a bijective homomorphism.

## 2.2 Evaluation homomorphisms

Let  $R$  be a ring, and consider the ring  $R[X]$  of polynomials in  $X$  with coefficients in  $R$ . If  $s$  is an element of  $R$ , then we can define a homomorphism  $R[X] \rightarrow R$  by **evaluation at  $s$** . More precisely, given an element of  $R[X]$  of the form  $P(X) = r_0 + \cdots + r_n X^n$  for some  $n$  and  $r_i \in R$ . Then  $P(s)$  for  $s \in R$  is defined to be  $P(s) = r_0 + \cdots + r_n s^n \in R$ . Consider the map  $\phi_s : R[X] \rightarrow R$  that sends  $\phi_s(P)$  to  $P(s)$ . In effect, it substitutes  $s$  for  $X$ . It is easy to check that this is in fact a ring homomorphism. More generally, if  $R$  and  $S$  are rings and  $f : R \rightarrow S$  is a homomorphism, and  $s$  is an element of  $S$ , then we can define a map

$$\phi_{s,f} : R[X] \rightarrow S,$$

by setting

$$\phi_{s,f}(r_0 + \cdots + r_n X^n) = f(r_0) + \cdots + f(r_n) s^n.$$

That is, by applying  $f$  to the coefficients and substituting  $s$  for  $X$ . Again, this is clearly a homomorphism. The evaluation homomorphisms  $\phi_{s,f}$  are a fundamental property of polynomial rings. In some sense, they are the reason polynomial rings are worth studying. In fact, the ring  $R[X]$  is uniquely characterised by the fact that homomorphisms from  $R[X]$  to  $S$  are in bijection with pairs  $(s, f)$ , where  $f : R \rightarrow S$  is a homomorphism and  $s$  is an element of  $S$ .

## 2.3 Images, kernels, and ideals

**Definition 2.4.** Let  $f : R \rightarrow S$  be a homomorphism. The **image** of  $f$  is the subset  $Im(f) = \{f(r) \mid r \in R\} \subseteq S$ . The **kernel** of  $f$  is the subset  $Ker(f) = \{r \in R \mid f(r) = 0\} \subseteq R$ .

The image of a homomorphism  $f : R \rightarrow S$  is easily seen to be a subring of  $S$ .

**Example.** If  $R$  is a subring of  $S$ ,  $f : R \rightarrow S$  is the inclusion and  $s$  lies in  $S$ , then the image of the map  $\phi_{s,f} : R[X] \rightarrow S$  is precisely the subring  $R[s]$  of  $S$ .

By contrast, the kernel of a homomorphism  $f$  is almost never a subring of  $R$ . For instance, subrings contain the identity. However, it is an ideal of  $R$ .

**Definition 2.5.** A nonempty subset  $I$  of  $R$  is an **ideal** of  $R$  if  $I$  is closed under addition, that is for all  $i, j \in I$ ,  $i + j \in I$ , and for all  $i \in I$ ,  $r \in R$ ,  $ri \in I$ .

Then one can verify, directly from the definition, that the kernel of any homomorphism  $f : R \rightarrow S$  is an ideal of  $R$ .

**Note.** Any ideal of  $R$  contains  $0_R$ , and conversely the subset  $\{0_R\}$  of  $R$  is an ideal, called the **zero ideal**. A homomorphism  $f : R \rightarrow S$  is injective iff its kernel is the zero ideal. Forward direction is easy. Conversely, if  $f(x) = f(y)$ ,  $f(x - y) = 0$ , so  $x - y \in \text{Ker}(f)$ . If  $\text{Ker}(f) = \{0\}$ ,  $x = y$ .

The kernel of the homomorphism  $\mathbb{Z} \rightarrow R$  is either the zero ideal, or the ideal of multiples of  $n$  in  $\mathbb{Z}$  for some positive  $n$ . We say that  $R$  has characteristic zero or characteristic  $n$ , respectively. If not zero, the characteristic of  $R$  is the smallest  $n$  such that the sum of  $n$  copies of  $1_R$  is equal to zero.

## 2.4 Ideals: examples and basic operations

If  $r$  is an element of  $R$ , then any ideal containing  $R$  contains any multiple  $sr$  of  $R$ , for any  $r$  in  $S$ . Conversely, one checks easily that the set  $\{sr \mid s \in R\}$  is an ideal of  $R$ . It is known as the **ideal of  $R$  generated by  $r$** , and denoted  $\langle r \rangle$ . An ideal generated by one element in this way is called a **principal ideal**.

**Note.** The ideal generated by  $1_R$ , or more generally by any element of  $R$  with a multiplicative inverse, is all of  $R$ . This ideal is called the **unit ideal** of  $R$ .

**Proposition 2.6.**  $R$  is a **field** iff the only ideals of  $R$  are the zero ideal  $\{0\}$  and the unit ideal  $R$ .

*Proof.* If  $R$  is a field, let  $I \subseteq R$  be a nonzero ideal. There exists  $r \in I \neq 0$ . Then for all  $s \in R$ ,  $(sr^{-1})(r) \in I$ , so  $s \in I$  for all  $s \in R$ . Conversely, if  $R$  has only zero ideal, unit ideal, let  $r \in R \neq 0$ , let  $I = \{sr \mid s \in R\}$ . This is an ideal not zero ideal, so it is all of  $R$ . In particular,  $1 \in I$ , so there exists  $s \in R$  such that  $sr = 1$ .  $\square$

More generally is the following.

**Definition 2.7.** If  $S$  is a subset of elements of  $R$ , then any ideal containing  $S$  consists of all elements of  $R$  the form  $r_0s_0 + \cdots + r_ns_n$  for some  $n \in \mathbb{Z}_{\geq 0}$ ,  $r_i \in R$ , and  $s_i \in S$ . The intersection of all these ideals is an ideal of  $R$ , known as the **ideal of  $R$  generated by  $S$** , and denoted  $\langle S \rangle$ . It is also the smallest ideal of  $R$  containing  $S$ .

If  $S$  has one element,  $\langle S \rangle$  is a principal ideal. We will show soon that any ideal of  $\mathbb{Z}$  is a principal ideal, as is any ideal of the ring  $k[X]$  for any field  $k$ . On the other hand, there are rings in which not every ideal is principal.

**Example.** The ideal  $\langle X, Y \rangle$  of  $k[X, Y]$  is not a principal ideal.

Given ideals  $I$  and  $J$  there are several ways to create new ideals.

1. If  $I, J$  are ideals, then the intersection  $I \cap J$  is an ideal. If  $I$  and  $J$  are given by generators, it might be hard to find generators for the intersection. Certainly it is not enough to intersect the generating sets.
2. The union of ideals is not usually an ideal. Taking  $R = \mathbb{Z}$ ,  $\langle 3 \rangle \cup \langle 5 \rangle$  contains 3, 5 but not  $3 + 5$ .
3. If  $I, J$  are ideals, then the sum  $I + J$  is an ideal, which are all expressions of the form  $i + j$  for  $i \in I$ ,  $j \in J$ . It is the smallest ideal containing both  $I$  and  $J$ , and also the ideal generated by  $I \cup J$ .
4. If  $I, J$  are ideals, the product  $I \cdot J$  or  $IJ$  is the ideal generated by elements of the form  $ij$  for  $i \in I$ ,  $j \in J$ . This may be strictly larger than the set of such products.

**Example.** Consider the product of the ideals  $I = \langle X, Y \rangle$  and  $J = \langle Z, W \rangle$  in  $R = k[X, Y, Z, W]$  for  $k$  a field. The product  $IJ = \langle XZ, XW, YZ, YW \rangle$  contains  $XZ + YW$ , but the latter is not a product of an element in  $I$  with an element in  $J$ .

**Note.** Let  $I, J$  be general ideals. The product of  $I$  and  $J$  is always contained in the intersection of  $I$  and  $J$ , but the two need not be equal, even in simple rings like  $\mathbb{Z}$ .  $\langle 3 \rangle \cdot \langle 3 \rangle = \langle 9 \rangle \subseteq \mathbb{Z}$  and  $\langle 3 \rangle \cap \langle 3 \rangle = \langle 3 \rangle$ .

## 2.5 Quotients

Let  $R$  be a ring and let  $I$  be an ideal of  $R$ . If  $x, y$  are elements of  $R$ , we say that  $x$  is **congruent to  $y$  modulo  $I$**  if  $x - y$  is in  $I$ . This is an equivalence relation on  $R$ . We denote the equivalence class of  $r$  by  $r + I$ , or the alternative notations  $[r]_I, \bar{r}$ . It is the set  $\{r + s \mid s \in I\}$ . Let  $R/I$  denote the set of equivalence classes on  $R$  modulo  $I$ . This set has the natural structure of a ring. The additive and multiplicative identities are  $0_R + I$  and  $1_R + I$ , respectively, and addition and multiplication are defined by  $(r + I) + (s + I) = (r + s) + I$  and  $(r + I) \cdot (s + I) = (rs + I)$  respectively. One has to check that these are well-defined, but this is not difficult. The ring  $R/I$  is called the **quotient** of  $R$  by the ideal  $I$ .

**Example.** If  $R = \mathbb{Z}$  and  $I$  is the ideal generated by  $n$ , then  $R/I$  is the ring  $\mathbb{Z}/n\mathbb{Z}$  that we have already seen.

**Note.** There is a **reduction modulo  $I$**  or **natural quotient** homomorphism  $R \rightarrow R/I$  defined by taking  $r$  to  $r + I$ . This homomorphism is surjective with kernel  $I$ .

We then have the following.

**Proposition 2.8** (Universal property of the quotient). Let  $I \subseteq R$  be an ideal and let  $f : R \rightarrow S$  be a homomorphism, and suppose that the kernel of  $f$  contains  $I$ . Then there is a unique homomorphism  $\bar{f} : R/I \rightarrow S$  such that for all  $r \in R$ ,  $\bar{f}(r + I) = f(r)$ .

*Proof.*  $\bar{f}$  is necessarily unique, as every element of  $R/I$  has the form  $r + I$  for some  $r$ . It thus suffices to show that it is well-defined and gives a homomorphism. If  $r + I = r' + I$ , then  $r - r' \in I$ , so  $f(r - r') = 0$  gives  $f(r) = f(r')$ . Thus  $\bar{f}$  is well-defined. Checking that it is a homomorphism follows from  $f$  is a homomorphism.  $\square$

**Note.** The kernel of  $\bar{f}$  in the above proposition is just the image of the kernel of  $f$  in  $R/I$ . If the kernel of  $f$  is equal to  $I$ , this image is the zero ideal and  $\bar{f}$  is injective. In particular, any homomorphism of  $R$  to  $S$  can be thought of as an isomorphism of some quotient of  $R$  with a subring of  $S$ .

**Example.** Let  $R \subseteq S$  be a subring,  $\alpha \in S$ , and  $\iota : R \rightarrow S$  be the inclusion map. Recall that we have an evaluation at  $\alpha$  by  $\phi_{\iota, \alpha} : R[X] \rightarrow S$ . Image of this is  $R[\alpha]$ . Let  $I = \text{Ker}(\phi_{\iota, \alpha})$ . Then  $\phi_{\iota, \alpha}$  descends to a map  $\phi_{\iota, \alpha} : R[\alpha]/I \rightarrow S$  that is injective with image  $R[\alpha]$ . So  $R[\alpha]$  is isomorphic to a quotient of  $R[X]$ .

Lecture 5  
Monday  
15/10/18

## 2.6 Prime and maximal ideals

**Definition 2.9.** An ideal  $I$  of  $R$  is **prime** if the quotient  $R/I$  is an integral domain. It is **maximal** if  $R/I$  is a field.

**Note.** As fields are integral domains, every maximal ideal is prime. The converse need not hold, of course. The zero ideal in  $\mathbb{Z}$  is prime but not maximal.

**Proposition 2.10.** An ideal  $I$  is prime iff for every pair of elements  $s, r$  in  $R$  such that  $rs$  is in  $I$ , either  $r$  is in  $I$  or  $s$  is in  $I$ .

*Proof.* This is just a restatement of the definition.  $R/I$  integral domain iff for all whenever two elements  $r + I$  and  $s + I$  in  $R/I$  satisfy  $(r + I)(s + I) = 0 + I$  in  $R/I$ , either  $r + I = 0 + I$  or  $s + I = 0 + I$  in  $R/I$ . This is the same as saying  $rs$  lies in  $I$  iff either  $r$  or  $s$  lies in  $I$ .  $\square$

**Proposition 2.11.** An ideal  $I$  is maximal iff the only ideals of  $R$  containing  $I$  are  $I$  and the unit ideal  $R$ .

This justifies the name maximal for such ideals.

*Proof.* First suppose that  $R/I$  is a field. Recall that  $R/I$  is a field iff only ideals of  $R/I$  are  $\{0\}$  and  $R/I$ . Given an ideal  $J \subseteq R/I$ , let  $\tilde{J}$  be the preimage of  $J$  under  $R \rightarrow R/I$ .  $\tilde{J}$  is an ideal containing  $I$  and contained in  $R$ . Then  $J$  is either the zero ideal of  $R/I$ , in which case  $\tilde{J}$  is contained in, and thus equal to,  $I$ , or  $J$  is all of  $R/I$ , in which case  $\tilde{J}$  contains  $I$  and an element of  $1_R + I$ , so  $\tilde{J}$  contains  $1_R$  and is thus the unit ideal of  $R$ . Conversely, if the only ideals of  $R$  containing  $I$  are  $I$  and the unit ideal, then for any  $r$  in  $R \setminus I$ , the ideal of  $R$  generated by  $I$  and  $r$  contains  $1_R$ . We can thus write  $1_R = rs + i$ , where  $i \in I$  and  $s \in R$ . This means that  $s + I$  and  $r + I$  are multiplicative inverses of each other in  $R/I$ , so  $R/I$  is a field.  $\square$



### 3 Factorisation

In these notes  $R$  always denotes an integral domain.

#### 3.1 Divisibility, units, associates, and irreducibles

**Definition 3.1.** Let  $r, s$  be elements of  $R$ . We say  $r$  **divides**  $s$ , written  $r \mid s$ , if there exists  $r' \in R$  with  $rr' = s$ , or, equivalently,  $s$  lies in the principal ideal  $\langle r \rangle$  generated by  $r$ . An element  $r$  that divides  $1_R$  is called a **unit** of  $R$ , or, equivalently,  $\langle r \rangle = R$ .

The set of units in  $R$  forms a group under multiplication denoted  $R^*$ . For any element  $r \in R$  and any unit  $u$  of  $R$ , both  $u$  and  $ur$  divide  $r$ .

**Definition 3.2.** The set of elements of  $R$  of the form  $ur$ , with  $r \in R^*$  are called **associates** of  $R$ , that is  $r, r'$  are associates if  $r = ur'$  for a unit  $u \in R^*$ .

This implies  $r \mid r'$ , that is there exists  $u'$  with  $u'u = 1$  and  $u'r = r'$ .

**Note.** The principal ideals  $\langle r \rangle$  and  $\langle r' \rangle$  are equal iff  $r$  and  $r'$  are associates.

**Definition 3.3.** A nonzero element  $r$  of  $R$  is called **irreducible** if  $r$  is not a unit and the only elements of  $R$  that divide  $r$  are the units and the associates of  $r$ .

#### 3.2 Unique factorisation domains

An interesting question is when elements of rings admit unique factorisations into irreducibles. To that end we define the following.

**Definition 3.4.** A **unique factorisation domain** (UFD) is a ring  $R$  in which

1. every nonunit, nonzero element  $r \in R$  admits a factorisation as a finite product of irreducibles in  $R$ , and
2. if  $r = p_1 \dots p_n = q_1 \dots q_m \in R$  are two factorisations of  $r$  as products of irreducibles  $p_i, q_i$ , then  $n = m$  and, up to permuting the  $q_i$ , each  $q_i$  is an associate of  $p_i$ .

**Example.** Both conditions can fail.

1. There are certainly domains in which 1 can fail, although they are somewhat exotic. One example is to take the rational polynomial ring  $R = \mathbb{C}[X^{\mathbb{Q}}]$  with coefficients in  $\mathbb{C}$ , whose entries are finite formal sums  $\sum_{i=0}^N a_i X^{n_i}$  where the  $a_i$  are in  $\mathbb{C}$  and the  $n_i$  are nonnegative rational numbers  $\mathbb{Q}_{\geq 0}$ . Any element of  $R$  is a polynomial in  $X^{1/n}$  for some  $n$ . The element  $X$  of this ring is not a unit, and also not a finite product of irreducibles. In  $\mathbb{C}[X^{1/n}]$ ,  $X$  factors as  $(X^{1/n})^n$ .  $X$  has no factorisation into irreducibles in  $R$ . We will show later that a very mild finiteness condition on a domain  $R$ , the condition that  $R$  is Noetherian, actually guarantees that 1 holds.
2. Even if 1 holds, 2 often fails. The classic example of this is  $R = \mathbb{Z}[\sqrt{-5}]$ , in which  $2, 3, 1 + \sqrt{-5}, 1 - \sqrt{-5}$  are all irreducibles, none are associates of each other, yet  $(2)(3) = (1 + \sqrt{-5})(1 - \sqrt{-5})$ .

Another way to interpret condition 2 is as follows.

**Definition 3.5.** We say an element  $r$  of  $R$  is **prime** if the principal ideal  $\langle r \rangle$  of  $R$  is a prime ideal. In other words, for any  $s, s'$  in  $R$ , if  $r$  divides  $ss'$ , then  $r \mid s$  or  $r \mid s'$ .

**Lemma 3.6.** Prime elements are irreducible.

*Proof.* If  $r$  is prime and  $s$  divides  $r$ , we can write  $r = ss'$ . Then since  $r$  divides  $ss'$  we have that either  $r$  divides  $s$ , in which case  $rs'' = s$ , then  $ss's'' = s$  and  $s's'' = 1$ , so  $r$  is an associate of  $s$ , or  $r$  divides  $s'$ , in which case  $s' = rs''$ , then  $r = sr s''$  and  $ss'' = 1$ , so  $r$  is an associate of  $s'$  and  $s$  is a unit.  $\square$

The converse is not necessarily true, but we have the following observation as a criteria for  $R$  to be a UFD.

**Proposition 3.7.** Let  $R$  be a domain in which condition 1 holds. Then condition 2 above holds for  $R$  iff every irreducible element of  $R$  is prime.

*Proof.* First suppose condition 2 holds, and let  $r$  be an irreducible element of  $R$ . If  $r$  divides  $ab$ , we can write  $rs = ab$  for some  $s \in R$ . Expanding out  $s, a, b$  as products of irreducibles we see that  $r$  is an associate of some irreducible dividing  $a$  or  $b$ , so  $r$  is prime. Conversely, if every irreducible element of  $R$  is prime, and we have  $p_1 \dots p_n = q_1 \dots q_m$  products of irreducibles, then, since  $p_1$  is prime, it divides the product  $q_1 \dots q_m$  and is thus an associate of some  $q_i$ . We can thus cancel  $p_1$  from the left and  $q_i$  from the right after introducing a unit on one side. This is possible because  $R$  is an integral domain. Repeating the process we find that, up to reordering the terms and multiplying by units, the two expressions coincide.  $\square$

### 3.3 Principal ideal domains

**Definition 3.8.** An integral domain  $R$  is a **principal ideal domain** (PID) if every ideal of  $R$  is a principal ideal.

**Theorem 3.9.** Every PID is a UFD.

We first show 1. It is true for units trivially.

**Lemma 3.10.** Let  $R$  be a PID. Then every nonzero nonunit  $r \in R$  has a irreducible divisor.

*Proof.* Fix  $r = r_0 \in R$ . We first show  $r$  has an irreducible factor. If  $r_0$  is irreducible we are done. Otherwise  $r_0$  is not irreducible, we can choose an  $r_1$ , not a unit nor an associate of  $r_0$ , such that  $r_1$  divides  $r_0$ , so  $r_0 = r_1 s_1$  with  $r_1, s_1$  not units. If  $r_1$  is not irreducible we choose  $r_2$  similarly, and repeat. If this process ever terminates we have found an irreducible divisor of  $r$ . Suffices to show this terminates. Suppose it does not terminate. We obtain an increasing tower of ideals

$$\langle r_0 \rangle \subsetneq \langle r_1 \rangle \subsetneq \dots$$

Let  $I$  be the union of all these ideals generated by  $r_0, r_1, \dots$ . Then  $I$  is an ideal, so it is generated by some element  $s \in I$ . Thus  $s$  divides  $r_i$  for all  $i$ . On the other hand,  $s$  lives in some  $\langle r_j \rangle$ , so  $r_j$  divides  $s$ . Thus  $s$  is an associate of  $r_j$ , and therefore an associate of  $r_i$  for all  $i > j$ , that is  $I \subseteq \langle r_j \rangle$ . This contradicts our construction because  $\langle r_{j+1} \rangle \subseteq I$  and  $\langle r_{j+1} \rangle \neq \langle r_j \rangle$ .  $\square$

Thus  $r$  has an irreducible divisor  $s_0$ .

**Lemma 3.11.** Let  $R$  be a PID. Every nonzero nonunit  $r \in R$  is a finite product of irreducibles.

*Proof.* Consider  $rs_0^{-1}$ . If this is a unit we are done. If not let  $s_1$  be an irreducible divisor of  $rs_0^{-1}$ . If  $r(s_0 s_1)^{-1}$  is a unit we are done, otherwise repeat. We obtain a sequence of irreducibles  $s_0, s_1, \dots$  such that  $s_0 \dots s_i$  divides  $r$  for all  $i$ , so  $r = r_0 s_0 = r_0 r_1 s_1 = \dots$  with  $r_0, r_1, \dots$  irreducible. If this process ever terminates we are done. Suppose it does not. Then we have a strictly increasing tower of ideals

$$\langle r \rangle \subsetneq \langle s_0 \rangle \subsetneq \langle s_1 \rangle \subsetneq \dots$$

This cannot continue forever. Arguing as above we arrive at a contradiction.  $\square$

Now we show 2.

*Proof of Theorem 3.9.* It suffices to show that in a PID every irreducible is prime. Let  $r \in R$  be irreducible, and suppose that  $r$  divides  $st$ . Want  $r \mid s$  or  $r \mid t$ . Let  $q$  be a generator of the ideal  $\langle r, s \rangle$  of  $R$ , so  $\langle r, s \rangle = \langle q \rangle$ . Then  $q$  divides  $r$ , so either  $q$  is a unit or  $q$  is an associate of  $r$ . If  $q$  is an associate of  $r$ , then since  $q$  divides  $s$ ,  $r$  divides  $s$ . on the other hand, if  $q$  is a unit, then the ideal generated by  $r$  and  $s$  is the unit ideal and  $1 \in \langle r, s \rangle$ , so we can write  $1 = xr + ys$  for  $x, y$  elements of  $R$ . We then have  $t = xrt + yst$ , and since  $r$  divides both  $yrt$  and  $xrt$ ,  $r$  divides  $t$ .  $\square$

### 3.4 Euclidean domains

One technique for proving that rings are PIDs is Euclid's algorithm. We formalize this in an abstract setting as follows.

**Definition 3.12.** Let  $R$  be an integral domain.

1. A **Euclidean norm** on  $R$  is a function  $N : R \rightarrow \mathbb{Z}_{\geq 0}$  such that for all  $a, b \in R$ , with  $b \neq 0$ , there exists  $q, r \in R$  such that  $a = qb + r$ , and either  $r = 0$  or  $N(r) < N(b)$ .
2. An integral domain  $R$  is called a **Euclidean domain** if there is a Euclidean norm on  $R$ .

**Theorem 3.13.** Any Euclidean domain is a PID.

*Proof.* Let  $R$  be a Euclidean domain,  $N$  be a Euclidean norm on  $R$ , and  $I \subseteq R$  a nonzero ideal of  $R$ . Let  $a \in I$  be a nonzero element such that  $N(a)$  is minimal, that is if  $b \in I$ ,  $b \neq 0$ , then  $N(b) \geq N(a)$ . Claim that  $I = \langle a \rangle$ . Let  $b \in I$ . Then there exists  $q, r$  such that  $b = aq + r$ , with either  $r = 0$  or  $N(r) < N(a)$ . So  $r = 0$  gives  $b = aq$ . Thus  $I = \langle a \rangle$ .  $\square$

*Proof.* Let  $R$  be a Euclidean domain,  $N$  be a Euclidean norm on  $R$ , and  $I \subseteq R$  be a nonzero ideal of  $R$ . Let  $n$  be the smallest integer such that there exists a nonzero element  $a \in I$  with  $N(a) = n$  minimal, that is if  $b \in I$  and  $b \neq 0$ , then  $N(b) \geq N(a)$ . Claim that  $I = \langle a \rangle$ . Then for any  $b \in I$ , we can write  $b = qa + r$  with  $N(r) < N(a)$  unless  $r = 0$ . But since  $N(a)$  is the smallest possible norm in  $I$ , we must have  $r = 0$ , so  $b = qa$ . Thus  $I$  is generated by  $a$  and we are done.  $\square$

### 3.5 Examples

**Example.**

1. The classic example of a Euclidean domain is  $\mathbb{Z}$ , with  $N(x) = |x|$  for  $x \in \mathbb{Z}$ .
2. The ring  $\mathbb{Z}[i]$  is a Euclidean domain, with  $N(z) = z\bar{z} = |z|^2$ , so  $N(x + yi) = |x + yi|^2 = x^2 + y^2$ . To see this, note that given  $a$  and  $b$  in  $\mathbb{Z}[i]$  for  $b \neq 0$ , set  $q' = a/b \in \mathbb{Q}[i]$ . Write  $q' = x' + iy'$  with  $x', y' \in \mathbb{Q}$ . Let  $x$  and  $y$  be the closest integers to  $x'$  and  $y'$ , such that  $|x - x'|, |y - y'| \leq 1/2$ , and set  $q = x + iy$  in  $\mathbb{Z}[i]$  and  $r = a - bq$ . Then

$$N(r) = |r|^2 = |a - bq|^2 = \left| a - b \left( \frac{a}{b} + (q - q') \right) \right|^2 = |b(q - q')|^2 = |b|^2 |q - q'|^2 \leq \frac{N(b)}{2}.$$

Similar arguments can be used to prove that  $\mathbb{Z}[\alpha]$  is a Euclidean domain for

$$\alpha = \sqrt{-2}, \quad \alpha = \frac{-1 + \sqrt{-3}}{2}, \quad \alpha = \frac{-1 + \sqrt{-7}}{2}.$$

Beyond this one needs other tricks and for most  $\alpha$  unique factorization fails.

3. A critical example is the polynomial ring  $K[X]$  for  $K$  a field. Here we can take  $N(P(X))$  to be the degree of  $P(X)$ . Then, given polynomials  $P(X), T(X) \in K[X]$  and  $T(X) \neq 0$ , we can use polynomial long division to write  $P(X) = Q(X)T(X) + R(X)$  for some  $Q(X)$  with the degree of  $R$  strictly less than that of  $T$ , unless  $T$  is constant, in which case we can make  $R = 0$ . To prove this, fix  $T(X)$ . If  $\deg(T(X)) = 0$ ,  $T(X)$  is constant, so  $T(X) = c \neq 0 \in K$ . Take  $Q(X) = c^{-1}P(X)$ , so  $R(X) = 0$ . Otherwise induct on  $\deg(P(X))$ . If  $\deg(P(X)) < \deg(T(X))$ , set  $R(X) = P(X)$  and  $Q(X) = 0$ . Suppose the claim is true for polynomials of degree  $n$  and  $P(X)$  has degree  $n + 1$ , so

$$P(X) = \sum_{i=0}^{n+1} a_i X^i, \quad T(X) = \sum_{i=0}^d b_i X^i,$$

for  $d < n + 1$ . Then  $S(X) = P(X) - (a_{n+1}/b_d)X^{n+1-d}T(X)$  has degree  $n$ . By inductive hypothesis there exists  $Q(X), R(X)$  with  $\deg(R(X)) < \deg(T(X))$  such that

$$S(X) = Q(X)T(X) + R(X) \quad \implies \quad P(X) = \left( \frac{a_{n+1}}{b_d} X^{n+1-d} + Q(X) \right) T(X) + R(X).$$

Later, will show if  $R$  UFD, then  $R[X]$  is also a UFD.

## 4 Chinese remainder theorem

In elementary number theory, let  $m_1, m_2 \in \mathbb{Z}$  be relatively prime and  $a_1, a_2 \in \mathbb{Z}$ . Then there exists  $a \in \mathbb{Z}$  such that

$$a \equiv a_1 \pmod{m_1}, \quad a \equiv a_2 \pmod{m_2}.$$

Moreover,  $a$  is unique up to congruence modulo  $m_1 m_2$ . Question is given ideals  $I_1, \dots, I_r$  and  $a_1, \dots, a_r \in \mathbb{R}$ , when can we find a  $a \in R$  with  $a \in a_1 + I_1, \dots, a_r + I_r$ ?

### 4.1 Products

**Definition 4.1.** Let  $R_1, \dots, R_n$  be rings. The **direct product**  $R \times \dots \times R_n$  is a ring whose elements are  $n$ -tuples  $(r_1, \dots, r_n)$  with  $r_i \in R_i$  for all  $i$ . The addition and multiplication are given componentwise.

$$(r_1, \dots, r_n) + (r'_1, \dots, r'_n) = (r_1 + r'_1, \dots, r_n + r'_n), \quad (r_1, \dots, r_n)(r'_1, \dots, r'_n) = (r_1 r'_1, \dots, r_n r'_n).$$

**Note.** The product comes with natural homomorphisms for all  $i$ ,  $\pi_i$ , **projection** onto the  $i$ -th factor, defined by

$$\pi_i(r_1, \dots, r_n) = r_i : R_1 \times \dots \times R_n \rightarrow R_i,$$

and the following universal property.

**Theorem 4.2** (Universal property of the product). Let  $S, R_1, \dots, R_n$  be any rings. For any homomorphisms  $f_1 : S \rightarrow R_1, \dots, f_n : S \rightarrow R_n$ , there exists a unique homomorphism  $f : S \rightarrow R_1 \times \dots \times R_n$  such that  $\pi_i \circ f = f_i$  for all  $i$ .

*Proof.* Given  $f_i$ , the homomorphism  $f$  is defined by  $f(s) = (f_1(s), \dots, f_n(s))$ . Then  $(\pi_i \circ f)(s) = f_i(s)$ . For uniqueness, if  $(\pi_i \circ g)(s) = f_i(s)$  for all  $i$ , then  $g(s) = (f_1(s), \dots, f_n(s)) = f(s)$ .  $\square$

More generally, if  $I$  is any index set, and for each  $i \in I$  we have a ring  $R_i$ , we can define the product  $\prod_i R_i$ . An element  $r$  of this product is a choice, for each  $i \in I$ , of an element of  $R_i$ . We write such an element as  $(r_i)_{i \in I}$ . For each  $j \in I$  we have a map  $\pi_j : \prod_i R_i \rightarrow R_j$  given by  $\pi_j((r_i)_{i \in I}) = r_j$ . Such a product satisfies a very similar universal property. For any collection  $f_i : S \rightarrow R_i$  of maps for each  $i \in I$ , we get a unique map  $f : S \rightarrow \prod_i R_i$  such that  $\pi_j \circ f = f_j$ .

### 4.2 The Chinese remainder theorem

Let  $R$  be a ring, and let  $I_1, \dots, I_r$  be a finite collection of ideals of  $R$ . We have the natural maps  $R \rightarrow R/I_1, \dots, R \rightarrow R/I_r$ , which are surjective with kernel  $I_j$ . Consider the product map

$$R \rightarrow \frac{R}{I_1} \times \dots \times \frac{R}{I_r}.$$

It is easy to see that the kernel of this map is the set of  $r \in R$  such that  $r$  maps to zero in  $R/I_j$  for all  $j$ . That is, the kernel is the intersection  $I_1 \cap \dots \cap I_r$ . Call this ideal  $J$ . We thus have an injective embedding

$$\frac{R}{J} \hookrightarrow \frac{R}{I_1} \times \dots \times \frac{R}{I_r}.$$

A natural question to ask is, what can we say about the image? In other words, given congruence classes modulo  $I_1, I_2, \dots$ , when is there a single element of  $R$  that lives in all those congruence classes simultaneously?

**Note.** Because the above map is injective, if one such element exists, then there is a unique congruence class modulo  $J$  that satisfies all of the required congruences.

Of course, without further hypotheses we cannot expect this map to be surjective. Think about what happens when  $I_1 = I_2$ , for instance. Nonetheless, we have the following.

**Definition 4.3.** We will say  $I_1, \dots, I_r$  are **pairwise relatively prime** if for each  $i \neq j$ , the sum  $I_i + I_j$  is the unit ideal in  $R$ .

(TODO Exercise: if  $R = \mathbb{Z}$ , then  $I_i = \langle n_i \rangle$ , and  $\{I_i\}$  is pairwise relatively prime iff for all  $i \neq j$ ,  $n_i$  and  $n_j$  are relatively prime)

**Theorem 4.4.** Let  $R$  be a ring and  $I_1, \dots, I_r$  be pairwise relatively prime ideals. Then the natural map

$$\frac{R}{J} \hookrightarrow \frac{R}{I_1} \times \cdots \times \frac{R}{I_r}$$

is an isomorphism.

*Proof.* We have to prove it is surjective. Fix any tuple  $(c_1, \dots, c_r)$  of elements of  $R$ . We need to find  $c \in R$  such that  $c \in c_i + I_i$  for all  $i$ . It suffices to construct, for each  $i$ , an element  $e_i$  of  $R$  such that  $e_i \equiv 1 \pmod{I_i}$  and  $e_i \equiv 0 \pmod{I_j}$  for  $i \neq j$ . Suppose we have such an element. Then the element  $c = c_1 e_1 + \cdots + c_r e_r$  is such that  $c \equiv c_j \pmod{I_j}$  for all  $j$ . Given  $i, j$  with  $i \neq j$ , we know that  $I_i + I_j$  is the unit ideal. That is, we can write  $a_{ij} + b_{ij} = 1$  with  $a_{ij} \in I_i$  and  $b_{ij} \in I_j$ . Then  $a_{ij} \equiv 1 \pmod{I_j}$  and  $a_{ij} \equiv 0 \pmod{I_i}$  as an element of  $R/I_1 \times \cdots \times R/I_r$ , so  $a_{ij}$  has zero in the  $i$ -th place and one in the  $j$ -th place. Then for any  $i$  we can take  $e_j = \prod_{i \neq j} a_{ij}$  and  $e_j \equiv 1 \pmod{I_j}$  and  $e_j \equiv 0 \pmod{I_i}$  for all  $i \neq j$ , so  $e_j$  has one only in the  $j$ -th place. So  $R \rightarrow R/I_1 \times \cdots \times R/I_r$  is surjective. The result follows.  $\square$

### 4.3 Examples

When  $R = \mathbb{Z}$ , then every ideal is principal, so we can write  $I_j = \langle n_j \rangle$  for all  $j$ . The condition that  $I_i + I_j$  is the unit ideal becomes the condition that  $n_i \in \mathbb{Z}$  are pairwise relatively prime. In this case the ideal  $J$  is generated by the product  $n$  of the  $n_i$ . Specialising, we find the version of the Chinese remainder theorem from elementary number theory.

**Theorem 4.5.** If  $\{n_j \in \mathbb{Z}\}$  is a finite collection of pairwise relatively prime integers, and  $n$  is their product, then for any  $c_1, \dots, c_r \in \mathbb{Z}$ , there exists  $c \in \mathbb{Z}$  unique up to congruence modulo  $n$  such that  $c$  is congruent to  $c_i \pmod{n_i}$  for all  $i$ .

Now let  $K$  be a field and take  $R = K[X]$ . If  $c_1, \dots, c_r \in K$  are distinct elements of  $K$ , the ideals  $I_i = \langle X - c_i \rangle \subseteq R$  are such that  $I_i + I_j = \langle X - c_i \rangle + \langle X - c_j \rangle$  contains  $c_i - c_j \in K^*$ , so contains 1. That is,  $I_i + I_j$  is the unit ideal in  $R$  and the ideals  $I_i$  are pairwise relatively prime. Moreover, for each  $i$ ,  $I_i$  is the kernel of the evaluation map  $f_i : R \rightarrow K$  by that takes  $P(X)$  to  $P(c_i)$ . Let  $f : R \rightarrow K^r = K \times \cdots \times K$  by  $P(X) \rightarrow (P(c_1), \dots, P(c_r))$ . Then the following diagram commutes.

$$\begin{array}{ccc} R & \xrightarrow{f} & K^r \\ \downarrow & & \uparrow \sim \\ \frac{R}{J} & \xrightarrow{\sim} & \frac{R}{I_1} \times \cdots \times \frac{R}{I_r} \end{array}$$

Chinese remainder theorem gives that  $f$  is surjective. We thus have an isomorphism of  $R/I_i$  with  $K$  that takes  $P(X)$  to  $P(c_i)$  for all polynomials  $P$ . We thus obtain the following.

**Theorem 4.6.** For any  $c_1, \dots, c_n \in K$ , there is a polynomial  $P(X)$  in  $R$ , unique up to congruence modulo  $(x - a_1) \cdots (x - a_n)$  such that  $P(a_i) = c_i$  for all  $i$ .

## 5 Fields and field extensions

Next we will use  $K[X]$  is a PID for  $K$  a field to study fields systematically.

## 5.1 Prime fields

Let  $K$  be a field. We have a unique ring homomorphism  $\iota : \mathbb{Z} \rightarrow K$  by  $n \geq 0 \mapsto n_K = 1_K + \cdots + 1_K$ . Let  $I$  be the kernel. Then  $\mathbb{Z}/I \hookrightarrow K$  so  $\mathbb{Z}/I$  is an integral domain, so  $I$  is a prime ideal. Thus  $I$  is either the zero ideal  $\{0\}$ , if  $K$  has characteristic zero, or the ideal  $\langle p \rangle$  for some prime  $p$  of  $\mathbb{Z}$ . In the former case  $I = \{0\}$ , the injection  $\mathbb{Z} \hookrightarrow K$  extends to an inclusion  $\mathbb{Q} \hookrightarrow K$  sending  $a/b \mapsto (\iota a)(\iota b^{-1}) = a_K/b_K$ . In the latter case  $I = \langle p \rangle$ , we get an injection  $\mathbb{Z}/p\mathbb{Z} \hookrightarrow K$ , which we often denote  $\mathbb{F}_p$  when we think of it as a field. Upshot is that every field  $K$  contains exactly one of  $\mathbb{Q}, \mathbb{F}_p$ , for  $p$  prime, in exactly one way depending on its characteristic. This field is called the **prime field** of  $K$ , and it is contained in  $K$  in a unique way.

## 5.2 Field extensions

The prime fields are in some sense the smallest possible fields. Once we know they exist, it makes sense to study fields by studying pairs  $K, L$  of fields such that  $K \subseteq L$  of fields, trying to relate  $L$  to  $K$ .

**Definition 5.1.** A **field extension** is a pair of fields  $K, L$  with  $K \subseteq L$ , and is often denoted  $L/K$ .

**Note.** Such an inclusion of fields  $L/K$  makes  $L$  into a  $K$ -vector space, that is a vector space over  $K$ .

**Definition 5.2.** We say that a field extension  $L/K$  is **finite** if  $L$  is finite-dimensional as a  $K$ -vector space. If this is the case, the **degree** of such an extension is the dimension of  $L$  as a  $K$ -vector space  $\dim_K L$ , and is denoted  $[L : K]$ .

**Proposition 5.3.** Let  $K \subseteq L \subseteq M$  be fields. Then  $M/K$  is finite iff  $M/L$  and  $L/K$  are both finite. If this is the case then  $[M : K] = [M : L][L : K]$ .

*Proof.* First suppose that  $M/K$  is finite. Then  $L$  is a  $K$ -subspace of  $M$ , so finite dimensional as a  $K$ -vector space. Moreover, there exists a  $K$ -basis  $m_1, \dots, m_r$ , and this basis spans  $M$  over  $K$  and thus also over  $L$ . Thus  $M$  is finite-dimensional as an  $L$ -vector space, so  $M/L$  is finite. Conversely, suppose  $L/K, M/L$  are finite. Let  $e_1, \dots, e_n$  be a  $K$ -basis for  $L$ , and let  $f_1, \dots, f_m$  be an  $L$ -basis for  $M$ . Then claim that

$$e_1 f_1, \dots, e_1 f_m, \dots, e_n f_1, \dots, e_n f_m$$

is a  $K$ -basis for  $M$ . Every element  $x$  of  $M$  can be expressed uniquely as  $c_1 f_1 + \cdots + c_m f_m$  with  $c_i \in L$ . Each  $c_i$  in turn can be expressed as  $d_{1,i} e_1 + \cdots + d_{n,i} e_n$  with  $d_{j,i} \in K$ . Thus we can express  $x$  as

$$d_{1,1} e_1 f_1 + \cdots + d_{n,1} e_n f_1 + \cdots + d_{1,m} e_1 f_m + \cdots + d_{n,m} e_n f_m.$$

In particular the set  $\{e_i f_j\}$  for  $1 \leq i \leq n$  and  $1 \leq j \leq m$  spans  $M$  over  $K$ . In this case the degree of  $L$  over  $K$  is  $n$  and the degree of  $M$  over  $L$  is  $m$ , so it remains to show that  $\{e_i f_j\}$  is linearly independent over  $K$ . Suppose we have elements  $d_{i,j}$  of  $K$  such that  $\sum_{i,j} d_{i,j} e_i f_j = 0$ . Then, regrouping, we find that  $\sum_j (\sum_i d_{i,j} e_i) f_j = 0$  is an  $L$ -linear combination of the  $f_j$  that is zero. Since the  $f_j$  are linearly independent over  $L$  we must have  $\sum_i d_{i,j} e_i = 0$  for all  $j$ . Since the  $e_i$  are linearly independent over  $K$  we must have  $d_{i,j} = 0$  for all  $i, j$ .  $\square$

## 5.3 Extensions generated by one element

Let  $L/K$  be a field extension, and let  $\alpha$  be an element of  $L$ .

**Definition 5.4.** We let  $K(\alpha)$  denote the subfield of  $L$  consisting of all elements of  $L$  that can be expressed in the form  $P(\alpha)/Q(\alpha)$ , where  $P$  and  $Q$  are polynomials with coefficients in  $K$  and  $Q(\alpha)$  is not zero. This is the smallest subfield of  $L$  containing  $K$  and  $\alpha$ .

Recall that if  $R, S$  are rings,  $f : R \rightarrow S$  is a homomorphism, and  $\alpha \in S$ , then have  $\phi_{f,\alpha} : R[X] \rightarrow S$  by  $\phi_{f,\alpha}(\sum_{i=1}^n r_i X^i) = \sum_{i=1}^n f(r_i) \alpha^i$ . We have a natural map  $K[X] \rightarrow K(\alpha) \subseteq L$ , inclusion on  $K$ , that takes a polynomial  $P(X)$  to  $P(\alpha)$ . It is a ring homomorphism. Let  $I$  be the kernel of this homomorphism. We then get an injection of  $K[X]/I$  into the field  $K(\alpha)$ . Thus  $K[X]/I$  is an integral domain, so  $I$  is a prime ideal of  $K[X]$ . Since  $K[X]$  is a PID, every nonzero prime ideal is maximal. (TODO Exercise) There are

thus two cases. In the first  $I$  is the zero ideal that is not maximal. That is, there is no nonzero polynomial  $Q$  in  $K[X]$  such that  $Q(\alpha)$  is zero in  $L$ . We say that  $\alpha$  is **transcendental** over  $K$  in this case. In the second  $I$  is an ideal  $\langle Q \rangle$  for  $Q \in K[X]$  a nonzero irreducible polynomial that is a maximal ideal of  $K[X]$ . In this case we say  $\alpha$  is **algebraic** over  $K$ .

**Definition 5.5.**  $K(X)$  is the **field of rational functions** on  $X$ ,

$$K(X) = \left\{ \frac{P(X)}{Q(X)} \mid P, Q \in K[X], Q \neq 0 \right\} / \sim.$$

Assume first that  $\alpha$  is transcendental over  $K$ , that is  $I = \{0\}$ . Recall  $I = \{P(X) \in K[X] \mid P(\alpha) = 0\}$ . So in this case there is no nonzero polynomial  $P \in K[X]$  with  $P(\alpha) = 0$ . In this case the map taking  $P(X)$  to  $P(\alpha)$  is an injection of  $K[X]$  into  $K(\alpha) \subseteq L$ . In particular every nonzero element of  $K[X]$  gets sent to a nonzero, hence invertible, element of  $L$ . Thus the map from  $K[X]$  to  $L$  extends to an injective map from the field of fractions of  $K[X]$ , which we denote  $K(X)$ , to  $L$ . This map takes  $P(X)/Q(X)$  to  $P(\alpha)/Q(\alpha)$ . By definition of  $K(\alpha)$ , this map is surjective so the image of this map is  $K(\alpha)$ . In particular  $K(X)$  and  $K(\alpha)$  are isomorphic. Thus the following diagram holds.

$$\begin{array}{ccc} K(X) & \xrightarrow[\sim]{g} & K(\alpha) \\ \uparrow \subseteq & \nearrow f & \uparrow \subseteq \\ K[X] & & \end{array} \quad f : P(X) \mapsto P(\alpha) \quad g : \frac{P(X)}{Q(X)} \mapsto \frac{P(\alpha)}{Q(\alpha)}$$

**Note.** In this case  $K(\alpha)$  is infinite dimensional as a  $K$ -vector space. It contains a subspace isomorphic to  $K[X]$ , for instance.

If  $\alpha$  is algebraic over  $K$ , then  $I$  is a nonzero maximal ideal of the PID  $K[X]$ , so it is generated by a single irreducible polynomial  $Q(X)$  in  $K[X]$ . As a consequence, since the units in  $K[X]$  are just the constant polynomials, the polynomial  $Q(X)$  is well-defined up to a constant factor. It is called the **minimal polynomial** of  $\alpha$ . By definition, it divides every polynomial  $P(X)$  such that  $P(\alpha) = 0$ . Since  $\langle Q(X) \rangle$  is maximal, the ring  $K[X] / \langle Q(X) \rangle$  is a field. Recall that for any  $P \in K[X]$ , can write  $P(X)$  uniquely as  $A(X)Q(X) + R(X)$  with  $\deg(R) < \deg(Q)$ . So  $1, \dots, X^{\deg(Q)-1}$  are a  $K$ -basis of  $K[X] / \langle Q(X) \rangle$ . So its dimension as a  $K$ -vector space is equal to the degree of  $Q(X)$ . The map  $K[X] \rightarrow K(\alpha) \subseteq L$  descends to an injection of  $K[X] / \langle Q(X) \rangle$  into  $L$ . Since its image is a subfield of  $K(\alpha)$  containing  $K$  and  $\alpha$ , this map is an isomorphism of  $K(\alpha)$  with  $K[X] / \langle Q(X) \rangle$ . Thus in this case the extension  $K(\alpha)/K$  is a finite extension, of degree equal to the degree of  $Q(X)$ . Thus the following diagram holds.

$$\begin{array}{ccc} K[X] & & L \\ \uparrow \subseteq & \searrow f & \uparrow \subseteq \\ \frac{K[X]}{\langle Q(X) \rangle} & \xrightarrow[\sim]{g} & K(\alpha) \end{array} \quad f : P(X) \mapsto P(\alpha) \quad g : [R(X)]_{\langle Q(X) \rangle} \mapsto R(\alpha)$$

To summarise, extend  $K$  by a single element by

1. building  $K[X]$ , and
2. either passing to field of fractions  $K(X)$  to form a transcendental extension, or choosing an irreducible polynomial  $Q$  to form an algebraic extension  $K[X] / \langle Q(X) \rangle$ .

Slightly informally, instead of  $K[X] / \langle Q(X) \rangle$ , we sometimes write  $K(\alpha)$ , where  $\alpha$  is a root of  $Q(X)$ .

**Definition 5.6.** An extension  $L/K$  is **algebraic** if every element of  $L$  is algebraic over  $K$ .

An observation is that if  $L/K$  is finite, then  $L/K$  is algebraic. Suppose not. Let  $\alpha \in L$  be transcendental over  $K$ .  $K[X]$  is a polynomial ring in  $K(\alpha)$  contained in  $L$ , so  $L/K$  is not finite.

**Corollary 5.7.** Let  $L/K$  be a field extension for  $\alpha, \beta \in L$  algebraic over  $K$ . Then  $\alpha + \beta$ ,  $\alpha\beta$  are algebraic over  $K$ .

*Proof.*  $[K(\alpha) : K] = \deg(\alpha)$  and  $[K(\alpha, \beta) : K(\alpha)] \leq \deg(\beta)$ , so  $[K(\alpha, \beta) : K] \leq (\deg(\alpha))(\deg(\beta))$ . Now  $K \subseteq K(\alpha + \beta) \subseteq K(\alpha, \beta)$ , so  $\deg(\alpha + \beta)$  over  $K$  is less than  $(\deg(\alpha))(\deg(\beta))$ . Similarly for  $\alpha\beta$ .  $\square$

**Corollary 5.8.** If  $L/K$ , then the subset  $L^{alg}$  of elements of  $L$  algebraic over  $K$  is a field.

*Proof.* If  $a_0 + \cdots + a_n \alpha^n = 0$  then  $a_0 (\alpha^{-1})^n + \cdots + a_n = 0$ .  $\square$

**Example.**  $\overline{\mathbb{Q}} \subseteq \mathbb{C}$  is the subfield of elements of  $\mathbb{C}$  that are algebraic over  $\mathbb{Q}$ .