

# M4P33 Algebraic Geometry

Lectured by Dr Genival Da Silva Jr  
Typeset by David Kurniadi Angdinata

Spring 2019

## Contents

<b>0</b>	<b>Introduction</b>	<b>3</b>
<b>1</b>	<b>Affine varieties</b>	<b>4</b>
<b>2</b>	<b>Projective varieties</b>	<b>8</b>
<b>3</b>	<b>Morphisms</b>	<b>11</b>

## 0 Introduction

I will not follow a particular book, but everything I am going to say will be contained in one of the following books.

Lecture 1  
Friday  
11/01/19

- I Shafarevich, Basic algebraic geometry, 1974
- R Hartshorne, Algebraic geometry, 1977
- J Harris, Algebraic geometry: a first course, 1992

# 1 Affine varieties

*Notation 1.1.*

- $R$  is a commutative ring with unity.
- $K$  is a field.
- $K[x_1, \dots, x_n]$  is the ring of polynomials in  $n$  variables.
- $\mathbb{A}^n$  is  $K^n$  as a set.

**Definition 1.2.** Let  $S \subseteq K[x_1, \dots, x_n]$  then

$$Z(S) = \{x \in \mathbb{A}^n \mid \forall f \in S, f(x) = 0\}$$

is called the **zero locus** of  $S$ . Subsets of  $\mathbb{A}^n$  that are of this form are called **affine varieties**.

*Remark 1.3.* Some authors call **algebraic set** the object  $Z(S)$ . We will not follow this notation.

**Example 1.4.**

- Single points  $p = (p_1, \dots, p_n)$ .  $p = Z(S)$  where  $S = \{x_1 - p_1, \dots, x_n - p_n\}$ .
- $\mathbb{A}^n = Z(0)$ .
- $\emptyset = Z(1)$ .
- Subspaces of  $\mathbb{A}^n = K^n$ .
- If  $X = Z(f_1, \dots, f_n) \subseteq \mathbb{A}^n$  and  $Y = Z(g_1, \dots, g_m) \subseteq \mathbb{A}^n$  are affine varieties then

$$X \times Y = Z(f_1, \dots, f_n, g_1, \dots, g_m) \subseteq \mathbb{A}^{n+m}$$

is a variety.

*Remark 1.5.* If  $S \subseteq K[x_1, \dots, x_n]$  and  $I = \langle S \rangle$  then  $Z(S) = Z(I)$ .

**Theorem 1.6** (Hilbert's basis theorem). *If  $R$  is Noetherian then  $R[x]$  is Noetherian.*

**Corollary 1.7.** *Every ideal in  $K[x_1, \dots, x_n]$  is finitely generated.*

**Definition 1.8.** Let  $X \subseteq \mathbb{A}^n$  then

$$I(X) = \{f \in K[x_1, \dots, x_n] \mid \forall x \in X, f(x) = 0\}.$$

**Example 1.9.**  $I(p) = I((p_1, \dots, p_n)) = \langle x_1 - p_1, \dots, x_n - p_n \rangle$ .

Goal is

$$\begin{array}{ccc} \{\text{affine varieties in } \mathbb{A}^n\} & \leftrightarrow & \{\text{ideals of } K[x_1, \dots, x_n]\} \\ X & \mapsto & I(X) \\ Z(J) & \leftarrow & J \end{array}.$$

$$Z(I(X)) = X \text{ but } I(Z(J)) \supseteq J.$$

**Example 1.10.**  $J = \langle x^2 + 1 \rangle \subseteq \mathbb{R}[x] = I(\emptyset) = I(Z(x^2 + 1))$ .

**Proposition 1.11.**

- If  $X \subseteq Y$  then  $I(Y) \subseteq I(X)$ . If  $I \subseteq J$  then  $Z(J) \subseteq Z(I)$ .
- $X \subseteq Z(I(X))$  and  $S \subseteq I(Z(S))$ .
- If  $X$  is affine then  $Z(J(X)) = X$ . If  $X = Z(S)$  then take  $Z$  of  $S \subseteq I(Z(S))$ .

**Example 1.12.** Let  $J \subseteq \mathbb{C}[x]$ .  $J = \langle f \rangle$ , where  $f = (x - x_1)^{k_1} \dots (x - x_n)^{k_n}$ .

**Definition 1.13.** Let  $I \subseteq K[x_1, \dots, x_n]$  be an ideal.

$$I \subseteq \sqrt{I} = \{f \in K[x_1, \dots, x_n] \mid \exists n \in \mathbb{N}, f^n \in I\}.$$

If  $\sqrt{I} = I$ , we say  $I$  is a **radical ideal**. (Exercise:  $\sqrt{I}$  is an ideal,  $I \subseteq \sqrt{I}$ , and  $\sqrt{I} = \bigcap_{p \text{ prime}} p$ )

**Theorem 1.14** (Hilbert's Nullstellensatz).  $I(Z(J)) = \sqrt{J}$ . If  $\sqrt{J} = J$  then

$$\begin{array}{ccc} \{\text{affine varieties}\} & \leftrightarrow & \{\text{radical ideals}\} \\ X & \mapsto & I(X) \\ Z(J) & \leftarrow & J \end{array}.$$

**Proposition 1.15.**

1.  $Z(S) \cup Z(T) = Z(ST)$ .
2.  $\bigcap_i Z(S_i) = Z(\bigcup_i S_i)$ .
3.  $Z(0) = \mathbb{A}^n$  and  $Z(1) = \emptyset$ .

*Proof.*

1. If  $p \in Z(S) \cup Z(T)$ , then  $f(p) = 0$  for  $f \in S$  or  $f \in T$ , so  $f(p) = 0$  for  $f \in ST$ , where

$$ST = \left\{ \sum_{i \in I, I \text{ finite}} s_i t_i \right\} \subseteq S \cap T,$$

with equality if  $S + T = R$ . If  $p \in Z(ST)$ , there exists  $f$  such that  $f(p) = 0$  for  $f \in S$  or  $f(p) = 0$  for  $f \in T$ , so  $p \in Z(S) \cup Z(T)$ .

□

**Definition 1.16.** The **Zariski topology** on  $\mathbb{A}^n$  is the topology generated by closed sets of the form  $Z(S)$ . By the above proposition this is a topology.

**Example 1.17.**  $\mathbb{A}^1$  is not Hausdorff.

**Definition 1.18.** A topological space  $X$  is **irreducible** if it cannot be expressed as a union  $X = A \cup B$ , where  $A$  and  $B$  are proper and closed subsets.  $\emptyset$  is not considered irreducible.

**Example 1.19.**  $\mathbb{A}^1$ .

**Example 1.20.** Any non-empty open set of irreducible  $X$  is dense and irreducible. Suppose  $A$  is open then  $X = A^c \cup \overline{A}$ . Since  $X$  is irreducible then  $A^c = X$ , a contradiction, or  $\overline{A} = X$ . Suppose  $A$  is reducible. Let  $A = (A \cap B) \cup (A \cap C)$ , where  $B$  and  $C$  are closed. Then  $X = A^c \cup (B \cup C)$ .  $A^c = X$  or  $B \cup C = X$ , which are contradictions.

**Example 1.21.** If  $A$  is irreducible then  $\overline{A}$  is also irreducible. Suppose  $\overline{A}$  is not irreducible.  $\overline{A} = (\overline{A} \cap B) \cup (\overline{A} \cap C)$ . Take  $\bigcap A$ ,  $A = (A \cap B) \cup (A \cap C)$ , a contradiction.

**Definition 1.22.** An affine variety is **irreducible** if it is irreducible as a topological space.

*Remark 1.23.* A **quasi-affine variety** is an open set of an affine variety.

**Proposition 1.24.**

1.  $I(X \cup Y) = I(X) \cap I(Y)$ .
2.  $Z(I(X)) = \overline{X}$  for any  $X \subseteq \mathbb{A}^n$ .

*Proof.*

1. If  $f \in I(X \cup Y)$  then  $f(p) = 0$  for all  $p \in X \cup Y$ , so  $f \in I(X)$  and  $f \in I(Y)$ .
2. We know that  $X \subseteq Z(I(X))$  hence  $\overline{X} \subseteq Z(I(X))$ . Now, let  $Y$  be a closed set containing  $X$ , that is  $X \subseteq Y$ . Then

$$I(Y) \subset I(X) \implies Z(I(X)) \subset Z(I(X)) = Y,$$

so any closed set containing  $Y$  contains  $Z(I(X))$ .

□

**Proposition 1.25.**  $X$  is irreducible if and only if  $I(X)$  is prime.

*Proof.*

$\implies$  Let  $f, g \in I(X)$ .

$$X \subseteq Z(fg) = Z(f) \cup Z(g) \implies X = (X \cap Z(f)) \cup (X \cap Z(g)).$$

$Z(f) \subseteq X$ , so  $f \in I(X)$ , or  $Z(g) \subseteq X$ , so  $g \in I(X)$ .

$\Leftarrow$  Exercise.

□

**Example 1.26.**  $\mathbb{A}^n$ .

**Definition 1.27.** If  $X \subseteq \mathbb{A}^n$ , the **coordinate ring** of  $X$  is

$$A(X) = \frac{A}{I(X)} = \frac{K[x_1, \dots, x_n]}{I(X)}.$$

**Example 1.28.** Let  $f \in K[x_1, \dots, x_n]$  be irreducible. If  $n = 3$ ,  $Z(f)$  is a surface. If  $n = 2$ ,  $Z(f)$  is a curve.

**Example 1.29.** Let  $y - x^2 \in K[x, y]$ . Then

$$\begin{aligned} A(X) &= \frac{K[x, y]}{\langle y - x^2 \rangle} \cong K[x, x^2] \rightarrow K[x] \\ \sum_{i,j} a_{ij} x^i x^{2j} &= \sum_{i,j} a_{ij} x^{2j+i} \mapsto \sum_n b_n x^n \end{aligned}$$

**Example 1.30.** Let  $xy - 1 \in K[x, y]$ . Then

$$A(X) = \frac{K[x, y]}{\langle xy - 1 \rangle} \cong K\left[x, \frac{1}{x}\right].$$

$A(X)$  cannot be  $K[x]$ .

**Definition 1.31.** A **Noetherian** topological space  $X$  is a topological space such that if

$$C_1 \supseteq C_2 \supseteq \dots$$

is a decreasing chain of closed sets then there is a  $k$  such that  $C_k = C_{k+1} = \dots$ .

**Example 1.32.**  $\mathbb{A}^n$ . Recall that if  $A \subset B$  then  $I(B) \subset I(A)$ . So using the definition above,

$$I(C_1) \subseteq I(C_2) \subseteq \dots$$

Since  $K[x_1, \dots, x_n]$  is Noetherian then  $I(C_i)$  stabilises. So  $I(C_k) = I(C_{k+1}) = \dots$ , but taking  $Z$ , we recover  $C_k$  so  $C_k$  stabilises as well.

Lecture 3  
Tuesday  
15/01/19

**Theorem 1.33.** *If  $X$  is Noetherian then any non-empty closed subset can be expressed as a finite union of irreducible closed sets  $X = Y_1 \cup \dots \cup Y_n$ . Moreover, if we require that  $Y_i \subseteq Y_j$  then this expression is unique.*

*Proof.* Let  $C$  be the collection of closed sets that do not satisfy that property. Let  $Y$  be a minimum closed inside  $C$ , in particular  $Y$  is reducible, so  $Y = Y' \cup Y''$ , for  $Y', Y''$  closed. Hence  $Y', Y'' \notin C$ , so they can be expressed as a finite union of irreducibles, a contradiction. If  $Y_i \not\subseteq Y_j$ , then suppose

$$Y_1 \cup \dots \cup Y_n = X_1 \cup \dots \cup X_n.$$

Then  $Y_1 \subset X_1 \cup X_n$ , in particular  $Y_1 = \bigcup_j (Y_1 \cap X_j)$ , so there is a  $j$  such that  $Y_1 \cap X_j = Y_1$ , so  $Y_1 \subset X_j$ . We can assume  $j = 1$  and repeat the same argument to find that  $Y_1 = X_1$ , so consider  $\overline{Y \setminus Y_1} = Y_2 \cup \dots \cup Y_n$ . But

$$Y_2 \cup \dots \cup Y_n = X_2 \cup \dots \cup X_n,$$

and the result follows by induction.  $\square$

**Corollary 1.34.** *Any affine variety in  $\mathbb{A}^n$  can be expressed equally as a union of irreducible algebraic varieties.*

**Definition 1.35.** The **dimension** of a topological space is the supremum of  $n$  where

$$Y_0 \subset \dots \subset Y_n$$

is a sequence of irreducible closed sets.

**Example 1.36.** Dimension of  $\mathbb{A}^1$  is one.

**Definition 1.37.** Let  $A$  be a ring and  $\mathfrak{p}$  be a prime ideal, then the **height** of  $\mathfrak{p}$  is the supremum of  $n$  where

$$\mathfrak{p}_1 \subset \dots \subset \mathfrak{p}_n \subset \mathfrak{p},$$

where  $\mathfrak{p}_i$  are prime. The **Krull dimension** of  $A$  is

$$\sup_{\mathfrak{p} \text{ prime}} \text{height}(\mathfrak{p}).$$

**Proposition 1.38.** *If  $Y$  is affine then  $\dim(Y) = \dim(A(Y))$ .*

*Proof.* Let  $C$  be a closed and irreducible set  $C \subset Y$ , then  $I(C) \supset I(Y)$ , then  $I(C)$  is prime.  $\square$

**Proposition 1.39.** *Let  $K$  be a field and  $B$  be an integral domain which is a finitely generated algebra, then*

- $\dim(B)$  is the transcendence degree of  $K(B)$  over  $K$ , and
- if  $\mathfrak{p} \subseteq B$  is prime, then

$$\text{height}(\mathfrak{p}) + \dim\left(\frac{B}{\mathfrak{p}}\right) = \dim(B).$$

*Proof.* Atiyah Macdonald chapter 11.  $\square$

**Proposition 1.40** (Krull Hauptidealsatz). *Let  $A$  be a Noetherian ring and  $f \in A$  not a zero divisor and not a unit. Then every prime ideal containing  $f$  has height one.*

*Proof.* Atiyah Macdonald page 122.  $\square$

**Proposition 1.41.** *A Noetherian integral domain  $A$  is a UFD if and only if every prime ideal  $I$  of height one is principal.*

**Theorem 1.42.** *An irreducible variety  $Y \subseteq \mathbb{A}^n$  has dimension  $n - 1$  if and only if  $Y = Z(f)$  where  $f$  is an irreducible polynomial in  $K[x_1, \dots, x_n]$ .*

*Proof.*

$\implies$  If  $Y$  has dimension  $n - 1$  then  $I(Y)$  has height one, by the above proposition  $I(Y) = \langle f \rangle$ , so  $Y = Z(f)$ .

$\impliedby$  Let  $I = I(Y)$  then  $I$  is prime, by the Krull Hauptidealsatz we have that  $I$  has height one, so  $\dim(Y) = n - 1$ .  $\square$

## 2 Projective varieties

**Definition 2.1.** The **projective space**  $\mathbb{P}^n$  is defined as

$$\mathbb{P}^n = \frac{\mathbb{A}^{n+1} \setminus \{0\}}{\{x \sim \lambda x \mid \lambda \in K^*\}}.$$

A point in  $\mathbb{P}^n$  is written as  $[a_0 : \dots : a_n] = \overline{(a_0, \dots, a_n)}$ .

**Definition 2.2.** A **graded ring**  $R$  is a ring together with a decomposition

$$R = \bigoplus_{d \geq 0} R_d,$$

where  $R_d$  are abelian groups and  $R_k \cdot R_t \subseteq R_{k+t}$ .

**Example 2.3.**  $K[x_0, \dots, x_n]$  is a graded ring, where  $R_d$  are monomials of degree  $d$ .

*Notation 2.4.* Let  $A$  be  $K[x_0, \dots, x_n]$  without the grading and  $S$  be  $K[x_0, \dots, x_n]$  as a graded ring.

**Definition 2.5.** An ideal  $I \subseteq S$  is **homogeneous** if

$$I = \bigoplus_{d \geq 0} (I \cap S_d).$$

If  $f = f_0 + \dots + f_d$ , then  $f_i \in I$ .

*Remark 2.6.*  $I$  is homogeneous if and only if  $I = \langle f_0, \dots, f_n \rangle$ , where  $f_i$  are homogeneous.

**Lemma 2.7.** If  $I, J$  are homogeneous then

1.  $I + J$  is homogeneous,
2.  $IJ$  is homogeneous,
3.  $I \cap J$  is homogeneous, and
4.  $\sqrt{I}$  is homogeneous.

*Proof.*

4. Let  $f = f_0 + \dots + f_d \in \sqrt{I}$  then

$$f^n = (f_0 + \dots + f_d)^n = f_d^n + \dots \in I \quad \implies \quad f_d^n \in I \quad \implies \quad f_d \in \sqrt{I},$$

so  $f - f_d \in \sqrt{I}$ , by induction  $f_i \in \sqrt{I}$ .

□

**Definition 2.8.** If  $f$  is homogeneous of degree  $k$  then

$$f(\lambda \cdot x) = \lambda^k \cdot f(x),$$

in particular  $f(x) = 0$  if and only if  $f(\lambda \cdot x) = 0$ , so it makes sense to define

$$Z(f) = \{x \in \mathbb{P}^n \mid f(x) = 0\}.$$

More generally, if  $I \subseteq S$  is a homogeneous ideal then

$$Z(I) = \{x \in \mathbb{P}^n \mid f \in I \text{ homogeneous, } f(x) = 0\}.$$

**Definition 2.9.** A subset  $X \subseteq \mathbb{P}^n$  is called a **projective variety** if  $X = Z(T)$  for some homogeneous ideal  $T$ .



**Proposition 2.10.**

- $Z(S) \cup Z(T) = Z(ST)$ .
- $\bigcap_{\alpha} Z(S_{\alpha}) = Z(\bigcup_{\alpha} S_{\alpha})$ .
- $Z(0) = \mathbb{P}^n$  and  $Z(1) = \emptyset$ .

**Definition 2.11.** We define the **Zariski topology** on  $\mathbb{P}^n$  by taking closed sets to be  $Z(T)$  for some  $T$ .

**Definition 2.12.**

- A projective variety is **irreducible** if it is an irreducible topological space.
- An open subset of a projective variety is called a **quasi-projective variety**.
- The **dimension** of a projective variety is its dimension as a topological space.
- If  $T \subseteq S$  then

$$I(T) = \langle f \in S \mid f \text{ homogeneous, } \forall p \in T, f(p) = 0 \rangle.$$

**Definition 2.13.** If  $X$  is a projective variety the **homogeneous coordinate ring** is

$$S(X) = \frac{S}{I(X)}.$$

**Definition 2.14.** If  $f \in S$  is linear and homogeneous, we call  $Z(f)$  a **hyperplane**.

**Proposition 2.15.**

$$\begin{aligned} \phi_i : U_i = \mathbb{P}^n \setminus Z(x_i) &\rightarrow \mathbb{A}^n \\ [x_0 : \dots : x_n] &\mapsto \left( \frac{x_0}{x_i}, \dots, \frac{x_n}{x_i} \right) \end{aligned}$$

is a homeomorphism in the Zariski topology.

*Proof.* Let  $\phi = \phi_0$  and  $U = U_0$ , let  $C \subseteq \mathbb{A}^n$  be a closed set then we claim that  $\phi^{-1}(C)$  is closed. Indeed, let  $C = Z(S)$ , then  $\phi^{-1}(C) = Z(S') \cup U$  where

$$S' = \left\{ x_0^d \cdot f\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \mid f \in S \right\}.$$

Similarly, let  $A \subseteq U$  is closed, we claim that  $\phi(A)$  is closed. Let  $\overline{A}$  be its closure in  $\mathbb{P}^n$ , then  $\overline{A} = Z(B)$ , so  $\phi(A) = Z(B')$  where

$$B' = \{f(1, x_1, \dots, x_n) \mid f \in B\}.$$

So we conclude that  $\phi$  is a homeomorphism. □

Note that  $\langle 1 \rangle = S$  and  $\langle x_0, \dots, x_n \rangle \subsetneq S$  map to  $\emptyset$  under  $Z$ . So in order to have a one-to-one correspondence we need the following.

- $Z(I) = \emptyset$  if and only if  $\sqrt{I} \supseteq \langle x_0, \dots, x_n \rangle$ . If we consider  $Z(I)$  in  $\mathbb{A}^{n+1}$ , note that  $x \in Z(I)$  if and only if  $\lambda x \in Z(I)$ . So  $Z(I) = \emptyset$  if and only if  $Z(I) \subseteq \{0\}$ . So  $\sqrt{I} \supseteq \langle x_0, \dots, x_n \rangle$ .
- $I(Z(J)) = \sqrt{J}$  if  $Z(J) \neq \emptyset$ , since  $I(Z(J)) = I(Z_a(J)) = \sqrt{J}$ .

**Corollary 2.16.**

$$\begin{aligned} \{ \text{projective varieties} \} &\longleftrightarrow \{ \text{homogeneous radical ideals not } \langle x_0, \dots, x_n \rangle \}, \\ \{ \text{irreducible projective varieties} \} &\longleftrightarrow \{ \text{homogeneous radical prime ideals} \}. \end{aligned}$$

**Example 2.17.**  $\mathbb{P}^n$  is irreducible.

Lecture 5  
Monday  
21/01/19

**Proposition 2.18.**

- $\mathbb{P}^n$  is Noetherian, that is satisfies the descending chain condition.
- Every projective variety can be written as a unique union of irreducible projective varieties. We call **irreducible components** the irreducible varieties in that decomposition.

**Theorem 2.19.** Let  $Y \subseteq \mathbb{P}^n$  be an irreducible projective variety. Then

$$\dim(S(Y)) = \dim(Y) + 1.$$

*Proof.* Let

$$\begin{aligned} \phi_i : U = \mathbb{P}^n \setminus Z(x_i) &\rightarrow \mathbb{A}^n \\ [x_0 : \dots : x_n] &\mapsto \left( \frac{x_0}{x_i}, \dots, \frac{x_n}{x_i} \right), \end{aligned}$$

and  $Y_i = \phi_i(Y \cap U_i)$ . Let

$$\begin{aligned} K[x_1, \dots, x_n] &\rightarrow (S(Y)_{x_i})_0 \\ f(x_1, \dots, x_n) &\mapsto \frac{x_i^{\partial f} f\left(\frac{x_1}{x_i}, \dots, \frac{x_n}{x_i}\right)}{x_i^{\partial f}}, \end{aligned}$$

then

$$A(Y_i) = \frac{K[x_1, \dots, x_n]}{I(Y_i)} \cong (S(Y)_{x_i})_0,$$

moreover  $S(Y)_{x_i} \cong A(Y_i)[x_i, x_i^{-1}]$ . So

$$\dim(S(Y)) = \dim(S(Y)_{x_i}) = \dim(A(Y_i)[x_i, x_i^{-1}]) = \text{tra}(K(Y_i)(x_i)) = \dim(Y_i) + 1.$$

Therefore if  $Y_i \neq \emptyset$ ,  $\dim(Y_i) = \dim(S(Y)) - 1$  for all  $i$ , but since  $U_i$  cover  $Y$  we have  $\dim(Y) = \max\{\dim(Y_i)\}$ . (Exercise: if  $\{U_n\}_n$  is a finite cover of a topological space  $Y$  then  $\dim(Y) = \max\{\dim(Y_i)\}$ ) Since  $\dim(Y_i)$  are the same if  $Y_i \neq \emptyset$ , we conclude that  $\dim(Y) = \dim(Y_d)$  for some  $d$ .  $\square$

**Proposition 2.20.** Every Noetherian topological space is compact.

*Proof.* Let  $X$  be a Noetherian topological space and let  $\{U_n\}$  be a cover of  $X$ . So consider  $C$ , the collection of the union of finitely many open sets of  $\{U_n\}$ . Since  $X$  is Noetherian  $C$  has a maximum element, say  $U_1 \cup \dots \cup U_n$ . If  $U_1 \cup \dots \cup U_n \subsetneq X$  then there is  $x \in X$  not in the union, and we can find another  $U_{\alpha_0} \ni x$ . But then

$$U_1 \cup \dots \cup U_n \cup U_{\alpha_0} \supsetneq U_1 \cup \dots \cup U_n,$$

a contradiction. So  $X = U_1 \cup \dots \cup U_n$ .  $\square$

**Corollary 2.21.**  $\mathbb{P}^n$ ,  $\mathbb{A}^n$ , affine varieties, and projective varieties are all compact in the Zariski topology.

**Definition 2.22.** A variety  $X$  is **complete** if for any other variety  $Y$ , the projection  $X \times Y \rightarrow Y$  is closed.

**Example 2.23.**  $\mathbb{P}^n$  is complete.  $\mathbb{A}^n$  is not complete.

Lecture 6  
Tuesday  
22/01/19

### 3 Morphisms

**Definition 3.1.** Suppose  $Y$  is a quasi-affine variety and  $p \in Y$ . We say that a function  $f : Y \rightarrow \mathbb{A}^1$  is **regular** at  $p$  if there are  $g, h \in K[x_1, \dots, x_n]$  and  $U \ni p$  such that  $f = g/h$  in  $U$  with  $h \neq 0$ . A function is **regular** if it is regular for every  $p \in Y$ .

**Example 3.2.** Local is not global. Let  $X = Z(x_1x_4 - x_2x_3) \subseteq \mathbb{A}^4$  and  $U = X \setminus Z(x_2, x_4)$ . Then

$$\begin{aligned} \phi : \quad U &\rightarrow \mathbb{A}^1 \\ (x_1, x_2, x_3, x_4) &\mapsto \begin{cases} \frac{x_1}{x_2} & x_2 \neq 0 \\ \frac{x_3}{x_4} & x_4 \neq 0 \end{cases} \end{aligned}$$

is a regular function.

**Definition 3.3.** Let  $Y$  be a quasi-projective variety,  $f : Y \rightarrow \mathbb{A}^1$ , and  $p \in Y$ . We say that  $f$  is **regular** at  $p$  if there are  $g, h$  homogeneous polynomials of the same degree and an open set  $U \ni p$  such that  $f = g/h$  on  $U$  and  $h \neq 0$ .

**Lemma 3.4.** A regular function is continuous.

*Proof.* It is enough to show that  $f^{-1}(p)$  is closed. Since  $f$  is regular  $f = g/h$  on some neighbourhood  $U$ , then  $f^{-1}(p) \cap U = Z(g - ph) \cap U$ .  $\square$

**Remark 3.5.** If  $X$  is irreducible then  $f = g$  on  $U \subseteq X$ , then  $f = g$  on  $X$ . Because the set where  $f - g = 0$  is closed and dense.

**Definition 3.6.** We will use the term **variety** to denote an affine, quasi-affine, projective, or quasi-projective variety.

**Definition 3.7.** A **morphism** is  $f : X \rightarrow Y$  if  $f$  is continuous and for every  $U \subseteq Y$  and every function  $g : U \rightarrow \mathbb{A}^1$  the composition  $g \circ f$  is regular.

**Remark 3.8.**

- Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  then the composition  $g \circ f$  of these two morphisms is the composition of  $f$  and  $g$  as functions.
- A morphism  $f : X \rightarrow Y$  is an **isomorphism** if there is a morphism  $g : Y \rightarrow X$  such that  $f \circ g = id$  and  $g \circ f = id$ .

**Definition 3.9.** Let  $X$  be a variety. Denote the set of all regular functions of  $X$  by  $\mathcal{O}(X)$ . If  $p \in X$  the **local ring** at  $p \in X$  is

$$\mathcal{O}_p = \varinjlim_{U \ni p} (\mathcal{O}(U)).$$

An element of  $\mathcal{O}_p$  is a pair  $(U, f)$ , where  $p \in U$  and  $f$  is regular at  $p$ , moreover  $(U, f) \sim (V, g)$  if  $f = g$  on  $U \cap V$ .

**Definition 3.10.** Let  $Y$  be an irreducible variety, the **function field**  $K(Y)$  of  $Y$  is the field whose elements are pairs  $(U, f)$  where  $U$  is open and  $f$  is regular on  $U$ , and

$$(U, f) + (V, g) = (U \cap V, f + g).$$

**Remark 3.11.**

- $K(Y)$  is indeed a field for if  $(U, f) \neq 0$  then  $U^{-1} = U \setminus Z(f)$ , so  $(U^{-1}, 1/f)$  is the inverse to  $(U, f)$ .
- $K(Y)$  is the quotient field of  $A(Y)$  or  $S(Y)$ .
- $\mathcal{O}(Y) \hookrightarrow \mathcal{O}_p \hookrightarrow K(Y)$  for all  $p \in Y$ .

**Theorem 3.12.** *If  $Y \subseteq \mathbb{A}^n$  is an irreducible affine variety with coordinate ring  $A(Y)$  then*

1.  $\mathcal{O}(Y) = A(Y)$ ,
2. for all  $p \in Y$ , let  $\mathfrak{m}_p = \{f \in A(Y) \mid f(p) = 0\}$  then we have a one-to-one correspondence

$$\{ \text{points of } Y \} \quad \rightsquigarrow \quad \{ \text{maximal ideals of } A(Y) \},$$

3. for all  $p \in Y$ ,  $\mathcal{O}_p \cong A(Y)_{\mathfrak{m}_p}$  and  $\dim(\mathcal{O}_p) = \dim(Y)$ , and
4.  $K(Y)$  is the quotient field of  $A(Y)$ .

*Proof.*

1. Notice that there is a natural map  $A \rightarrow \mathcal{O}(Y)$  with kernel  $I(Y)$ , so there is an injection  $A(Y) \hookrightarrow \mathcal{O}(Y)$ , that is

$$A(Y) \subseteq \mathcal{O}(Y) \subseteq \bigcap_{p \in Y} \mathcal{O}_p = \bigcap_{\mathfrak{m}_p} A(Y)_{\mathfrak{m}_p} = A(Y),$$

so  $A(Y) = \mathcal{O}(Y)$ .

2. We know that points of  $Y$  correspond to maximal ideals  $\mathfrak{m}_p \supseteq I(Y)$ . Taking the quotient, we get maximal ideals inside  $A(Y)$ .
3. There is a natural map  $A(Y)_{\mathfrak{m}_p} \rightarrow \mathcal{O}_p$ , which is injective by  $\alpha : A(Y) \hookrightarrow \mathcal{O}(Y)$ , and it is surjective by definition of  $\mathcal{O}_p$ . Moreover,  $\dim(\mathcal{O}_p) = \dim(A_p)_{\mathfrak{m}_p} = \text{height}(\mathfrak{m}_p) = \dim(Y)$ .
4. The quotient field of  $A(Y)$  is the quotient field of  $\mathcal{O}_p$  for all  $p$ , by 3, which is  $K(Y)$  by definition.

□

**Theorem 3.13.** *Let  $Y \subseteq \mathbb{P}^n$  be irreducible and projective. Then*

1.  $\mathcal{O}(Y) = K$ ,
2. for all  $p \in Y$ ,  $\mathfrak{m}_p$  as before,  $\mathcal{O}_p \cong (S(Y)_{\mathfrak{m}_p})_0$ , and
3.  $K(Y) \cong (S(Y)_{(0)})_0$ .

*Proof.* Recall that

$$\begin{aligned} \phi_i : U_i = \mathbb{P}^n \setminus Z(x_i) &\rightarrow \mathbb{A}^n \\ [x_0 : \cdots : x_n] &\mapsto \left( \frac{x_0}{x_i}, \dots, \frac{x_n}{x_i} \right) \end{aligned}$$

gives  $\phi_i^* : A(Y_i) \cong (S(Y)_{x_i})_0$  and  $Y_i = \phi_i(Y \cap U_i)$ .

1.  $K \subseteq \mathcal{O}(Y)$ . Take  $f \in \mathcal{O}(Y)$ , so  $f$  is regular at each  $Y_i$ , but  $\mathcal{O}(Y_i) \cong A(Y_i)$ , also by  $\phi_i^*$ ,  $A(Y_i) \cong (S(Y)_{x_i})_0$ . Thus  $f = g_i/x_i^{n_i}$ , where  $n_i = \deg(g_i)$ , in particular  $x_i^{n_i}f \in S(Y)_{n_i}$ . Now, set  $N \geq \sum_i n_i$ , then  $S(Y)_N \cdot f \subseteq S(Y)_N$ , so we can iterate this process to obtain  $S(Y)_N \cdot f^q \subseteq S(Y)_N$ . In particular  $x_0^N f \in S$ , hence  $S(Y)[f]$  is contained in  $x_0^{-N} S(Y)$ . Therefore  $f$  is integral since  $S(Y)[f]$  is finitely generated. There are  $a_i \in S$  such that  $f^k + a_1 f^{k-1} + \cdots + a_k = 0$ . Since  $f$  is homogeneous of degree zero we can take the constant terms of  $a_i$  and still have an equation, hence  $a_i \in K$ .
2. Let  $p \in Y$ , then  $p \in Y_i$ , by the previous theorem we know that  $\mathcal{O}_p \cong A(Y_i)_{\mathfrak{m}_p}$ . By  $\phi_i^*$ ,  $\mathcal{O}_p \cong ((S(Y)_{x_i})_{\mathfrak{m}_p})_0$ , but since  $x_i \notin \mathfrak{m}_p$ , hence  $\mathcal{O}_p \cong (S(Y)_{\mathfrak{m}_p})_0$ .
3. Recall that the quotient field of  $Y$  is  $K(Y) = K(Y_i)$ , but  $K(Y_i)$  is the quotient field of the coordinate ring  $A(Y_i)$ , by  $\phi_i^*$ , this is  $(S(Y)_{(0)})_0$ .

□

Lecture 8  
Monday  
28/01/19

**Proposition 3.14.** *Let  $X$  be an irreducible variety and  $Y$  be an irreducible affine variety, then we have a bijection*

$$\alpha : \text{Hom}(X, Y) \xrightarrow{\sim} \text{Hom}(A(Y), \mathcal{O}(X)),$$

*the set of morphisms from  $X$  to  $Y$  to the set of  $K$ -algebra homomorphisms.*

*Proof.* Given a morphism  $\phi : X \rightarrow Y$ , by definition of morphism,  $\phi$  takes regular functions at  $Y$  to regular functions at  $X$ . So if  $f \in A(Y)$  then  $\phi \circ f \in \mathcal{O}(X)$ . Conversely, let  $h : A(Y) \rightarrow \mathcal{O}(X)$  be a homomorphism of  $K$ -algebras. Recall that  $A(Y) = A/I(Y) = k[x_1, \dots, x_n]/I(Y)$ . Take  $\bar{x}_i \in A(Y)$  and let  $y_i = h(\bar{x}_i) \in \mathcal{O}(X)$  and define

$$\begin{aligned} \psi : X &\rightarrow \mathbb{A}^n \\ p &\mapsto (y_1(p), \dots, y_n(p)) \end{aligned}$$

We claim that  $\text{Im}(\psi) \subseteq Y$ , but since  $Y = Z(I(Y))$ , it is enough to show that if  $f \in I(Y)$  then  $f(\psi(p)) = 0$ .

$$f(\psi(p)) = f(y_1(p), \dots, y_n(p)) = f(h(\bar{x}_1(p)), \dots, h(\bar{x}_n(p))) = h(f(x_1, \dots, x_n))(p) = 0.$$

□

**Lemma 3.15.** *If  $X, Y$  are as before then  $\psi : X \rightarrow Y$  is a morphism if and only if  $\psi_i = x_i \circ \psi$  are regular functions.*

*Proof.* Suppose  $\psi_i$  are regular functions, then if  $p$  is a polynomial  $p \circ \psi$  is regular, but since regular functions are quotients of polynomials, we conclude that  $f \circ \psi$  is regular for any regular function  $f$ . □

**Corollary 3.16.** *If  $X, Y$  are affine then  $X \cong Y$  if and only if  $A(X) \cong A(Y)$ .*

**Corollary 3.17.** *The correspondence  $X \mapsto A(X)$  induces an arrow reversing correspondence between the category of affine varieties and the category of  $K$ -integral domains.*

Lecture 9 is a problem class.

Lecture 9  
Tuesday  
29/01/19