

# M3P20 Geometry I: Algebraic Curves

Lectured by Dr Mattia Talpo  
Typeset by David Kurniadi Angdinata

Autumn 2018

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Complex plane curves</b>	<b>5</b>
<b>3</b>	<b>Projective space</b>	<b>9</b>
<b>4</b>	<b>Projective curves</b>	<b>12</b>
<b>5</b>	<b>Affine vs projective plane curves</b>	<b>15</b>
<b>6</b>	<b>Points at infinity</b>	<b>17</b>

# 1 Introduction

This course is intended as a first course in algebraic geometry, the area of mathematics that studies spaces defined by polynomial equations using algebra. It will focus on one dimensional algebraic varieties. The reference books for the course are the following.

1. F Kirwan, Complex algebraic curves, 1992
2. W Fulton, Algebraic curves: an introduction to algebraic geometry, 1969

Geometry is the study of shapes in suitable spaces, such as sets of points on the real line  $\mathbb{R}$ , lines and circles in  $\mathbb{R}^2$ , spheres in higher dimensional Euclidean spaces  $\mathbb{R}^n$ , etc. One way to think about shapes is to see them as the locus of zeroes defined by

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid f(x_1, \dots, x_n) = 0\} \iff \{f(x_1, \dots, x_n) = 0\} \subset \mathbb{R}^n$$

for some suitable function  $f$ .

## Example 1.1.

1. Circles  $\{f_1(x, y) = x^2 + y^2 - R^2 = 0\}$  in  $\mathbb{R}^2$  for some  $R \in \mathbb{R}$ .
2. The unit square with vertices at  $\{(\pm 1, 0), (0, \pm 1)\}$  in  $\mathbb{R}^2$  defined by  $\{f_2(x, y) = |x| + |y| - 1 = 0\}$ .
3. Spheres  $\{f_3(x_1, \dots, x_n) = x_1^2 + \dots + x_n^2 - R^2\}$  in  $\mathbb{R}^n$  for some  $R \in \mathbb{R}$ .

**Remark 1.2.** Every subset  $S \subseteq \mathbb{R}^n$  is the zero-set of some function. We can define  $\chi_S(x) = 1$  if  $x \notin S$ ,  $\chi_S(x) = 0$  if  $x \in S$ .

The class of functions used to define our shapes has great consequences on their geometry. For the circle,  $f_1$  is a polynomial so that it is differentiable and also  $C^\infty$ . For the square,  $f$  is continuous but not differentiable at  $\{(0, \pm 1), (\pm 1, 0)\}$ , the vertices of the square. The function  $\chi_S$  is not even continuous, unless  $S$  is empty, or the whole  $\mathbb{R}^n$ . As these examples illustrate, an underlying principle is the following equivalence.

**Fact 1.** Regularity properties of  $f$  are regularity properties of  $\{f = 0\}$ .

Such shapes are called **algebraic varieties**. Their geometric properties are intimately related to the algebraic properties of the defining polynomial equations.

## Example 1.3.

1. Let  $f(x)$  be a polynomial. Then the zero set of  $f(x)$ ,  $\{f(x) = 0\} \subseteq \mathbb{R}$  is a finite set of points in  $\mathbb{R}$ , and every finite set of points arises in this manner.
2. The circle is an algebraic variety.
3. Spheres in higher dimensions are algebraic varieties.

## Exercise 1.

1. Is  $\mathbb{Z} \subseteq \mathbb{R}$  an algebraic variety?
2. Is the unit square an algebraic variety?

**Definition 1.4.** Let  $K$  be a field, such as  $K = \mathbb{Q}, \mathbb{R}, \mathbb{C}$ . For  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$  a multi-index, denote by  $|\alpha| = \sum_{i=1}^n \alpha_i$  and by  $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ , a **monomial**. A **polynomial of degree  $d$**  in  $n$  variables with coefficients in  $K$  is a finite sum.

$$P(x_1, \dots, x_n) = \sum_{\alpha \in \mathbb{N}^n} a_\alpha x^\alpha,$$

where  $a_\alpha \in K$ ,  $a_\alpha = 0$  for all  $|\alpha| > d$  and  $a_\alpha \neq 0$  for some  $\alpha$  with  $|\alpha| = d$ . The set of polynomials of arbitrary degree in  $n$  variables with coefficients in  $K$  is denoted  $K[x_1, \dots, x_n]$ .

**Example 1.5.** Let  $n = 3$ .  $P(x_1, x_2, x_3) = 3 + x_1^2 x_2 + x_3^{10}$  for  $\alpha = (2, 1, 0)$  and  $\alpha = (0, 0, 10)$  has degree ten.

**Exercise 2.**

1. Show that  $K[x_1, \dots, x_n]$  is a ring, and that if  $P, Q$  are polynomials of degrees  $p, q$  respectively, then the degree of  $\lambda P + \mu Q$  for  $\lambda, \mu \in K$  is at most  $\max\{p, q\}$ . Give an example of polynomials  $P, Q \in K[x]$  such that  $\deg(P + Q) < \max\{\deg(P), \deg(Q)\}$ .
2. Show that  $(P \cdot Q)(x_1, \dots, x_n) = P(x_1, \dots, x_n)Q(x_1, \dots, x_n)$  is a polynomial  $P \cdot Q \in K[x_1, \dots, x_n]$  with  $\deg(P \cdot Q) = \deg(P) + \deg(Q)$ . What if  $P = 0$ ? What is  $\deg(0)$ ?

**Definition 1.6.** An **affine plane curve** defined over  $K$  is

$$C = \{(x, y) \in K^2 \mid P(x, y) = 0\} \subset K^2,$$

where  $P \in K[x, y]$ . More generally, an **algebraic variety**  $V \subset K^n$  is a subset of  $K^n$  defined as the locus

$$\{f_1 = \dots = f_k = 0\} \subset K^n,$$

where  $f_1, \dots, f_k \in K[x_1, \dots, x_n]$  are polynomials in  $n$  variables with coefficients in  $K$ .

**Example 1.7.**

1. Let  $a, b, c \in \mathbb{R}$  with  $(a, b) \neq (0, 0)$ , and let

$$f(x, y) = ax + by + c.$$

Then  $\{(x, y) \in \mathbb{R}^2 \mid f(x, y) = 0\}$  is a line.

2. Let  $a, b \in \mathbb{R}^* = \mathbb{R} \setminus \{0\}$  and

$$f(x, y) = \frac{x^2}{a^2} + \frac{y^2}{b^2} - 1.$$

The curve  $\{(x, y) \in \mathbb{R}^2 \mid f(x, y) = 0\}$  is an ellipse.

3. Let  $a, b \in \mathbb{R}^*$  and

$$g(x, y) = \frac{x^2}{a^2} - \frac{y^2}{b^2} - 1.$$

The curve  $\{(x, y) \in \mathbb{R}^2 \mid g(x, y) = 0\}$  is a hyperbola.

4. Spheres, quadrics such as ellipsoids, paraboloids, and hyperboloids in  $\mathbb{R}^3$  are all defined via a single polynomial equation of degree two. A line in  $\mathbb{R}^3$  can be defined by two equations in degree one.

The first property of algebraic curves is the following.

**Lemma 1.8.** The union of two affine plane curves is again an affine plane curve.

*Proof.* Let  $f_1, f_2 \in K[x, y]$  and let  $C_1 = \{f_1 = 0\}$  and  $C_2 = \{f_2 = 0\}$ . Then  $f_1 \cdot f_2 \in K[x, y]$  is a polynomial and

$$C_1 \cup C_2 = \{f_1 \cdot f_2 = 0\},$$

so that  $C_1 \cup C_2$  is an affine plane curve. □

**Exercise 3.** Write down an equation for the plane curve that is the union of the lines through any two vertices of the unit square.

Recall the following.

**Definition 1.9.** A polynomial  $P \in K[x_1, \dots, x_n]$  is **reducible** over  $K$  if there are non-constant polynomials  $Q, R \in K[x_1, \dots, x_n]$ , so  $\deg(Q), \deg(R) > 0$ , such that  $P = Q \cdot R$ . A polynomial  $P$  is **irreducible** if it is not reducible.

**Example 1.10.**  $x_1x_2$  is reducible,  $x_1 + x_2$  is irreducible.

**Fact 2.** Recall also that every polynomial  $P \in K[x_1, \dots, x_n]$  can be written as a product of irreducible factors  $P = f_1 \dots f_k$  in an essentially unique way up to multiplication by constants. We have

$$\{P = 0\} = \{f_1 = 0\} \cup \dots \cup \{f_k = 0\} \subseteq K^n,$$

so in particular every algebraic curve is a union of algebraic curves defined by irreducible polynomials.

In the course, we will consider questions such as the following.

1. When do polynomials  $f, g \in K[x, y]$  define the same affine plane curve?
2. What can be said about the intersection  $\{f = 0\} \cap \{g = 0\} \subset K^2$ ?

Very different questions can be approached through algebraic curves. For example, we can study integer solutions to some Diophantine equations.

**Example 1.11.** The unit circle is the curve

$$C = \{x^2 + y^2 = 1\} \subset \mathbb{R}^2.$$

Several parametrisations are known, such as

$$t \in [0, 2\pi) \mapsto (\cos t, \sin t) \in \mathbb{R}^2.$$

We can write down another parametrisation of  $C$  by considering lines through the point  $P = (-1, 0)$  using a stereographic projection. A line through  $P$  with slope  $t \in \mathbb{R}$  has equation

$$L_t = \{y = t(x + 1)\} \subset \mathbb{R}^2$$

and meets  $C$  in two points,  $P$  and  $P_t = (x(t), y(t))$ . We can determine the coordinate of  $P_t$  by solving the system

$$L_t \cap C = \begin{cases} y = t(x + 1) \\ x^2 + y^2 = 1 \end{cases}.$$

Replacing the value of  $y$  given by the first equation into the second yields two solutions for  $x(t)$ . The first one is  $x = -1$  and corresponds to the point  $P = (-1, 0)$ . The second is  $(x(t), y(t))$ , where

$$x(t) = \frac{1 - t^2}{1 + t^2}, \quad y(t) = \frac{2t}{1 + t^2}.$$

Note that when  $t \rightarrow \infty$ ,  $(x(t), y(t)) \rightarrow (-1, 0)$ , so that  $t \mapsto (x(t), y(t))$  is a parametrisation of  $C$  that identifies it with  $\mathbb{R} \cup \{\infty\}$ . The advantage of this parametrisation is that it is given by rational functions, that is  $x(t)$  and  $y(t)$  are of the form

$$t \mapsto \frac{p(t)}{q(t)},$$

where  $p, q$  are polynomials. One can use this parametrisation to get the general solution of the equation

$$x^2 + y^2 = z^2 \tag{1}$$

for  $x, y, z \in \mathbb{Z}$  coprime. If  $t = p/q \in \mathbb{Q}$ , where  $p, q \in \mathbb{Z}$  are coprime, then  $x(t), y(t) \in \mathbb{Q}$  becomes

$$x(t) = \frac{p^2 - q^2}{p^2 + q^2}, \quad y(t) = \frac{2pq}{p^2 + q^2}.$$

If  $x = p^2 - q^2$ ,  $y = 2pq$ , and  $z = p^2 + q^2$ ,  $x, y, z \in \mathbb{Z}$  satisfy (1). They are coprime precisely when  $p, q$  are coprime and not both odd. When  $p, q$  are coprime and both odd, then

$$x = \frac{p^2 - q^2}{2}, \quad y = pq, \quad \frac{p^2 + q^2}{2}$$

satisfy (1). Conversely, this is the general form of solutions in (1). Indeed, given  $x, y, z \in \mathbb{Z}$  coprime that satisfy (1),  $z \neq 0$  and

$$\frac{x^2}{z^2} + \frac{y^2}{z^2} = 1,$$

so that  $(x/z, y/z) \in \mathbb{C}$  and if  $(x, y, z) \neq (-1, 0, 1)$ , there is  $t \in \mathbb{R}$  such that  $(x/z, y/z) = (x(t), y(t))$ . But then since  $x/z, y/z \in \mathbb{Q}$ , we can take  $t \in \mathbb{Q}$  and  $x, y, z$  have the form above.

**Definition 1.12.** Let  $f \in \mathbb{R}[x, y]$  and let  $C = \{f = 0\}$ . A **rational point** of  $C$  is a point  $(x, y) \in C$ , that is  $f(x, y) = 0$ , such that  $x, y \in \mathbb{Q}$ .

**Example 1.13.** There are infinitely many rational points on the circle  $\{x^2 + y^2 = 1\} \subseteq \mathbb{R}^2$ , which can be described explicitly, and can be used to solve  $a^2 + b^2 = c^2$  for  $a, b, c \in \mathbb{Z}$ , a problem in number theory. Now take  $n \geq 3$  and consider

$$C = \{x^n + y^n - 1 = 0\}.$$

What are the rational points of  $C$ ? Write

$$x = \frac{a}{c}, \quad y = \frac{b}{c}, \quad a, b, c \in \mathbb{Z}, \quad c \neq 0.$$

Then

$$(x, y) \in C \iff a^n + b^n = c^n.$$

Fermat's Last Theorem by Wiles then states that there exists no solution with  $a, b \neq 0$ .

## 2 Complex plane curves

Let  $P \in \mathbb{R}[x, y]$  be a polynomial with coefficients in  $\mathbb{R}$ . A priori, it is natural to study the real plane curve  $C_{\mathbb{R}} = \{(x, y) \in \mathbb{R}^2 \mid P(x, y) = 0\}$ . However,  $P$  can also be seen as a polynomial with coefficients in  $\mathbb{C}$ , and it will often be simpler to study the complex plane curve  $C_{\mathbb{C}} = \{(x, y) \in \mathbb{C}^2 \mid P(x, y) = 0\}$ . We first explain some of the properties of algebraic curves that we would like to hold and explain why these properties do not necessarily hold for real plane curves and some unpleasant things happen.

**Fact 3.** Many real curves are so degenerate that they do not even have points, that is  $C_{\mathbb{R}} = \emptyset$ . If  $C_{\mathbb{R}} \neq \emptyset$ , the dimension of  $C_{\mathbb{R}}$  is difficult to determine.

**Example 2.1.** Let  $t \in \mathbb{R}$  and consider  $f_t(x, y) = x^2 + y^2 - t$  and the real plane curve  $C_t = \{f_t(x, y) = 0\} \subseteq \mathbb{R}^2$ . If  $t > 0$ ,  $C_t$  is a circle with radius  $\sqrt{t}$ , if  $t = 0$ , has  $C_0 = \{(0, 0)\}$ , and if  $t < 0$ ,  $C_t = \emptyset$ .

**Fact 4.** In general, it is not clear when two polynomials  $f, g \in \mathbb{R}[x, y]$  define the same real plane curve, that is when

$$\{(x, y) \in \mathbb{R}^2 \mid f(x, y) = 0\} = \{(x, y) \in \mathbb{R}^2 \mid g(x, y) = 0\}.$$

**Example 2.2.** Let  $f, g$  denote the polynomials

$$f(x, y) = x^2y + y^2 + x^3 + x, \quad g(x, y) = x^2 + 2xy + y^2.$$

Then, since  $f(x, y) = (x + 1) \cdot (x^2 + 1)$  and  $g(x, y) = (x + y)^2$ ,

$$\{(x, y) \in \mathbb{R}^2 \mid f(x, y) = 0\} = \{(x, y) \in \mathbb{R}^2 \mid g(x, y) = 0\}.$$

**Fact 5.** In general, it is hard to predict when a curve intersects a fixed line, or more generally when two real curves intersect.

**Example 2.3.** In the notation of Example 2.1, let  $C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 - 1\} \subset \mathbb{R}^2$  be the unit circle. Consider the line  $\{ax + by + c = 0\}$  for  $(a, b) \neq (0, 0)$ . Then, depending on  $(a, b, c) \in \mathbb{R}^3$ ,  $L \cap C$  consists of two points, one point, or is empty.

Lecture 2  
Thursday  
11/10/18

Most of these difficulties disappear when working with curves  $C_{\mathbb{C}} \subset \mathbb{C}^2$ , essentially because  $\mathbb{C}$  is algebraically closed, in other words the following theorem holds.

**Theorem 2.4** (Fundamental theorem of algebra). Let  $P \in \mathbb{C}[x]$  be a non-constant polynomial. Then  $P$  has at least one complex root, that is there exists  $\alpha \in \mathbb{C}$  such that  $P(\alpha) = 0$ .

A consequence of the fundamental theorem of algebra is that if  $P \in \mathbb{C}[x, y]$  is non-constant, then  $C = \{P = 0\}$  has infinitely many points. Assume without loss of generality that the polynomials in one variable  $P(\cdot, y)$  and  $P(x, \cdot)$  are not constant. This means that if

$$P(x, y) = \sum_{(r,s) \in \mathbb{N}^2} c_{r,s} x^r y^s,$$

there exist  $(r, s)$  and  $(r', s')$  in  $\mathbb{N}^2$ , which may be equal, such that  $r \neq 0$  and  $s' \neq 0$ , and  $c_{r,s} \neq 0$  and  $c_{r',s'} \neq 0$ . When  $y_0 \neq 0$  the polynomial  $P(x, y_0) \in \mathbb{C}[x]$  is non-constant and by Theorem 2.4, there exists  $x_0 \in \mathbb{C}$  such that  $P(x_0, y_0) = 0$ . In fact, for most choices of  $y_0 \neq y'_0$ , the polynomials  $P(\cdot, y_0)$  and  $P(\cdot, y'_0)$  are different, and have some distinct roots. It follows that  $\{P = 0\}$  contains infinitely many points.

**Example 2.5.** Let  $a, b, c \in \mathbb{C}$  with  $(a, b) \neq (0, 0)$ , and let  $f(x, y) = ax + by + c$ . If  $a \neq 0$ , for each  $y \in \mathbb{C}$ , there is precisely one solution of  $f(x, y) = 0$ , namely

$$x = -\frac{b}{a}y - \frac{c}{a}.$$

Thus  $\mathbb{C}^2 \supset \{f = 0\} = C \rightarrow \mathbb{C} \cong \mathbb{R}^2$  is an isomorphism. We will call  $C$  a **complex line**.

**Remark 2.6.** It is difficult to draw complex curves. Our intuition is for real vector spaces, and this makes complex curves hard to visualise. They are objects of real dimension two in  $\mathbb{C}^2 \cong \mathbb{R}^4$ , a four-dimensional real vector space.

**Example 2.7.** Let  $f(x, y) = x^2 + y^2$ . Then  $f(x, y) = (x + iy) \cdot (x - iy)$  and, as in Lemma 1.8,  $C = \{f = 0\} \subset \mathbb{C}^2$  is the union of the two complex lines  $\{x + iy = 0\}$  and  $\{x - iy = 0\}$ . When seen as  $\mathbb{R}$ -vector spaces, these two planes meet at exactly one point corresponding to  $(0, 0) \in \mathbb{R}^2 \subseteq \mathbb{C}^2$ , the only real point of  $C$ . It is difficult to imagine two planes meeting in one point, because our intuition relies on three-dimensional space  $\mathbb{R}^3$ , while  $\mathbb{C}^2 = \mathbb{R}^4$ .

Describing intersections is also easier.

**Example 2.8.** Consider  $C = \{x^2 + y^2 - 1 = 0\} \subseteq \mathbb{C}^2$  and  $L = \{ax + by + c = 0\} \subseteq \mathbb{C}^2$ . If  $b \neq 0$ , we determine the intersection  $C \cap L$  by solving the equation

$$x^2 + y^2 = 1, \quad y = -\frac{a}{b}x - \frac{c}{b}.$$

If  $a^2 = -b^2$  or  $c = 0$ , there are one or two solutions. Again, it is hard to imagine a two-dimensional real surface which meets a real plane in two points.

We now turn to the question of recognising when two polynomials define the same plane curve. Here again, working in  $\mathbb{C}$  is a simplification.

**Theorem 2.9** (Consequence of Hilbert's Nullstellensatz). Let  $f, g \in \mathbb{C}[x, y]$  be two polynomials. Then

$$\{f = 0\} = \{g = 0\}$$

if and only if there exist  $P_1, \dots, P_k \in \mathbb{C}[x, y]$ ,  $a_1, \dots, a_k, b_1, \dots, b_k \in \mathbb{Z}_{>0}$  and  $\lambda_1, \lambda_2 \in \mathbb{C}^*$  such that

$$f(x, y) = \lambda_1 P_1^{a_1} \dots P_k^{a_k}, \quad g(x, y) = \lambda_2 P_1^{b_1} \dots P_k^{b_k}. \quad (2)$$

*Proof.* Assume that (2) holds. Then by the proof of Lemma 1.8,

$$\{f = 0\} = \{P_1^{a_1} = 0\} \cup \dots \cup \{P_k^{a_k} = 0\} = \{P_1 = 0\} \cup \dots \cup \{P_k = 0\},$$

because if  $\alpha \in \mathbb{C}$  is such that  $\alpha^n = 0$ , then  $\alpha = 0$ . The same holds for  $\{g = 0\}$ . Therefore  $\{f = 0\} = \{g = 0\}$ . The second half of the proof needs tools of commutative algebra and is omitted.  $\square$

**Remark 2.10.** The theorem fails over  $\mathbb{R}$ . Let  $f(x, y) = x^2 + 1$  and  $g(x, y) = 1$ . Then  $\{f = 0\} = \{g = 0\} = \emptyset$  but the conclusion in (2) is not true.

Thus, the relation between the geometric shape  $C = \{f = 0\}$  in  $\mathbb{C}^2$  and the polynomial  $f \in \mathbb{C}[x, y]$  is more transparent than in  $\mathbb{R}$ . We will always work in  $\mathbb{C}$ . Let us introduce some important notions for the study of polynomials.

**Definition 2.11.** A polynomial  $f \in K[x, y]$  has **no repeated factors** over  $K$  if it cannot be written as a product of the form

$$f(x, y) = g(x, y)^2 \cdot h(x, y),$$

where  $g, h \in K[x, y]$  and  $g$  is non-constant.

**Exercise 4.** Show that this is equivalent to

$$f = P_1 \cdots P_k,$$

where  $P_1, \dots, P_k$  are distinct irreducible polynomials.

**Corollary 2.12.** Let  $f, g \in \mathbb{C}[x, y]$  be polynomials with no repeated factors. Then  $f, g$  define the same complex plane curve

$$\{f = 0\} = \{g = 0\}$$

if and only if there is a non-zero constant  $\lambda \in \mathbb{C}^*$  such that  $f = \lambda g$ .

*Proof.* Follows immediately from Theorem 2.9.  $\square$

**Remark 2.13.** If  $f = P_1^{a_1} \cdots P_k^{a_k}$  with  $P_i$  irreducible for all  $i$  and  $a_i \in \mathbb{N}$ , then  $\{f = 0\} = \{g = 0\}$  where  $g = P_1 \cdots P_k$ . We do not lose anything by only looking at  $f$  with no repeated factors.

Let  $C \subseteq \mathbb{C}^2$  be a complex plane curve. We have proved that, up to multiplication by  $\lambda \in \mathbb{C}^*$ , there is a unique non-constant polynomial  $f \in \mathbb{C}[x, y]$  with no repeated factors such that

$$C = \{f = 0\}.$$

It makes sense to define the following.

**Definition 2.14.** The **degree** of an affine curve  $C \subseteq \mathbb{C}^2$  is the degree of any polynomial with no repeated factors  $f$  such that  $C = \{f = 0\}$ , that is

$$\deg(C) = \deg(f).$$

**Example 2.15.** Lines have degree one, since they are defined by a linear polynomial. Conics have degree two.  $\{x^2y + y^2 + x + 1 = 0\}$  has degree 3, assuming it has no repeated factors.

Unless mentioned otherwise, in the first few weeks, we will assume that polynomials have no repeated factors.

**Definition 2.16.** Let  $f_1, f_2 \in \mathbb{C}[x, y]$  be polynomials with no repeated factors and let  $C_1 = \{f_1 = 0\}$  and  $C_2 = \{f_2 = 0\}$  be the associated complex curves. The curves  $C_1$  and  $C_2$  have **no common component** if there is no non-constant polynomial  $P$  that divides both  $f_1$  and  $f_2$ .

This is equivalent to saying that if  $f = P_1^{a_1} \cdots P_k^{a_k}$  and  $g = Q_1^{b_1} \cdots Q_l^{b_l}$  with  $P_i, Q_i$  irreducibles,  $P_i$  distinct, and  $Q_i$  distinct, then  $\lambda P_i \neq Q_j$  for all  $i, j, \lambda \in \mathbb{C}^*$ .

**Exercise 5.** Show that if  $C_1$  and  $C_2$  have no common component, then  $\deg(C_1 \cup C_2) = \deg(C_1) + \deg(C_2)$ .

**Exercise 6.** Let  $L, L'$  be the lines

$$L = \{ax + by + c = 0\} \subset \mathbb{C}^2, \quad L' = \{a'x + b'y + c' = 0\} \subset \mathbb{C}^2.$$

1. Show that  $L$  and  $L'$  meet at exactly one point if and only if  $ab' - a'b \neq 0$ .
2. Show that  $L = L'$  if and only if there exists  $\lambda \in \mathbb{C}$  such that  $\lambda \neq 0$  and

$$a' = \lambda a, \quad b' = \lambda b, \quad c' = \lambda c.$$

**Remark 2.17** (First aid topology).

1. A **topological space** is a set  $X$  with a collection of open subsets  $\{U_i \subset X\}$  such that
  - (a)  $\emptyset$  and  $X$  are open,
  - (b) any union  $\cup_{i \in I} U_i$  of open sets  $U_i$  is open, and
  - (c) any finite intersection  $\cap_{i=1}^k U_i$  of open sets  $U_i$  is open.
2. A **metric space**  $X$ , such as  $(\mathbb{C}^n, \|\cdot\|)$ , is a topological space. The open sets are given by arbitrary unions and finite intersections of the familiar open balls  $B(x, \epsilon) = \{z \in X \mid \|z - x\| < \epsilon\}$ .
3. A subset  $X \subset Y$  of a topological space  $Y$  inherits a topology from  $Y$ . The open sets of  $X$  are the sets  $X \cap U$ , where  $U \subset Y$  is an open set of  $Y$ .
4.  $X$  is **compact** if for all open covering  $X = \sum_{i \in I} U_i$  where  $U_i$  are open, there exists a finite subcovering  $\cup_{i_1, \dots, i_k} U_{i_j}$  for  $\{i_1, \dots, i_k\} \subseteq I$ .
5. The **Heine-Borel theorem** states that a subset  $X$  of  $\mathbb{R}^n$  or of  $\mathbb{C}^m$  is compact if and only if  $X$  is closed, that is its complement is open, and bounded for the usual norm.
6. A closed subset of a compact space is compact.
7. A map  $f : X \rightarrow Y$  between topological spaces is **continuous** if and only if  $f^{-1}(U)$  is open in  $X$  whenever  $U \subset Y$  is open. It follows that  $f^{-1}(F)$  is closed whenever  $F \subset Y$  is closed. In particular, if  $f \in \mathbb{C}[x_1, \dots, x_n]$  is a polynomial,  $f$  defines a map  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  that is continuous, and

$$f^{-1}(\{0\}) = \{f = 0\} \subset \mathbb{C}^n$$

is closed because  $\{0\}$  is a closed subset of  $\mathbb{C}$ .

In particular,  $\mathbb{C}^2$  is a topological space with the Euclidean distance in  $\mathbb{R}^4$  and the affine plane curve  $C = \{f = 0\} \subseteq \mathbb{C}^2 \cong \mathbb{R}^4$  inherits a topology. Open sets of  $C$  are  $U \cup C$  where  $U \subseteq \mathbb{C}^2$  is open. So algebraic curves have a natural topology.

**Lemma 2.18.** Let  $C \subset \mathbb{C}^2$  be an affine plane curve, then  $C$  is not compact.

*Proof.* Since  $f$  is a continuous function  $\mathbb{C}^2 \rightarrow \mathbb{C}$ ,  $C = \{f = 0\} = f^{-1}(\{0\})$ , and  $\{0\}$  is closed in  $\mathbb{C}$ ,  $C$  is closed in  $\mathbb{C}^2$ . We check that  $C \subset \mathbb{C}^2$  is not bounded. Assume that it is, then there is a constant  $M > 0$  such that  $C \subset B(0, M)$ , where the open ball is

$$B(0, M) = \left\{ |x|^2 + |y|^2 < M \right\}.$$

Want to show that some points in  $\mathbb{C}$  are outside this open ball. Let  $y_0 \in \mathbb{C}$  be such that  $|y_0| > M$  and assume we can arrange for  $g = f(\cdot, y_0)$  to be a non-constant polynomial of  $x$ . By the fundamental theorem of algebra,  $g$  has a root  $x_0 \in \mathbb{C}$  and the point  $(x_0, y_0) \in C$ . This is a contradiction, as  $(x_0, y_0) \notin B(0, M)$ .  $\square$

(TODO Exercise: what if  $f(x, y)$  happens to be a polynomial of  $y$  alone, so that this cannot be arranged?)



### 3 Projective space

Recall that it is difficult to determine when two affine plane curves  $C, C' \in \mathbb{C}^2$  intersect, and some curves do not in fact intersect even over  $\mathbb{C}$ . We want to fix that, and the key is adding points at infinity.

**Example 3.1.**

1. Consider two distinct lines

$$L_1 = \{ax + by + c = 0\}, \quad L_2 = \{a'x + b'y + c' = 0\}.$$

Then  $L_1$  and  $L_2$  meet at exactly one point if and only if

$$\det \begin{pmatrix} a & b \\ a' & b' \end{pmatrix} \neq 0.$$

But we can pretend that parallel lines meet at a point at infinity corresponding to the direction vector.

2. Consider the asymptotic curve and line

$$C = \{xy - 1 = 0\}, \quad L = \{x = 0\}.$$

Then  $C$  and  $L$  do not meet, but again we can pretend that they meet at a point at infinity.

A heuristic trick is to introduce a variable  $z$ .

1. Replace  $x$  by  $x/z$  and  $y$  by  $y/z$ .
2. Solve for  $z = 0$ .

**Example 3.2.**

1. Consider the lines

$$L_1 = \{x + y + 1 = 0\}, \quad L_2 = \{x + y - 1 = 0\}.$$

Clearly  $L_1$  and  $L_2$  do not meet. Let us apply the trick. By 1 and 2 we get

$$\begin{cases} x/z + y/z + 1 = 0 \\ x/z + y/z - 1 = 0 \end{cases} \implies \begin{cases} x + y + z = 0 \\ x + y - z = 0 \end{cases} \implies \begin{cases} x + y = 0 \\ x + y = 0 \end{cases}.$$

We get that the point  $(1, -1, 0)$  is a common solution. This will be called the point at infinity.

2. Consider the asymptotic curve and line

$$C = \{xy - 1 = 0\}, \quad L = \{x = 0\}.$$

Apply 1 and 2 to get

$$\begin{cases} xy - z^2 = 0 \\ x/z = 0 \end{cases} \implies \begin{cases} xy = 0 \\ x = 0 \end{cases}.$$

We get that  $(0, 1, 0)$  is a common solution. Again, this will be called the point at infinity.

To make this formal, we introduce the projective plane  $\mathbb{P}^2$ . We will add points at infinity to  $\mathbb{C}^2$ , in such a way that asymptotic curves meet at infinity. We will then compactify an affine plane curve  $C$  so that the two compactifications are compatible, that is

$$(C \subseteq \mathbb{C}^2) \hookrightarrow (\overline{C} \subseteq \mathbb{P}^2).$$

**Notation 3.3.** Fix  $n \geq 0$  and  $\mathbb{C}^{n+1}$ . Let  $\underline{0} = (0, \dots, 0) \in \mathbb{C}^{n+1}$  be the origin of the  $(n+1)$ -dimensional complex Euclidean space. We will denote

$$W = \mathbb{C}^{n+1} \setminus \{0\},$$

that a point  $x \in W$  is given by  $x = (x_0, \dots, x_n)$  where  $x_0, \dots, x_n \in \mathbb{C}$  are not all zero. We define the equivalence relation on  $W$ , for any  $x, y \in W$  by

$$x \sim y \iff \exists \lambda \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}, x = \lambda y.$$

**Exercise 7.** Show that  $\sim$  is an equivalence relation on  $W$ .

**Notation 3.4.** Given  $x \in W$ , we denote

$$[x] = \{y \in W \mid x \sim y\}.$$

For simplicity, if  $x = (x_0, \dots, x_n)$  we will denote  $[x] = [x_0, \dots, x_n]$  instead of  $x = [(x_0, \dots, x_n)]$ .

**Definition 3.5.** The  $n$ -dimensional projective space  $\mathbb{P}_{\mathbb{C}}^n$  or  $\mathbb{P}^n(\mathbb{C})$  or simply  $\mathbb{P}^n$  is defined as the quotient of  $W$  by  $\sim$ , that is

$$\mathbb{P}^n = \frac{W}{\sim} = \{[x] \mid x \in W = \mathbb{C}^{n+1} \setminus \{0\}\}.$$

The coordinates of  $\mathbb{P}^n$  are  $[x] \in \mathbb{P}^n$  except  $[0, \dots, 0]$  and  $[\lambda x_0, \dots, \lambda x_n] = [x_0, \dots, x_n]$  for  $\lambda \in \mathbb{C}^*$ . In other words, in  $\mathbb{P}^n$ , two points  $[x_0, \dots, x_n]$  and  $[y_0, \dots, y_n]$  are the same point if and only if there exists a non-zero constant  $\lambda$  such that

$$x_0 = \lambda y_0, \quad \dots, \quad x_n = \lambda y_n.$$

**Exercise 8.** Show that  $[x] = [y]$  if and only if  $x \sim y$ . Show that if  $y \notin [x]$  then  $[x] \cap [y] = \emptyset$ .

**Example 3.6.** The point  $[1, 2, i]$  is the same as the point  $[i, 2i, -1]$ .

**Exercise 9.** Show that there exists a bijection between  $\mathbb{P}^n$  and the set of all the one-dimensional subspaces of  $\mathbb{C}^{n+1}$ . In fact, if  $V$  is a finite-dimensional vector space over  $\mathbb{C}$  without the choice of a basis, we can define the associated projective space  $\mathbb{P}(V)$  as the set of one-dimensional linear subspaces of  $V$ .

**Example 3.7.** For any non-zero  $x \in \mathbb{C}$  we have  $[x] = [1]$ . So  $\mathbb{P}^0 = \mathbb{C}^1 \setminus \{0\} / \sim$  is a point  $\mathbb{C}^0 = \{[1]\}$ .

**Notation 3.8.** For any  $i = 0, \dots, n$ , denote the **affine chart**

$$U_i = \{[x] = [x_1, \dots, x_n] \in \mathbb{P}^n \mid x_i \neq 0\} \subseteq \mathbb{P}^n.$$

**Lemma 3.9.**  $\mathbb{P}^n = U_0 \cup \dots \cup U_n$ .

*Proof.* Take  $[x] = [x_0, \dots, x_n] \in \mathbb{P}^n$  in  $\mathbb{P}^n$  then  $x \in W$  and in particular  $x = (x_0, \dots, x_n)$  where at least one of the coefficients is non-zero, say  $x_i \neq 0$ . Then  $[x] \in U_i$ . Thus any  $[x] \in \mathbb{P}^n$  is contained in the union of  $U_0, \dots, U_n$ .  $\square$

**Lemma 3.10.** Pick  $i = 0, \dots, n$ . Define  $\phi_i : \mathbb{C}^n \rightarrow U_i$  by

$$\phi_i(y_1, \dots, y_n) = [y_1, \dots, y_i, 1, y_{i+1}, \dots, y_n].$$

Then  $\phi_i$  is a bijection and its inverse  $\rho_i : U_i \rightarrow \mathbb{C}^n$  is given by

$$\rho_i[x_0, \dots, x_n] = \left( \frac{x_0}{x_i}, \dots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \dots, \frac{x_n}{x_i} \right).$$

*Proof.* First note that both  $\phi_i$  and  $\rho_i$  is well-defined, indeed, if

$$(y_1, \dots, y_n) \in \mathbb{C}^n$$

then

$$(y_1, \dots, y_i, 1, y_{i+1}, \dots, y_n) \in W$$

and therefore  $[y_1, \dots, y_i, 1, y_{i+1}, \dots, y_n] \in \mathbb{P}^n$ . Similarly, if  $[x_0, \dots, x_n] = [x'_0, \dots, x'_n]$  then it follows that  $\rho_i[x_0, \dots, x_n] = \rho_i[x'_0, \dots, x'_n]$ . Thus, it is enough to show that both  $\phi_i \circ \rho_i$  and  $\rho_i \circ \phi_i$  coincide with the identity. We have

$$\rho_i(\phi_i(y_1, \dots, y_n)) = \rho_i(y_1, \dots, y_i, 1, y_{i+1}, \dots, y_n) = \left(\frac{y_1}{1}, \dots, \frac{y_{i-1}}{1}, \frac{y_i}{1}, \dots, \frac{y_n}{1}\right) = (y_1, \dots, y_n).$$

Similarly,

$$\phi_i(\rho_i[x_0, \dots, x_n]) = \phi_i\left(\frac{x_0}{x_i}, \dots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \dots, \frac{x_n}{x_i}\right) = \left[\frac{x_0}{x_i}, \dots, \frac{x_{i-1}}{x_i}, 1, \frac{x_{i+1}}{x_i}, \dots, \frac{x_n}{x_i}\right] = [x_0, \dots, x_n].$$

Thus they are inverses.  $\square$

**Example 3.11.** Let  $n = 2$  and  $[x_0, x_1, x_2] \in \mathbb{P}^2$ . Let  $\phi_1 : \mathbb{C}^2 \rightarrow U_1$  be defined by  $(0, 0) \mapsto [0, 1, 0]$  and  $\phi_2 : \mathbb{C}^2 \rightarrow U_2$  be defined by  $(0, 0) \mapsto [0, 0, 1]$ . Check that

$$\rho_i(\phi_i(y_1, \dots, y_n)) = \rho_i([y_1, \dots, 1, \dots, y_n]) = (y_1, \dots, y_n).$$

The previous two lemmas can be used to define a topology on  $\mathbb{P}^n$ . Let  $U \subseteq \mathbb{P}^n$ , then  $U$  is open if and only if  $\phi_i^{-1}(U \cap U_i) \subseteq \mathbb{C}^n$  is open in  $U_i \cong \mathbb{C}^n$  for any  $i = 0, \dots, n$ .

**Exercise 10.** Show that  $U_i \subseteq \mathbb{P}^n$  is open in  $\mathbb{P}^n$  for all  $i = 0, \dots, n$ .

**Exercise 11.** We can define another topology on  $\mathbb{P}^n$  as the quotient topology induced by the map  $\pi : W \rightarrow \mathbb{P}^n$  defined by  $(x_0, \dots, x_n) \mapsto [x_0, \dots, x_n]$ . A subset  $U \subseteq \mathbb{P}^n$  is open if and only if its preimage  $\pi^{-1}(U) \subseteq W$  is open in  $W$ . Show that this indeed defines a topology, and that this topology coincides with the one defined above using the maps  $\phi_i$ . Check that with this topology on  $\mathbb{P}^n$ ,  $\pi$  is continuous.

**Exercise 12.** Prove that  $\mathbb{P}^n$  is compact. Restrict the projection  $\pi$  of the previous exercise to the  $(n+1)$ -dimensional sphere  $S^{n+1} \subseteq W$ , and check that this restriction is surjective and continuous. Since  $S^{n+1}$  is compact, it follows that  $\mathbb{P}^n$  is compact as well.

**Example 3.12.**

1.  $\mathbb{P}^1 = \{[x_0, x_1] \mid (x_0, x_1) \in \mathbb{C}^2 \setminus \{0\}\}$  is the union of two copies  $U_0$  and  $U_1$  of  $\mathbb{C}^1$ . The intersection of  $U_0$  and  $U_1$  is  $U_0 \cap U_1 = \{[x_0, x_1] \mid x_0 \neq 0, x_1 \neq 0\}$  can be identified with  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$  via the map  $[x_0, x_1] \mapsto x_1/x_0$ . Using this identification and the maps  $\rho_i$ , the inclusion  $U_0 \cap U_1 \subseteq U_0$  is the map  $\mathbb{C}^* \rightarrow \mathbb{C}^1$  sending  $z$  to itself, and the inclusion  $U_0 \cap U_1 \subseteq U_1$  is the map  $\mathbb{C}^* \rightarrow \mathbb{C}^1$  given by  $z \mapsto 1/z$ .  $\mathbb{P}^1$  is glued together from two copies of  $\mathbb{C}$  along  $U_0 \cap U_1$  by these inclusions, so  $\mathbb{P}^1 = \mathbb{C}^1 \cup \{\infty\}$ . Over the real numbers,  $\mathbb{P}^1(\mathbb{R})$  is built up in the same way from two copies of  $\mathbb{R}^1$ , and can be identified with the circle  $S^1$ .
2.  $\mathbb{P}^2$  is the union of three copies of  $U_0 \cong U_1 \cong U_2 \cong \mathbb{C}^2$ , and the intersection can be described similarly. At infinity we will have a line. More generally,  $\mathbb{P}^n$  can be described similarly.

In practice, we will view the affine complex plane  $\mathbb{C}^2$  as being embedded in  $\mathbb{P}^2$  as one of the open sets  $U_i$ . For example, we identify

$$(x, y) \in \mathbb{C}^2 \iff [x, y, 1] \in \mathbb{P}^2.$$

What is the complement of this embedding, that is the points at infinity?

**Notation 3.13.** For any  $i = 0, \dots, n$ , denote

$$\mathcal{P}_i = \{[x_0, \dots, x_n] \in \mathbb{P}^n \mid x_i = 0\} \subseteq \mathbb{P}^n.$$

**Lemma 3.14.** For any  $i = 0, \dots, n$ , we have

$$\mathbb{P}^n = U_i \sqcup \mathcal{P}_i.$$

Moreover if we define  $f_i : \mathbb{P}^{n-1} \rightarrow \mathcal{P}_i$  by

$$f_i [z_0, \dots, z_{n-1}] \mapsto [z_0, \dots, z_{i-1}, 0, z_i, \dots, z_{n-1}]$$

then  $f_i$  is a bijection.

*Proof.* TODO Exercise: both statements are easy to check. □

In conclusion, we have that

$$\mathbb{P}^n = U_i \sqcup \mathcal{P}_i \cong \mathbb{C}^n \sqcup \mathbb{P}^{n-1} = \dots = \mathbb{C}^n \cup \dots \cup \mathbb{C}^0.$$

$\mathcal{P}_i$  is called the **hyperplane at infinity**.

**Example 3.15.**

1. We have already seen that  $\mathbb{P}^0$  is a point.
2.  $\mathbb{P}^1 \cong \mathbb{C}^1 \cup \mathbb{P}^0$ . In other words,  $\mathbb{P}^1$  is obtained by adding a point at infinity to the complex line  $\mathbb{C}$ . It is a way to compactify the real plane. If  $\mathbb{C}^1$  above is  $U_0$ , this point at infinity is the origin of the other open subset  $U_1 \cong \mathbb{C}^1$ . One can show that there is an bijection which is also a homeomorphism between  $\mathbb{P}^1$  and the sphere  $S^1$  in  $\mathbb{R}^3$  by the identification  $r \mapsto 1/r$ .  $\mathbb{P}^1$  is called the projective line.
3.  $\mathbb{P}^2 \cong \mathbb{C}^2 \cup \mathbb{P}^1$ . Thus  $\mathbb{P}^2$  is obtained by adding a projective line at infinity to the complex plane  $\mathbb{C}^2$ .  $\mathbb{P}^2$  is called the projective plane.

Note that the point at infinity or the line at infinity is not unique, it depends on the choice of the coordinate and in our settings, it depends on  $i$ . In the future, we will often fix  $i$ , and this will give us a unique choice of the point or line at infinity.

Lecture 5  
Thursday  
18/10/18

## 4 Projective curves

Recall that an algebraic curve  $C \subseteq \mathbb{C}^2$  is given by  $C = \{f(x, y) = 0\}$  for  $f \in \mathbb{C}[x, y]$ . In this section, we want to define projective curves similarly in  $\mathbb{P}^2$  and then compactify these affine curves. If we try to define a plane projective curve in the same way as an affine curve, that is as  $\{f = 0\} \subseteq \mathbb{P}^2$ , where  $f$  is a polynomial in  $\mathbb{C}[x_0, x_1, x_2]$ , the first hurdle we encounter is that  $f$  does not define a function on  $\mathbb{P}^2$ .

**Example 4.1.**

1. Let  $f(x, y, z) = x^2 + y^2 - z^2$ . Then  $f(1, 1, 1) = 1 + 1 - 1 = 1 \neq 4 = 4 + 4 - 4 = f(2, 2, 2)$ , so that  $f$  does not define a function on  $\mathbb{P}^2$ . It makes no sense to talk about  $f(1, 1, 1)$ , because in  $\mathbb{P}^2$ ,  $[1, 1, 1] = [2, 2, 2]$ , so the value would not be well-defined. However, the locus where  $f$  vanishes in this case is well-defined. If  $f(x, y, z) = 0$ , then  $f(\lambda x, \lambda y, \lambda z) = 0$  for all  $\lambda \in \mathbb{C}^*$ , so that the subset  $\{[x, y, z] \in \mathbb{P}^2 \mid f(x, y, z) = 0\}$  is well-defined.
2. Let  $g(x_0, x_1, x_2) = x_0^2 + x_1$ , then  $g(i, 1, 0) = 0$ , but  $g(2i, 2, 0) \neq 0$ , so that in this case, the vanishing locus  $\{[x_0, x_1, x_2] \in \mathbb{P}^2 \mid g(x_0, x_1, x_2) = 0\}$  is not even well-defined.

The example shows that if we want to define projective curves as

$$\{[x_0, x_1, x_2] \in \mathbb{P}^2 \mid f(x_0, x_1, x_2) = 0\},$$

then  $f \in \mathbb{C}[x_0, x_1, x_2]$  has to satisfy some additional properties.

**Definition 4.2.** A polynomial  $f \in \mathbb{C}[x_0, \dots, x_n]$  is **homogeneous** if all its monomials have the same degree  $d \in \mathbb{N}$ , that is

$$f(x_0, \dots, x_n) = \sum_{|\alpha| = \sum_{i=0}^n \alpha_i = d} a_\alpha x^\alpha, \quad \alpha \in \mathbb{N}^{n+1}.$$

**Example 4.3.**

1. The polynomial  $f(x, y, z) = x^2 + y^2 - z^2$  is homogeneous of degree two, and the polynomial  $g(x_0, x_1, x_2) = x_0^2 + x_1$  is not homogeneous.
2.  $f(x, y) = Ax^3 + Bx^2y + Cxy^2 + Dy^3$  is homogeneous of degree three, and all the homogeneous polynomials of degree three in  $x, y$  can be written in this form.
3.  $g(x, y, z) = x^4 - 2x^2yz + yz^3$  is homogeneous of degree four.

**Lemma 4.4.** Let  $P \in \mathbb{C}[x_0, \dots, x_n]$  be a polynomial. If  $P$  is homogeneous of degree  $d$ , then

$$P(\lambda x_0, \dots, \lambda x_n) = \lambda^d P(x_0, \dots, x_n),$$

for all  $\lambda \in \mathbb{C}$ , and all  $(x_0, \dots, x_n) \in \mathbb{C}^{n+1}$ .

*Proof.* Let  $P$  be a homogeneous polynomial of degree  $d$ . Then  $P = M_1 + \dots + M_k$ , where each  $M_i$  is a monomial of degree  $d$ , that is a product  $a_i \cdot x^{\alpha_i}$  where  $\alpha_i \in \mathbb{N}^{n+1}$  and  $a_i \in \mathbb{C}$ . For each monomial  $M = a_\alpha x^\alpha = a_\alpha x_0^{\alpha_0} \dots x_n^{\alpha_n}$  of degree  $d$ , for  $a_\alpha \in \mathbb{C}$  and  $\alpha \in \mathbb{N}^{n+1}$ , we have

$$M(\lambda x_0, \dots, \lambda x_n) = a_\alpha (\lambda^{\alpha_0} x_0^{\alpha_0}) \dots (\lambda^{\alpha_n} x_n^{\alpha_n}) = \lambda^{\sum_{i=0}^n \alpha_i} a_\alpha x^\alpha = \lambda^d a_\alpha x^\alpha = \lambda^d M(x_0, \dots, x_n),$$

because  $\sum_{i=0}^n \alpha_i = |\alpha| = d$ . For the arbitrary homogeneous polynomial  $P$ , write  $P = \sum_{i=1}^k M_i$ . Thus

$$P(\lambda x_0, \dots, \lambda x_n) = \sum_{i=1}^k M_i(\lambda x_0, \dots, \lambda x_n) = \lambda^d \sum_{i=1}^k M_i(x_0, \dots, x_n) = \lambda^d P(x_0, \dots, x_n).$$

□

**Exercise 13.** Prove the converse implication for the previous lemma, that is if  $P \in \mathbb{C}[x_0, \dots, x_n]$  and

$$P(\lambda x_0, \dots, \lambda x_n) = \lambda^d P(x_0, \dots, x_n),$$

for all  $\lambda \in \mathbb{C}$ , and all  $(x_0, \dots, x_n) \in \mathbb{C}^{n+1}$ , then  $P$  is homogeneous of degree  $d$ .

**Proposition 4.5.** Let  $P$  be a homogeneous polynomial. Then

$$\{[x_0, \dots, x_n] \in \mathbb{P}^n \mid P(x_0, \dots, x_n) = 0\}$$

is well-defined.

*Proof.* We have to check that if  $(x_0, \dots, x_n) \sim (y_0, \dots, y_n)$ ,  $P(x_0, \dots, x_n) = 0$  if and only if  $P(y_0, \dots, y_n) = 0$ . This follows immediately from Lemma 4.4, because by definition of  $\sim$ ,  $(y_0, \dots, y_n) = (\lambda x_0, \dots, \lambda x_n)$  for some  $\lambda \in \mathbb{C}^*$ , so  $P(x_0, \dots, x_n) = 0$  if and only if  $P(\lambda x_0, \dots, \lambda x_n) = \lambda^d P(x_0, \dots, x_n) = 0$ . □

**Notation 4.6.** Unless mentioned otherwise, arbitrary polynomials will be denoted  $f, g, h, \dots$  while homogeneous polynomials will be denoted  $P, Q, R, \dots$

Let  $P_1, \dots, P_k \in \mathbb{C}[x_0, \dots, x_n]$  be homogeneous polynomials. Then the vanishing locus

$$\{P_1 = \dots = P_k = 0\} \subseteq \mathbb{P}^n$$

is a **projective variety**. These are the main object of study of algebraic geometry. In this class, we will focus on plane projective curves.

**Definition 4.7.** A complex projective plane algebraic curve is

$$C = \{[x_0, x_1, x_2] \in \mathbb{P}^2 \mid P(x_0, x_1, x_2) = 0\} \subseteq \mathbb{P}^2,$$

where  $P$  is a non-constant homogeneous polynomial  $P \in \mathbb{C}[x_0, x_1, x_2]$ .

From now on  $n = 2$ , so consider  $\mathbb{P}^2$ .

**Example 4.8.** The hyperplanes at infinity  $\mathcal{P}_i = \{x_i = 0\} \subseteq \mathbb{P}^2$  for  $i = 0, 1, 2$  are projective curves of degree one, where  $x_0$  is a homogeneous polynomial of degree one.

**Definition 4.9.** A **projective line** is a projective curve defined by a homogeneous polynomial  $ax_0 + bx_1 + cx_2 = 0$  of degree one.

We have seen that in the case of affine plane curves  $\{f = 0\} \subseteq \mathbb{C}^2$ , the irreducible factors of the polynomial  $f$  were important when studying curves. The next lemma ensures that the same type of results hold for projective curves.

**Lemma 4.10.** Let  $P \in K[x_0, \dots, x_n]$  be a non-zero homogeneous polynomial. Assume that  $P = Q \cdot R$ , where  $Q, R \in K[x_0, \dots, x_n]$ . Then the polynomials  $Q$  and  $R$  are homogeneous.

*Proof.* TODO Exercise. □

**Remark 4.11.** As in the case of affine plane curves, we can therefore make sense of irreducible projective plane curves  $C = \{P = 0\} \subset \mathbb{P}^2$  with  $P$  irreducible. When  $P$  is reducible  $P = P_1^{a_1} \dots P_k^{a_k}$  with  $P_i$  distinct irreducible polynomials and  $a_i \in \mathbb{N}_{>0}$ , the projective curves  $C_i = \{P_i = 0\}$  in  $\{P = 0\} = \{P_1 = 0\} \cup \dots \cup \{P_k = 0\}$  are called the **irreducible components** of  $C = \{P = 0\}$ .

**Remark 4.12.** Hilbert's Nullstellensatz, and Theorem 2.9, still holds. In particular, if  $P, Q \in \mathbb{C}[x, y, z]$  are homogeneous polynomials with no repeated factors,  $\{P = 0\} = \{Q = 0\} \subseteq \mathbb{P}^2$  if and only if  $P = \lambda Q$  for  $\lambda \in \mathbb{C}^*$ .

**Definition 4.13.** Let  $C \subseteq \mathbb{P}^2$  be a projective plane curve and  $P$  any homogeneous polynomial with no repeated factor such that  $C = \{P = 0\} \subseteq \mathbb{P}^2$ . The **degree** of  $C$  is the degree of  $P$ .

**Exercise 14.** Show that the union of two projective curves  $C_1, C_2 \subseteq \mathbb{P}^2$  is a projective curve. More generally, show that if  $X_1, X_2 \subseteq \mathbb{P}^2$  are projective varieties, then  $X_1 \cup X_2 \subseteq \mathbb{P}^n$  is again a projective variety.

**Exercise 15.** Show that if  $X = \{P = 0\} \subseteq \mathbb{P}^n$  for a non-zero homogeneous polynomial  $P \in \mathbb{C}[x_0, \dots, x_n]$ , then

$$X = \{x_0 P = \dots = x_n P = 0\}.$$

**Lemma 4.14.** Let  $C \subseteq \mathbb{P}^2$  be a projective curve. Then  $C$  is compact.

Recall that affine curves are never compact.

*Proof.* Recall that a closed subset of a compact is compact itself. Since  $\mathbb{P}^2$  is compact by Exercise 12, we only need to prove that  $C$  is closed, or equivalently, that  $\pi^{-1}(C) \subseteq W = \mathbb{C}^3 \setminus \{0\}$  is closed, where  $\pi : W \rightarrow W/\sim = \mathbb{P}^2$  by  $(x_0, x_1, x_2) \mapsto (x_0, x_1, x_2)$ , by Remark 2.17. Assume that  $C = \{P = 0\}$  for a homogeneous polynomial  $P$ . Then,  $P : W \rightarrow \mathbb{C}$  is a continuous map, so that  $\pi^{-1}(C) = P^{-1}(\{0\}) = \{P(x_0, x_1, x_2) = 0\} \cap W$  is a closed subset of  $W$ . □

**Remark 4.15.** You may have heard of the Zariski topology for algebraic varieties. The Zariski topology is defined on affine space  $\mathbb{C}^n$  or on projective space  $\mathbb{P}^n$  as the topology whose closed sets are of the form  $V(S) = \{\underline{x} \in \mathbb{C}^n \mid f(\underline{x}) = 0\}$  where  $S$  is a set of polynomials or homogeneous polynomials  $f \in \mathbb{C}[x_0, \dots, x_n]$ . In this course, we do not work in the Zariski topology, but in the classical metric or Euclidean topology on  $\mathbb{C}^{n+1}$  and in the topology this induces on  $\mathbb{P}^n$  as outlined above. One reason for that is that the Zariski topology is not Hausdorff, or separated.

Lecture 6 is a problem class.

## 5 Affine vs projective plane curves

We now show how to projectivise affine curves, and study the relationship between affine and projective curves.

**Notation 5.1.** We will typically use coordinates  $x, y$  on  $\mathbb{C}^2$  and  $x_0, x_1, x_2$  on  $\mathbb{P}^2$ .

Starting with an affine curve  $C = \{f = 0\} \subseteq \mathbb{C}^2$  for  $f \in \mathbb{C}[x, y]$  non-constant, want to construct a projective curve  $\overline{C} \subseteq \mathbb{P}^2$ , a projectivisation of  $C$ . Recall that we have identified  $\mathbb{C}^2$  with the open subset  $U_2 = \{x_2 \neq 0\} \subseteq \mathbb{P}^2$ , via the function  $\phi : U_2 \rightarrow \mathbb{C}^2$  defined by  $[x_0, x_1, x_2] \mapsto (x_0/x_2, x_1/x_2)$  and its inverse  $(x, y) \mapsto [x, y, 1]$ . We now want to identify  $U_2 \cap \overline{C} \subseteq \mathbb{C}^2$  with the original curve  $C$  for a suitable projective plane curve  $\overline{C} = \{P = 0\} \subseteq \mathbb{P}^2$  for  $P$  homogeneous, picking  $\mathbb{C}^2 \cong U_2 \subseteq \mathbb{P}^2$  to be the points not at infinity.

**Example 5.2.** Idea is to let  $x = x_0/x_2$  and  $y = x_1/x_2$ .  $f(x, y) = x^3 + y + 1 = (x_0/x_2)^3 + (x_1/x_2) + 1$  and clear denominators to get  $P(x_0, x_1, x_2) = x_0^3 + x_1x_2^2 + x_2^3$ , homogeneous of degree three. Check that restricting  $P$  to  $U_2 \cong \mathbb{C}^2$  I get back  $f$ . Restricting  $P$  to  $U_2 \cong \mathbb{C}^2$  gives  $P(x, y, 1) = x^3 + y + 1$ .  $\{P = 0\} = \overline{C}$  is the projectivisation of  $\{f = 0\} = C$ .

**Theorem 5.3.** There is a bijective correspondence between

1. projective plane curves  $\overline{C} = \{P = 0\} \subseteq \mathbb{P}^2$  that do not contain the line at infinity  $\mathcal{P}_2 = \{x_2 = 0\}$ , and
2. affine curves  $C = \{f = 0\} \subseteq \mathbb{C}^2$ .

The bijection  $C \leftrightarrow \overline{C}$  is obtained by

$$P \mapsto [f : (x, y) \mapsto P(x, y, 1)], \quad f \mapsto [P : (x_0, x_1, x_2) \mapsto x_2^d \cdot f(x_0/x_2, x_1/x_2)],$$

for  $d = \deg(f)$ . The affine curve  $\{f = 0\} \subseteq \mathbb{C}^2$  is  $\phi(U \cap \overline{C})$ , where  $\overline{C} = \{P = 0\}$ .

**Notation 5.4.** In general,  $P$  is called the **homogenisation** of the polynomial  $f$ .  $\overline{C}$  is the **projectivisation** of  $C$ .

*Proof.* Let  $\overline{C} = \{P = 0\} \subseteq \mathbb{P}^2$  be a projective curve that does not contain  $\{x_2 = 0\}$  as in the statement.  $\{x_2 = 0\}$  is not contained in  $\overline{C}$  if and only if  $P$  contains at least one monomial without  $x_2$ . Then  $P$  contains at least one monomial without  $x_2$ , so that  $f : (x, y) \mapsto P(x, y, 1)$  is a polynomial in  $x, y$ . If not,  $P = x_2 \cdot Q$  and  $\{P = 0\} = \{x_2 = 0\} \cup \{Q = 0\}$ . So  $f(x, y)$  has degree equal to  $d = \deg(P)$ . Under the identification  $\phi : U \rightarrow \mathbb{C}^2$  defined in the previous section,

$$\phi(U \cap \overline{C}) = \{(x, y) \in \mathbb{C}^2 \mid f(x, y) = 0\}.$$

Conversely, if  $C = \{f = 0\}$ , then  $C$  is the image by  $\phi$  of the intersection of  $U$  and  $\overline{C}$ , where  $\overline{C} = \{P = 0\}$ , where the polynomial  $P$  is defined by  $P(x_0, x_1, x_2) = x_2^d \cdot f(x_0/x_2, x_1/x_2)$ . Check that  $P$  is a well-defined homogeneous polynomial of degree  $d$ .  $\square$

Look at an example where  $\overline{C}$  does contain  $\{x_2 = 0\}$ .

**Example 5.5.** Let  $P = x_0x_2^2 + x_1^2x_2 = x_2(x_0x_2 + x_1^2)$ . Then  $f(x, y) = P(x, y, 1) = x + y^2$ , so  $P(x_0, x_1, x_2) = (x_0/x_2 + (x_1/x_2)^2)x_2^2 = x_0x_2 + x_1^2$ .

**Remark 5.6.** To intersect  $\{P = 0\}$  and  $\{Q = 0\}$  in  $\mathbb{P}^2$ , solve  $P = 0$  and  $Q = 0$  in  $x_0, x_1, x_2$  to get homogeneous coordinates of points of intersection, where  $\underline{0}$  is not a valid solution and  $(\lambda x_0, \lambda x_1, \lambda x_2)$  is the same as  $(x_0, x_1, x_2)$ .

**Example 5.7.**

1. Let  $\overline{C}$  be the projective curve  $\overline{C} = \{P(x_0, x_1, x_2) = x_0^2 + x_1^2 + x_2^2 = 0\}$ . Then  $C = \phi(\overline{C} \cap U_2)$  is  $\{f = 0\} \subseteq \mathbb{C}^2$ , where  $f(x, y) = x^2 + y^2 + 1$ .

2. Let  $C$  be the affine curve  $x^2y + y - 1 = 0$ . Then  $\overline{C}$  is defined by  $x_2^3 \left( (x_0/x_2)^2 (x_1/x_2) + x_1/x_2 - 1 \right) = x_0^2x_1 + x_1x_2^2 - x_2^3$ .

**Exercise 16.** Let  $i = 1$  or  $2$ , recall that  $\phi_i$  denotes the homeomorphism  $\phi : U_i \rightarrow \mathbb{C}^2$ . Show that if  $\overline{C} \subseteq \mathbb{P}^2$  is a projective curve that does not contain  $\mathcal{P}_i = \{x_i = 0\}$ , then

$$C_i = \phi_i(\overline{C} \cap U_i) \subseteq \mathbb{C}^2$$

is an affine curve.

Intersect projectivisations to find points of intersection at infinity.

**Example 5.8.**

1. Consider the parallel lines

$$L_1 = \{x + y + 1 = 0\}, \quad L_2 = \{x + y - 1 = 0\}.$$

The corresponding projective lines are given by

$$\overline{L}_1 = \{x_0 + x_1 + x_2 = 0\}, \quad \overline{L}_2 = \{x_0 + x_1 - x_2 = 0\},$$

then solve to give

$$\begin{cases} x_0 + x_1 = -x_2 \\ x_0 + x_1 = x_2 \end{cases} \implies \begin{cases} x_0 + x_1 = 0 \\ x_2 = 0 \end{cases}.$$

Inside  $\mathbb{P}^2$ , the intersection  $L_1 \cap L_2$  consists of exactly one point

$$p = [1, -1, 0].$$

Thus, the two projective lines meet at one point  $p \in \mathcal{P}_0$ , that is  $p$  is a point at infinity.

2. Similarly, let

$$C = \{xy - 1 = 0\}, \quad L = \{x = 0\},$$

then the projectivisations are

$$\overline{C} = \{x_0 \cdot x_1 - x_2^2 = 0\}, \quad \overline{L} = \{x_0 = 0\},$$

then intersect to give

$$\begin{cases} x_0 = 0 \\ x_0x_1 - x_2^2 = 0 \end{cases} \implies \begin{cases} x_0 = 0 \\ x_2 = 0 \end{cases}.$$

As above, in  $\mathbb{P}^2$ ,  $\overline{C} \cap \overline{L}$  consists of a single point

$$\{[0, 1, 0]\}$$

lying on  $\mathcal{P}_2 = \{x_2 = 0\}$ , the hyperplane at infinity.

**Example 5.9.** Projective conics over  $\mathbb{R}$  are defined by degree two equations. In  $\mathbb{R}^2$ , smooth conics are three kinds, ellipses, hyperbolas, and parabolas. Passing to  $\mathbb{P}^2(\mathbb{R})$ , these three kinds of curves become the same. Ellipses have no new points at infinity, hyperbolas have two new points at infinity, and parabolas have one new point at infinity.

1. Let  $f(x, y) = x^2 - y^2 + 1$ . Projectivising,  $P(x_0, x_1, x_2) = x_0^2 - x_1^2 + x_2^2$ . Restrict to  $U_1 = \{x_1 \neq 0\} \subseteq \mathbb{R}^2$ .  $\overline{C} \cap U_1$ , the equation of the unit circle  $g(x, y) = x^2 - 1 + y^2 = x^2 + y^2 - 1$  is obtained by setting  $x_1 = 1$ .
2. Consider the parabola  $C$  with equation  $y = x^2$  in  $\mathbb{R}^2$ . Projectivising, this becomes the curve  $\overline{C} = \{x_1x_2 = x_0^2\} \subseteq \mathbb{P}^2(\mathbb{R})$ . The intersection with the line at infinity  $x_2 = 0$  gives  $x_0^2 = 0$ , so the only point is  $[0, 1, 0]$ . The square suggests some kind of tangency, and indeed  $\overline{C}$  is tangent to  $x_2 = 0$ . In the chart  $x_0 \neq 0$ , the equation becomes  $xy = 1$ , and we see a hyperbola. In the chart  $x_1 \neq 0$  we have again a parabola,  $y = x^2$ .

Lecture 7  
Monday  
22/10/18



3. Consider the unit circle  $C = \{x^2 + y^2 - 1 = 0\} \subseteq \mathbb{R}^2$ . Projectivising we obtain  $\overline{C} = \{x_0^2 + x_1^2 - x_2^2 = 0\} \subseteq \mathbb{P}^2(\mathbb{R})$ . The intersection with the line at infinity  $x_2 = 0$  is empty as we could expect. In the chart  $x_0 \neq 0$  we obtain the curve  $1 + x^2 - y^2 = 0$ , which is a hyperbola, and similarly in  $x_1 \neq 0$ .

**Exercise 17.** Let  $a, b, c, d, e, f \in \mathbb{C}$ , with  $(a, b, c) \neq (0, 0, 0)$ , and define

$$C = \{ax^2 + bxy + cy^2 + dx + ey + f = 0\}.$$

Define the projectivisation  $\overline{C}$  of  $C$  and determine its points on the line at infinity.

**Exercise 18.**

1. Show that there exists a unique projective line  $L \subseteq \mathbb{P}^2$  through two distinct points  $P, Q \in \mathbb{P}^2$ .
2. Show that two distinct projective lines in  $\mathbb{P}^2$  meet in exactly one point.

**Exercise 19.** Let  $P_1, P_2, P_3$  be three points of  $\mathbb{P}^2$ , and denote

$$P_i = [b_{i,1}, b_{i,2}, b_{i,3}], \quad i = 1, 2, 3$$

for their coordinates. Let  $B = (b_{i,j})$  be the associated  $3 \times 3$  matrix. Show that  $P_1, P_2, P_3$  lie in the same projective line if and only if  $\det(B) = 0$ .

## 6 Points at infinity

Given an affine curve  $C \subseteq \mathbb{C}^2$  and its projectivisation  $\overline{C} \subseteq \mathbb{P}^2$ , we study the points at infinity of  $C$ , that is the points of  $\overline{C}$  that do not lie on  $C$ . Let us first recall the notation in use. Denote by  $U_2$  the open set of  $\mathbb{P}^2$  defined by  $\mathbb{C}^2 \cong U_2 = \{x_2 \neq 0\} \subseteq \mathbb{P}^2$  and let  $\mathcal{P}_2 = \mathbb{P}^2 \setminus U_2 = \{x_2 = 0\}$  be the corresponding hyperplane or line at infinity. The homeomorphism  $\phi : \mathbb{C}^2 \rightarrow U_2$  is defined by  $(x, y) \mapsto [x, y, 1]$ , and its inverse  $\psi : U_2 \rightarrow \mathbb{C}^2$  is the map  $\psi : [x_0, x_1, x_2] \mapsto [x_0/x_2, x_1/x_2]$ . Let  $C$  be the affine curve

$$C = \{(x, y) \in \mathbb{C}^2 \mid f(x, y) = 0\},$$

where  $f \in \mathbb{C}[x, y]$  is a polynomial of degree  $d$  with no repeated factors. The projectivisation of  $C$  is then the curve  $\overline{C}$  such that  $\overline{C} \cap \{x_2 \neq 0\} = \phi(C)$ , or equivalently such that  $\psi(\overline{C} \cap U_2) = C$ . The curve  $\overline{C}$  is

$$\overline{C} = \{[x_0, x_1, x_2] \in \mathbb{P}^2 \mid P(x_0, x_1, x_2) = 0\},$$

where  $P$  is the homogeneous polynomial

$$P(x_0, x_1, x_2) = x_2^d f(x_0/x_2, x_1/x_2).$$

Let  $f_i \in \mathbb{C}[x, y]$  be the homogeneous polynomials of degree  $0 \leq i \leq d$  with

$$f(x, y) = f_0(x, y) + \cdots + f_d(x, y),$$

and note that

$$P(x_0, x_1, x_2) = f_0(x_0, x_1)x_2^d + \cdots + f_d(x_0, x_1).$$

**Definition 6.1.** The **points at infinity** of  $C$  are the points on  $\overline{C} \setminus \phi(C)$ . We will often just write  $\overline{C} \setminus C$ .

How many points at infinity can there be? If  $\{x_2 = 0\} \subseteq \overline{C}$ , if and only if  $x_2$  divides  $P(x_0, x_1, x_2)$ , then there are infinitely many. Assume  $\{x_2 = 0\} \not\subseteq \overline{C}$ . Look at  $P(x_0, x_1, 0)$ , and find solutions of  $P(x_0, x_1, 0) = 0$  in  $\mathbb{P}^1 \cong \mathcal{P}_2$ . The points at infinity of  $C$  are thus

$$\overline{C} \setminus C = \overline{C} \cap \mathcal{P}_2 = \{[x_0, x_1, 0] \in \mathbb{P}^2 \mid P(x_0, x_1, 0) = 0\}.$$

From the above equation, and under the identification  $\mathcal{P}_2 \cong \mathbb{P}^1$ , this becomes

$$\overline{C} \setminus C = \{[x_0, x_1] \in \mathbb{P}^1 \mid f_d(x_0, x_1) = 0\},$$

that is a projective variety in  $\mathcal{P}_2 \cong \mathbb{P}^1$ . It is natural to expect that such a projective variety consists of a finite number of points. This is what we will prove next.

**Lemma 6.2.** Let  $Q \in \mathbb{C}[x_0, x_1]$  be a non-zero homogeneous polynomial of degree  $d \geq 1$ . Then there exist complex numbers  $\alpha_i, \beta_i \in \mathbb{C}$  with  $(\alpha_i, \beta_i) \neq (0, 0)$  for  $i = 1, \dots, d$  such that

$$Q(x_0, x_1) = \prod_{i=1}^d (\alpha_i x_0 + \beta_i x_1).$$

Therefore,  $\{[x_0, x_1] \in \mathbb{P}^1 \mid Q(x_0, x_1) = 0\} = \{P_i = [-\beta_i, \alpha_i]\} \subseteq \mathbb{P}^1$ . Note that the  $P_i$  need not be distinct.

*Proof.* Let us pass to  $U_1 \cong \mathbb{C}^1 \subseteq \mathbb{P}^1$ , one of the two affine charts. Write  $Q(x_0, x_1) = \sum_{r=0}^d a_r x_0^r x_1^{d-r}$  and let  $e = \max\{r \mid a_r \neq 0\}$  so

$$Q(x_0, x_1) = x_1^d \cdot \sum_{r=0}^e a_r \left(\frac{x_0}{x_1}\right)^r.$$

Define  $f \in \mathbb{C}[x]$  as  $f(x) = \sum_{r=0}^e a_r x^r$ . By the fundamental theorem of algebra, there are complex numbers  $\lambda_1, \dots, \lambda_e$  such that

$$f(x) = a_e \cdot \prod_{i=1}^e (x - \lambda_i),$$

where  $\lambda_i$  are the roots of  $f$ , and hence

$$Q(x_0, x_1) = x_1^d \cdot a_e \cdot \prod_{i=1}^e \left(\frac{x_0}{x_1} - \lambda_i\right) = a_e \cdot x_1^{d-e} \cdot \prod_{i=1}^e (x_0 - \lambda_i x_1).$$

This proves the lemma, if we set

$$(\alpha_i, \beta_i) = \begin{cases} (1, -\lambda_i) & i \leq e \\ (0, a_e) & i = e+1 \\ (0, 1) & i > e+1 \end{cases}.$$

If  $e = d$ , then set

$$(\alpha_i, \beta_i) = \begin{cases} (1, -\lambda_i) & i < e \\ (a_e, -a_e \lambda_e) & i = e \end{cases}.$$

The description that  $\{Q = 0\} \subseteq \mathbb{P}^1$  is clear, once we note that  $(\alpha_i, \beta_i) \neq (0, 0)$ . □

We have proved the following.

**Theorem 6.3.** Let  $\overline{C} \subseteq \mathbb{P}^2$  be a projective curve defined by a polynomial of degree  $d$  that does not contain  $\mathcal{P}$ . Then  $\overline{C} \cap \mathcal{P}$  is a non-empty set of at most  $d = \deg(Q)$  points.

**Definition 6.4.** Let  $C = \{f = 0\} \subseteq \mathbb{C}^2$  for  $f \in \mathbb{C}[x, y]$  be an affine curve of degree  $d$  and let  $\overline{C}$  be its projectivisation, where, as usual,  $C = \psi(\overline{C} \cap U)$ . Denote

$$f = f_d + \dots + f_0, \quad \deg f_i = i, \quad f_d \neq 0,$$

the decomposition of  $f$  into its homogeneous polynomial parts, so that

$$P(x_0, x_1, x_2) = f_d(x_0, x_1) + \dots + f_0(x_0, x_1) x_2^d = \sum_{i=0}^d x_2^{d-i} f_i(x_0, x_1).$$

Then, if  $(\alpha_i, \beta_i) \in \mathbb{C}^2 \setminus \{0\}$  for  $i = 1, \dots, d$  are such that

$$f_d(x_0, x_1) = \prod_{i=1}^d (\alpha_i x_0 + \beta_i x_1),$$

the affine lines  $\{\alpha_i x + \beta_i y = 0\} \subseteq \mathbb{C}^2$  are the **asymptotes** of  $C$ . The points at infinity at  $x_2 = 0$  of  $C$  depend on the equation  $f_d(x_0, x_1) = 0$  and are

$$\overline{C} \setminus C = \{[-\beta_i, \alpha_i]\} \subseteq \mathcal{P}.$$

Each asymptote meets  $\overline{C}$  at the point  $[-\beta_i, \alpha_i] \in \mathbb{P}^1$ , the line at infinity. (TODO Exercise: check this)

**Remark 6.5.** It is clear from Lemma 6.2 that asymptotes are well-defined. Note that the asymptotes are lines in  $\mathbb{C}^2$  because  $(\alpha_i, \beta_i) \neq (0, 0)$ .