

Tetromino tilings with line clears

Jaehyeok Kim

September 19, 2026

Abstract

Walkup proved in 1965 that a $w \times n$ rectangle can be tiled by T-tetrominoes exactly when $4 \mid w$ and $4 \mid n$, and a line of later work measured how badly other rectangles fail. We ask the same question in a *dynamic* setting: pieces fall one at a time under gravity, and any row that becomes full is deleted. A *perfect clear* is a play that starts from the empty board and returns to it.

Line clears collapse the static obstruction almost completely. Four of the seven tetromino types — I, T, J and L — admit a perfect clear at every width, and O admits one at every even width; we determine the minimum number of pieces in each case, for all widths. The two exceptions are S and Z, which admit no perfect clear at any width, alone or together. The classification extends to every subset of the seven types.

The contrast with the static theory is sharpest for T, where Walkup’s condition reappears *inside* the dynamic construction: the part of the construction that uses no line clears is exactly a static tiling, so its width must be a multiple of four, and the line clears cover precisely the residues that static tiling forbids.

1 Introduction

A tetromino is a polyomino with four cells; there are seven of them up to translation, conventionally named I, O, T, S, Z, J and L. Which rectangles can be tiled by copies of a single one?

For the T-tetromino the answer is a classical theorem of Walkup [1]: a $w \times n$ rectangle admits a tiling by T-tetrominoes if and only if $4 \mid w$ and $4 \mid n$. Korn and Pak [3] later analysed the structure of those tilings. Rectangles that fail Walkup’s condition cannot be tiled at all, so Hochberg [2] asked how close one can get: the *gap number* is the least number of monominoes that must be left uncovered. He proved lower bounds of a strikingly rigid kind — width 5 forces at least one monomino every five columns, width 7 one every seven columns — and showed the gap number of T is one of 5, 6, 7, 9; Torres and Vallejo [4] later brought the cap down to six.

All of that is static. The tetrominoes are also the pieces of Tetris, and there the board is not a passive region: pieces fall under gravity, and a row that becomes completely full is deleted, with everything above it dropping by one. The natural question is what those deletions buy.

Definition 1. A *perfect clear* at width w is a play that begins with an empty board of width w and, after finitely many placements and the deletions they trigger, ends with the board empty again.

Our main result is that line clears dissolve the static obstruction for five of the seven pieces and for no others.

piece	static	dynamic (this paper)
I	$4 \mid w, 4 \mid n$ needed for a full tiling	every width, $n = w/4$ or w
O	only even widths	every even width, $n = w/2$
T	Walkup: $4 \mid w$ and $4 \mid n$	every width, $n = w$
J, L		every width, $n = w/2, w$ or $2w$
S, Z	conjectured unbounded gap number	never, at any width

The T row is where the two theories meet most sharply, and it is worth stating the mechanism now, because it is the shape of the whole paper. Our construction for T (Theorem 7) splits the board into column-disjoint *units*. All but the last unit must fill their rows without triggering any clear — that is, they must *statically tile* a $x \times 4$ rectangle, so Walkup forces $4 \mid x$. Widths 5, 6 and 7 therefore cannot be built this way at all. They are exactly the widths the final unit handles, and the final unit is the one allowed to use line clears. The dynamic rule covers precisely the residues the static theorem forbids.

Two further remarks place this work. Hoogeboom and Kusters [8] characterised which board configurations are reachable from empty when *all seven* piece types are available; with the full set the only obstruction is a divisibility condition on the number of cells, so the question is interesting only when the supply of pieces is restricted. That is the case we treat. Separately, Tetris is hard: deciding how much of a given board can be cleared is NP-complete [9], and this remains true for a single piece type [10]. Our question is different in kind — not the complexity of clearing a given board, but which piece sets can empty an *empty* board at all — and it turns out to have a clean answer.

2 The model

We fix the rules precisely, because one of our results is that they matter less than one might expect.

The board of width w is a finite set $B \subseteq \{0, \dots, w-1\} \times \mathbb{Z}_{\geq 0}$ of occupied cells; we write (r, c) for the cell in row r and column c , with row 0 at the bottom. Play starts from $B = \emptyset$. A move chooses a piece type from the available set P , chooses one of its rotations and a horizontal position, and places its four cells as a set X , subject to:

- (L1) $X \cap B = \emptyset$ — the four cells were empty;
- (L2) X shifted down by one row is not a legal position — the piece cannot fall further;
- (L3) after the placement, every full row is deleted simultaneously, and each surviving cell descends by the number of deleted rows below it.

Condition (L2) says a piece locks only when supported. It says nothing about how the piece reached its position. Real implementations differ here: hard drop (choose a column, fall straight down), soft drop with horizontal movement (allowing tucks), and rotation systems with wall kicks (allowing spins) permit successively more placements. Writing $\text{HD} \subseteq \text{SD} \subseteq \text{SRS}$ for the three sets of reachable placements, our results are insensitive to the choice:

- every construction in this paper was found under hard drop only, so it remains valid in the larger models;
- every impossibility proof uses only (L1), (L2), (L3) and the shape of the pieces, never the path by which a piece arrived, so it holds in all three.

Finally, if a perfect clear uses n pieces and deletes R rows in total, then counting cells gives

$$4n = Rw, \tag{1}$$

since every cell that is ever placed is eventually deleted, and each deletion removes w cells.

3 Two tools

The first tool does almost all of the work.

Lemma 2 (Column counting). *In a perfect clear at width w that deletes R rows in total, every column receives exactly R cells.*

Proof. A cell never changes column: placement fixes its column, and the descent in (L3) is vertical. A deletion removes one cell from each of the w columns, since the deleted row is full. The board ends empty, so every cell placed in column c is eventually deleted, and deletions remove cells from column c one at a time, once per deleted row. Hence column c receives exactly R cells. $\square$

Lemma 2 is purely combinatorial: it uses (L3) and nothing else, so it is independent of the rotation system. Summing over columns recovers (1).

To use it we record, for each piece type, its *column profiles*: the tuple of cells contributed to consecutive columns, one tuple per rotation.

piece	column profiles
I	(1, 1, 1, 1), (4)
O	(2, 2)
T	(1, 2, 1), (1, 3), (3, 1)
S, Z	(1, 2, 1), (2, 2)
J, L	(2, 1, 1), (1, 1, 2), (1, 3), (3, 1)

If x_c denotes the number of pieces placed with a given profile starting at column c , Lemma 2 becomes a system of linear equations in the x_c , one per column, each with right-hand side R . We refer to it as the *counting system*. Note that S and Z have the same profiles, as do J and L; counting alone can never distinguish them. The reason is structural.

Lemma 3 (Mirror). *The board and the rules are symmetric under left–right reflection, which exchanges J with L and S with Z. Consequently J and L behave identically at every width and every piece count, as do S and Z.*

Proof. Reflecting every placement of a legal play gives a legal play: (L1) and (L2) are preserved because reflection is a bijection on cells respecting rows, and a row is full if and only if its reflection is. Reflection maps the rotations of J onto those of L and the rotations of S onto those of Z. $\square$

4 Single piece types

Throughout this section P consists of one type.

4.1 S and Z

Theorem 4. *S alone admits no perfect clear, at any width. The same holds for Z alone.*

Proof. Let h_c be the number of horizontal S pieces (profile (1, 2, 1)) whose leftmost column is c , and v_c the number of vertical ones (profile (2, 2)). The counting system reads

$$h_c + 2h_{c-1} + h_{c-2} + 2v_c + 2v_{c-1} = R \quad (0 \leq c \leq w-1),$$

with the convention that a variable vanishes when its piece would leave the board. Column 0 gives $h_0 + 2v_0 = R$ and column 1 gives $h_1 + 2h_0 + 2v_1 + 2v_0 = R$; subtracting, $h_1 + h_0 + 2v_1 = 0$,

so $h_0 = h_1 = v_1 = 0$ and $2v_0 = R$. The same subtraction at columns $c+2, c+3$ propagates this: by induction $h_c = 0$ for all c , $v_c = 0$ for odd c , and $v_c = R/2$ for even c .

If w is odd, then $w-1$ is even, so $v_{w-1} = R/2$; but a vertical piece starting at column $w-1$ would occupy column w , so $v_{w-1} = 0$ and $R = 0$. There is no nonempty play.

If w is even the counting system is consistent, and we need the geometry. Every piece is a vertical S starting at an even column, so the columns split into independent blocks $\{2k, 2k+1\}$, no piece straddling two blocks. A vertical S occupies relative rows $\{1, 2\}$ in its left column and $\{0, 1\}$ in its right one: only the *right* column reaches relative row 0. Since every piece starts at an even column, every cell in an even column sits at relative row at least 1 above the piece's base, and the base is at absolute row ≥ 0 ; so no placement ever fills cell $(0, 0)$.

Nor can a cell descend into it. A cell reaches row 0 only if all the rows below its current position are deleted, which requires row 0 itself to be deleted, which requires $(0, 0)$ to be occupied. The two statements — “ $(0, 0)$ is never filled” and “row 0 is never deleted” — therefore support each other, and a simultaneous induction on the moves establishes both.

The first piece placed in the block $\{0, 1\}$ rests on the floor, since nothing else can occupy those columns, and so leaves a cell in row 0. Row 0 is never deleted, and a cell leaves the board only when its row is deleted; that cell is permanent, and the board never empties. The statement for Z follows from Lemma 3. $\square$

4.2 O and I

Theorem 5. *O alone admits a perfect clear exactly at even widths, with $n = w/2$ pieces and $R = 2$.*

Proof. With o_c the number of O pieces starting at column c , the counting system is $2o_c + 2o_{c-1} = R$. Column 0 gives $2o_0 = R$, column 1 gives $2o_1 = 0$, and inductively $o_c = R/2$ for even c and 0 for odd c . If w is odd then $o_{w-1} = 0$ because the piece would leave the board and $o_{w-2} = 0$ because $w-2$ is odd, so column $w-1$ gives $R = 0$. If w is even, place $w/2$ pieces side by side: rows 0 and 1 fill and are deleted. That $R = 2$ cannot be improved is part of Theorem 10. $\square$

Theorem 6. *I alone admits a perfect clear at every width w , with $n = w/4$ pieces and $R = 1$ when $4 \mid w$, and $n = w$ pieces and $R = 4$ otherwise.*

Proof. For the construction, place one vertical I in each column: rows 0 to 3 fill and are deleted, using $n = w$ pieces. If $4 \mid w$, lay $w/4$ horizontal I pieces along row 0 instead, deleting one row. The minimality of R in each case follows from Theorem 10 below. $\square$

4.3 T

This is the case where the static theory reappears, so we give it in full.

Theorem 7. *T alone admits a perfect clear at every width $w \geq 4$, and the minimum number of pieces is exactly $n = w$, with $R = 4$.*

Proof of the lower bound. By (1) it suffices to show $R \geq 4$. Let h_c, p_c, q_c count the pieces with profile $(1, 2, 1)$, $(1, 3)$ and $(3, 1)$ respectively starting at column c . The counting system is

$$N_j := h_j + 2h_{j-1} + h_{j-2} + p_j + 3p_{j-1} + 3q_j + q_{j-1} = R.$$

We use only $j = 0, 1, 2$, which exist as soon as $w \geq 3$.

If $R = 1$, then $N_0 = 1$ forces $(h_0, p_0, q_0) = (1, 0, 0)$ or $(0, 1, 0)$; the first gives $N_1 \geq 2h_0 = 2$ and the second $N_1 \geq 3p_0 = 3$, both exceeding 1.

If $R = 2$, then $3q_0 \leq 2$ forces $q_0 = 0$ and $(h_0, p_0) \in \{(2, 0), (1, 1), (0, 2)\}$; then $N_1 \geq 2h_0 + 3p_0$ equals 4, 5 or 6, all exceeding 2.

If $R = 3$, the solutions of $N_0 = 3$ are $(3, 0, 0)$, $(2, 1, 0)$, $(1, 2, 0)$, $(0, 3, 0)$ and $(0, 0, 1)$. For the first four, $N_1 \geq 2h_0 + 3q_0 \geq 6$. The remaining case has $q_0 = 1$ and $h_0 = p_0 = 0$, whence $N_1 = h_1 + p_1 + 3q_1 + 1 = 3$, so $q_1 = 0$ and $(h_1, p_1) \in \{(2, 0), (1, 1), (0, 2)\}$; but then $N_2 \geq 2h_1 + 3p_1 \geq 4 > 3$.

So $R \geq 4$ and $n = Rw/4 \geq w$. $\square$

Proof of the construction. Write the four rotations of T as

$$\begin{aligned} U &= \{(0, 0), (0, 1), (0, 2), (1, 1)\}, & D &= \{(0, 1), (1, 0), (1, 1), (1, 2)\}, \\ L &= \{(0, 1), (1, 0), (1, 1), (2, 1)\}, & R &= \{(0, 0), (1, 0), (1, 1), (2, 0)\}, \end{aligned}$$

with cells given as (row, column) offsets from the landing position, and write $X@c$ for a placement of X whose leftmost column is c .

We decompose the board into units occupying disjoint intervals of columns. Because a piece never leaves its unit's columns, the landing row of each placement, and whether a row of that unit is full, depend only on that unit's own contents. Two kinds of unit are needed, with $R = 4$ throughout:

The *stacking unit* S , of width 4, is $U@0, L@2, R@0, D@1$. With clears switched off it fills rows 0 to 3 of its four columns exactly, using 4 pieces.

The *final units* E_m , of width $m \in \{4, 5, 6, 7\}$ and m pieces:

$$\begin{aligned} E_4 &: U@0 \ L@2 \ R@0 \ D@1 \\ E_5 &: U@0 \ D@2 \ R@0 \ L@3 \ D@1 \\ E_6 &: U@1 \ L@4 \ D@2 \ R@0 \ D@0 \ D@3 \\ E_7 &: U@0 \ U@3 \ D@2 \ L@5 \ D@4 \ R@0 \ D@1 \end{aligned}$$

With clears switched on, each empties its own columns after deleting four rows.

Now write $w = 4a + m$ with $m \in \{4, 5, 6, 7\}$ and $a \geq 0$; this is possible for every $w \geq 4$, taking $m \equiv w \pmod{4}$ in the stated range. Play a copies of S in the columns $[0, 4a)$, then E_m in the columns $[4a, w)$.

While the copies of S are played, the columns of E_m are empty, so no row of the board is full and no deletion occurs; each copy of S therefore fills rows 0 to 3 of its own columns, as required. When E_m begins, every column outside it is full in rows 0 to 3, so a row of the board is full exactly when E_m fills that row within its own columns. Hence E_m evolves exactly as it would on a board of width m on its own, deleting four rows; those deletions remove the rows of the copies of S as well, and the board is empty. The piece count is $4a + m = w$, matching the lower bound. $\square$

Remark 8 (Where Walkup reappears). The requirement on a stacking unit — fill rows 0 to 3 of x columns with no deletion — says exactly that the $x \times 4$ rectangle admits a static tiling by T-tetrominoes. By Walkup's theorem this forces $4 \mid x$. So widths 5, 6 and 7 admit no stacking unit at all, and those are precisely the widths carried by the final unit, the one permitted to use line clears. The dynamic rule covers exactly the residues the static theorem forbids.

Walkup's theorem does not, however, settle whether the 4×4 tiling can be *reached* under gravity: support is not a concern of static tiling, and a tiling in which some piece must be inserted beneath another would be unusable here. That it can be reached — by the four placements listed above — is the one genuinely dynamic ingredient of the construction.

4.4 J and L

Theorem 9. *J alone admits a perfect clear at every width $w \geq 4$, with minimum piece count*

$$n = \begin{cases} w/2 & (w \equiv 0 \pmod{4}), \quad R = 2, \\ w & (w \equiv 2 \pmod{4}), \quad R = 4, \\ 2w & (w \text{ odd}), \quad R = 8. \end{cases}$$

The same holds for L .

Sketch. The construction is again a unit decomposition, with $A = (2, 1, 1)$, $B = (1, 1, 2)$, $P = (1, 3)$ and $Q = (3, 1)$ denoting the rotations of J by their profiles. The units are

AB	width 4,	$R = 2$,	A@0, B@1
PQ	width 2,	$R = 4$,	P@0, Q@0
PQPQ	width 2,	$R = 8$,	(P@0, Q@0) twice
U_5	width 5,	$R = 8$,	A@0 P@1 P@0 P@3 Q@0 Q@3 A@2 A@0 P@3 B@1

and the board is built as $AB^{w/4}$ when $4 \mid w$, as $PQ^{w/2}$ when $w \equiv 2 \pmod{4}$, and as U_5 followed by $(w-5)/2$ copies of PQPQ when w is odd. The verification is the same two steps as in Theorem 7 and is a finite check on four units, independent of w . Minimality follows from Theorem 10. L follows by Lemma 3. $\square$

5 Lower bounds at every width

The minimality claims above assert, for each piece and each width, that the counting system has no solution for smaller R . For a fixed width this is a finite check. The following observation upgrades it to all widths at once.

Theorem 10. *For each piece type and each R , the set of widths at which the counting system is solvable is eventually periodic in w , and the period can be computed. The resulting minimum values of R are*

piece	minimum R
I	1 if $4 \mid w$, else 4
O	2 if w even; no solution if w odd
T	4 at every width
S, Z	2 if w even; no solution if w odd
J, L	2 if $4 \mid w$; 4 if $w \equiv 2 \pmod{4}$; 8 if w odd

Proof. Sweep the columns from left to right, maintaining as state the vector of contributions already committed to the columns not yet processed. A profile spans at most four columns, so the state has at most three components, each an integer in $[0, R]$ — any larger value can never be reduced and is dead. The state space is therefore finite. Processing one column is a fixed relation on states, so it induces a deterministic map on *sets* of states; iterating it from the initial set $\{(0, \dots, 0)\}$ gives a sequence of subsets of a finite set, which must eventually cycle.

The rightmost few columns are special — a profile may no longer fit — but the sequence of restricted maps applied there depends only on how many columns remain, not on w . Hence solvability at width w is obtained by applying a fixed tail map to the $(w-k)$ -th term of an eventually periodic sequence, where $k \leq 3$ is the number of special columns, and is therefore eventually periodic in w . Computing the pre-period and period gives the table. $\square$

The entries for S and Z deserve emphasis, because they explain the shape of Theorem 4.

Corollary 11. *At every even width, S alone passes the counting test with $R = 2$. Consequently no argument based on Lemma 2 alone can rule out S , and the geometric step in the proof of Theorem 4 is unavoidable.*

Among all seven pieces, this is the only place where counting and geometry part company: for I, O, T, J and L the counting bound is achieved by an explicit construction, and for S and Z at odd widths counting already gives a contradiction. The single gap is S and Z at even widths, and it is exactly the gap Theorem 4 fills.

6 Sets of pieces

Suppose now that a set P of types is available, and the player may use any of them in any order.

Theorem 12. $\{S, Z\}$ together admits no perfect clear, at any width.

Sketch. S and Z have the same column profiles, so the counting argument of Theorem 4 applies verbatim to their total counts: no horizontal piece is used, every piece is vertical and starts at an even column, and odd widths are excluded outright. At even widths the columns again split into independent blocks $\{2k, 2k+1\}$, and the width- w problem reduces to a width-independent question about one block.

Let F_0, F_1 be the sets of filled rows in the two columns of a block and put $d_j = |F_0 \cap [0, j]| - |F_1 \cap [0, j]|$. Placing a vertical Z at height r increments d_r and d_{r+1} ; placing a vertical S decrements them; deleting row m removes d_m from the sequence and changes no other value. One then checks four invariants of the reachable states: $|F_0| = |F_1|$; the occupied rows form an initial segment; $|d_j| \leq 1$ for all j ; and $d_0 \neq 0$ whenever the block is nonempty. The last one says row 0 of the block is never full, so it is never deleted, and the cell the first piece leaves in row 0 is permanent. $\square$

Theorem 13. A set P of tetromino types admits a perfect clear at width w if and only if P meets $\{I, T, J, L\}$, or w is even and $O \in P$.

Proof. If P contains one of I, T, J, L , use that type alone (Theorems 6, 7, 9). Otherwise $P \subseteq \{O, S, Z\}$. All three of these have profile set $\{(1, 2, 1), (2, 2)\}$, so the counting derivation in the proof of Theorem 4 applies to their total counts without change: horizontal pieces are unused, every piece starts at an even column, and at odd widths $R = 0$. At even widths, if $O \in P$ then $w/2$ copies of O clear two rows; if not, then $P \subseteq \{S, Z\}$ and Theorems 4 and 12 apply. $\square$

Remark 14. It is tempting to argue that since $\{S, Z\}$ is the only unusable pair, every set of three or more types must work. That is false: $\{O, S, Z\}$ has three types and fails at every odd width. The reason is that O carries a width restriction of its own, so “only one bad pair” says nothing about sets built from the three restricted types.

7 Restricted supply: the seven-bag randomiser

Modern Tetris does not let the player choose the next piece. Pieces arrive in *bags*: each block of seven consecutive pieces is a permutation of the seven types. A single slot of *hold* lets the player set one arriving piece aside.

This changes the question, because every construction above repeats one type.

Theorem 15. Under the seven-bag constraint, with the order inside each bag at the player’s choice, the minimum number of pieces in a perfect clear is

w	4	5	6	7	8	9	10	11	12	13	14
unrestricted	1	5	3	7	2	9	5	11	3	13	7
seven-bag	1	5	3	7	4	9	10	11	9	13	14

The pattern is that the constraint is free at odd widths and costly at even ones. At an odd width the unrestricted minimum already has $R = 4$, and four rows give room to place seven different types; at an even width the unrestricted optimum is one or two rows, and a thin region cannot absorb variety. A single row admits only horizontal I pieces, so $w/4 \geq 2$ is immediately fatal; two rows survive to $w = 8$ and fail from $w = 10$.

Width 16 continues the pattern and is the most expensive case we have settled: the unrestricted minimum is 4, and the seven-bag minimum is 12. Both $n = 4$ and $n = 8$ are impossible — every admissible multiset was decided — while $n = 12$ admits five, among them one copy each of I and L with two of every other type.

The harder question is what the *order* inside the bags costs, and here we can report less. Fix $w = 10$ and $n = 10$, the setting of the standard opener. Ten placements consume at most eleven arrivals, so only the first four pieces of the second bag matter, giving $P(7, 4) = 840$ prefixes to consider for each of the $7! = 5040$ orders of the first bag.

We sampled 100 first bags uniformly from the 5040 with a fixed seed and settled all 840 prefixes for each: 84,000 cases. Nine fail. They fall in three first bags, three apiece:

LJTIOZS : OLSZ, LOSZ, LSOZ
 LOTJISZ : OJZS, JOZS, JZOS
 TLJOISZ : OJZS, JOZS, JZOS

The other 97 first bags admit a perfect clear for every prefix, so the failure rate over the sample is $9/84,000 \approx 1.1 \times 10^{-4}$. An earlier first bag, TJLISZO, was settled separately and fails on four prefixes (OJZS, OLZS, LOZS, JOZS).

One structural fact halves the work for anyone repeating this.

Lemma 16. *The hold slot is empty on an initial segment of the play and never afterwards, so a play stashes at most once.*

Proof. A move either places the arriving piece and leaves the hold unchanged, or places the held piece and puts the arriving one in its place, or — only from an empty hold — stashes the arriving piece without placing anything. The last two leave the hold occupied, and a stash requires it empty. $\square$

Write $\text{HG}_m(h; s_1 s_2 \dots)$ for the set of length- m sequences the player can place starting from hold h with $s_1, s_2, \dots$ arriving, each step placing either the held piece (the arrival replacing it) or the arrival (the hold unchanged).

Theorem 17. *If the arrival sequence $q_0 q_1 \dots q_{L-1}$ has $L \geq n + 1$, then the set of placement sequences of length n the player can achieve is*

$$q_1 \cdot \text{HG}_{n-1}(q_0; q_2 q_3 \dots) \cup q_0 \cdot \text{HG}_{n-1}(q_1; q_2 q_3 \dots),$$

which is symmetric in q_0 and q_1 . Exchanging the first two arrivals therefore changes neither the achievable placement sequences nor whether a perfect clear exists.

Proof. The two terms are the two continuations after stashing q_0 immediately, so they are achievable. For the converse, note that inside HG the move “place the held piece, keep the arrival” repeated from hold q_1 emits $q_1, q_2, q_3, \dots$ in order. By Lemma 16 a play either never stashes, giving $q_0 q_1 \dots q_{n-1}$, which is q_0 followed by $n - 1$ such repetitions and needs the arrival q_n — the one place where $L \geq n + 1$ is used — or stashes once, at q_k say, giving $q_0 \dots q_{k-1}$ followed by $\text{HG}_{n-k}(q_k; q_{k+1} \dots)$; for $k = 0$ this is the displayed union, for $k = 1$ its second term, and for $k \geq 2$ it is reached inside that second term by $k - 1$ repetitions before branching freely. $\square$

So the 5040 first bags fall into 2520 classes of two, and a bad first bag always has a bad partner. The hypothesis is not decorative: with $L = n$ no stash is possible and the play is forced, and among sequences of four pieces at width 4 there are 66 pairs differing only in their first two entries where one clears and the other does not.

Theorem 17 also explains a stronger effect we first met as a coincidence. Exchanging q_0 and q_1 leaves the stash-immediately subtree identical state by state, so a search that stashes first and

succeeds inside that subtree visits exactly the same states for both orders; across 1,684 solvable cases the two orders agreed on the state count to the digit. The no-stash branch is *not* identical — an I placed before an O reaches boards an O placed before an I cannot — and it is explored only when the search fails. On the first failing pair we tested, the two orders agreed on the verdict, as the theorem requires, and differed in state count, 38,542,538 against 38,537,388.

A caution about the computation is worth recording, because it changed the answer. Our first pass used a state cap of 8×10^6 and reported *no* failures at all: every one of the nine was sitting in a case that hit the cap, undecided. They need between 3.8×10^7 and 5.2×10^7 states to settle. A capped search is not a negative result.

Across all thirteen known failures the second-bag prefix is a permutation of $\{O, S, Z\}$ together with one of J or L — no exception so far. A tempting sharper rule, that the failing prefixes end in ZS, is false: the endings observed are ZS eight times, SZ twice, OS twice and OZ once, and the prefix LSOZ fails after LJTIOSZ but succeeds after TJLISZO. Failure is a joint property of the two bags, not of the second one alone.

What the four failing first bags do share is that S and Z both arrive late — in position five or later. Fifteen of our hundred samples have that property and three of them fail; the other eighty-five do not fail at all. The condition also survived a test made before the computation finished: the last four undecided cases all lay in first bags where S or Z arrives early, so the condition predicted they would succeed, and all four did. This is consistent with the mechanism one would guess — a vertical I is the main way to resolve an S/Z staircase inside four rows, and an I arriving after the staircase has been built cannot undo it, which is the direction Burgiel’s theorem [6] points in.

We claim no novelty for this section. The community’s own perfect-clear solvers work in a more general model — ours is hard-drop only, so an impossibility here is an impossibility *in our model*, and spins or tucks might rescue some of these cases. The point is the shape of the answer: perfect clears are almost always available, failures are rare and structured, and the well-known openers exist to save search and memorisation rather than to reach positions that are otherwise unreachable.

A A parity identity

The following is folklore among competitive players, where it is used to decide whether a perfect clear is still reachable; see the Hard Drop wiki [12]. We record it because it has a short proof, generalises to every width, and explains a natural false conjecture.

Colour the board like a chessboard and set

$$\varphi(B) = \sum_{(r,c) \in B} (-1)^{r+c}.$$

Every tetromino except T covers two cells of each colour, so placing it leaves φ unchanged; a T covers three of one colour and one of the other, changing φ by ± 2 , hence by $2 \bmod 4$.

Consider a deletion event removing full rows $j_1 < \dots < j_k$. For a surviving cell at row r , let $s(r)$ be the number of deleted rows below it; the cell descends by $s(r)$, multiplying its contribution by $(-1)^{s(r)}$. Put

$$D = \#\{\text{surviving cells with } s(r) \text{ odd}\}, \quad A = \sum_{i=1}^k \varphi(\text{row } j_i),$$

noting that φ of a full row is 0 when w is even and $(-1)^j$ when w is odd. Then

$$\Delta\varphi = -2 \sum_{s(r) \text{ odd}} (-1)^{r+c} - A \equiv -2D - A \pmod{4},$$

since modulo 4 the first sum depends only on the parity of the number of cells involved. Summing over a whole perfect clear, where φ starts and ends at 0, gives $2 \#T \equiv 2 \sum D + \sum A \pmod{4}$, that is

$$\#T \equiv \sum_{\text{events}} D + \frac{1}{2} \sum_{\text{events}} A \pmod{2}.$$

For even w the correction A vanishes and the identity reads: *the parity of the number of T pieces equals the parity of the total number of cells left above the deleted rows*. In particular a perfect clear with no deletions at all uses an even number of T pieces — which is the false conjecture one arrives at by forgetting that rows disappear.

Remark 18. D must be counted per deleted row, or equivalently as the number of cells with an odd number of deleted rows below them. Reading it as “the cells above the deleted rows” is correct only for single-row deletions.

Computations

All constructions and all counting claims in this paper are reproducible from the accompanying repository, which contains for each numerical statement the command that produces it. The identity of Appendix A was checked against all 4,758,805 perfect clears at widths 4, 5, 6, 7 and 8, by two independent implementations agreeing on the number of plays in every case.

References

- [1] D. W. Walkup, Covering a rectangle with T -tetrominoes, *Amer. Math. Monthly* **72** (1965) 986–988.
- [2] R. Hochberg, The gap number of the T -tetromino, *Discrete Math.* **338** (2015) 130–138.
- [3] M. Korn and I. Pak, Tilings of rectangles with T -tetrominoes, *Theoret. Comput. Sci.* **319** (2004) 3–27.
- [4] Torres and Vallejo, Fragments that reduce the gap cap of the T -tetromino to six, *Sci. Eng. J.* **17** (2024) 128–133.
- [5] M. Reid, Klarner systems and tiling boxes with polyominoes, *J. Combin. Theory Ser. A* **111** (2005) 89–105.
- [6] H. Burgiel, How to lose at Tetris, *Math. Gaz.* **81** (1997) 194–200.
- [7] J. Brzustowski, *Can you win at Tetris?*, MSc thesis, University of British Columbia, 1992.
- [8] H. J. Hoogeboom and W. A. Kusters, How to construct Tetris configurations, *Int. J. Intell. Games Simul.* **3** (2004) 97–105.
- [9] E. D. Demaine, S. Hohenberger and D. Liben-Nowell, Tetris is hard, even to approximate, in *COCOON 2003*, 351–363.
- [10] MIT Hardness Group, Tetris is hard with just one piece type, in *FUN 2026*, LIPIcs **366**, 32:1–32:16.
- [11] D. Baccherini and D. Merlini, Combinatorial analysis of Tetris-like games, *Discrete Math.* **308** (2008) 4165–4176.
- [12] Parity, Hard Drop Tetris Wiki.