Thesis Title

Thesis Subtitle



Joab Gray Kloppers

Department of Electrical Engineering
University of Cape Town
Rondebosch, Cape Town
South Africa


Supervisor
RA **Verrinder**

Co-supervisor(s)
A/Prof D **Pillay**, G **Maniel**

August 2026

BSc (Eng) final year project report submitted in partial fulfilment of the requirements for the degree of Bachelor of Science in Electrical and Computer Engineering in the Department of Electrical Engineering at the University of Cape Town

Keywords: keyword1; keyword2; keyword3



Declaration

I, **Joab Gray Kloppers**, hereby:

1. know that plagiarism is wrong. Plagiarism is to use another's work and pretend that it is one's own.
2. have used the **Harvard** convention for citation and referencing. Each contribution to, and quotation in, this report from the work(s) of other people has been attributed, and has been cited and referenced. Any section taken from an internet source has been referenced to that source.
3. declare that:
    a. this report is my own work, and apart from the normal guidance from my supervisor(s), I have received no assistance except as acknowledged.
    b. neither the substance nor any part of the above report has been submitted in the past, or is being, or is to be submitted for a degree at this University or at any other university.
    c. I have not paid a third party to complete my work on my behalf.
    d. my use of artificial intelligence software has been limited to:

    (specify which tool was used and how you used AI to assist with this project).

4. have not allowed and will not allow anyone to copy my work with the intention of passing it off as their own work.
5. acknowledge that copying someone else's work, or part of it, is wrong, and declare that this is my own work.
6. grant the University of Cape Town free license to reproduce the above thesis in whole or in part, for the purpose of research.

Word count [run latexmk] words (excluding preamble, appendices and bibliography)
Page count 12 pages (excluding preamble, appendices and bibliography)



Joab Gray Kloppers
Department of Electrical Engineering
University of Cape Town

Tue 11 August, 2026

Abstract

Thesis Title

Joab Gray **Kloppers**

Tue 11 August, 2026

Acknowledgements

Contents

Declaration . . . . . **i**
Abstract . . . . . **ii**
Acknowledgements . . . . . **iii**
List of Figures . . . . . **v**
List of Tables . . . . . **vi**
List of Symbols . . . . . **vii**
List of Symbols . . . . . **vii**
List of Acronyms and Glossary . . . . . **viii**
Chapter 1 Introduction **1**
1.1 Background 1
1.1.1 Biological Activity 1
1.1.2 Technical application 1
1.2 Problem statement 2
1.3 Project objectives 2
1.4 Scope, limitations and assumptions 2
1.5 Plan of development 2
Chapter 2 Literature review **3**
2.1 Biology 3
2.2 Technology 3
Chapter 3 Theoretical development **5**
Chapter 4 Design description **6**
Chapter 5 Experimental design **7**
Chapter 6 Results **8**
Chapter 7 Discussion **9**
Chapter 8 Conclusions **10**
Chapter 9 Further work **11**
Chapter A GitHub repository **12**
A.1 Title for appendix A.1 12
Appendix B: Ethics approval **13**
Appendix C: Topic description **14**
Appendix D: ECSA Graduate Attributes **15**
Project mapping to ECSA Graduate Attributes 15
Bibliography **16**

List of Figures

1.1 A simplified block diagram of a general communication system. 1

2.1 Custom incubation and flow through chamber by Schenone et al [1] 4

List of Tables

1.1 Table title 2

List of Symbols

Symbol	Description	Unit
$\boldsymbol{B}$	Magnetic flux density	T
$\boldsymbol{E}$	Electric field	V/m
$\boldsymbol{J}$	Current density	A/m^2
t	Time	s
ε_0	Permittivity of free space (8.85×10^{-12})	$C^2/N{\cdot}m^2$
μ_0	Permeability of free space ($4\pi \times 10^{-7}$)	N/A^2

Chapter 1

Introduction

1.1 Background

1.1.1 Biological Activity

Sand prawn activity has a large effect on the surrounding ecosystem. Pillay and Branch [2] explain that by physically burrowing into the sediment, they distribute buried sediment to the surface, alter the surface structure via the introduced mounds, which can increase surface area, and alter both chemical and structural properties of the sediment through burrowing and nutrient mixing. Venter et al [3] further shows that sand prawn bi-directional water pumping filters can reduce phytoplankton biomass by up to 70% in estuarine environments which can help with ecosystem resilience.

1.1.2 Technical application

Schenone et al [1] demonstrates that surface microtopography can be used as a reliable metric to predict biochemical processes. However common metrics for measuring surface roughness characteristics such as (list them here) are often invasive and result in the damage of the very structures under measurement. Azhar et al [4] demonstrated that the use of low cost RGB-D cameras can be used to produce depth images of the sediment surface, which can then be processed to analyse the surface properties.

Sub-subsection




Figure 1.1: A simplified block diagram of a general communication system. A communication system can be abstracted as follows: an *information source* produces a message to be sent through a *transmitter* which converts it to a signal which can be sent via a communication *channel*. The received signal is noisy and is converted back to the message by the *receiver* which is then delivered to the *destination*. Taken from shannon1948

Table 1.1: Table title. Extended description of tables contents.

Heading 1	**Heading 2**	
	Subheading 1 [units]	*Subheading 2* [units]

1.2 Problem statement

Equations are numbered and include the chapter number. As they are considered part of the text, they include punctuation, and each parameter must be described in the text below. Ensure that the units are stated and where possible use SI units. For example, the Maxwell equations are expressed as:

$$\nabla \cdot \boldsymbol{E} = \frac{\rho}{\epsilon_0}, \tag{1.1}$$

$$\nabla \cdot \boldsymbol{B} = 0, \tag{1.2}$$

$$\nabla \times \boldsymbol{E} = -\frac{\partial \boldsymbol{B}}{\partial t}, \tag{1.3}$$

$$\nabla \times \boldsymbol{B} = \mu_0 \boldsymbol{J} + \mu_0 \epsilon_0 \frac{\partial \boldsymbol{E}}{\partial t}, \tag{1.4}$$

where $\boldsymbol{B}$ is the magnetic flux density in T, $\boldsymbol{J}$ is the total current density in A/m^2, $\boldsymbol{E}$ is the electric field in V/m, μ_0 is the permeability of free space ($4\pi \times 10^{-7}$ H/m), and ϵ_0 is the permittivity of free space (8.85 $\times 10^{-12}$ C^2/N·m^2).

1.3 Project objectives

The Global Positioning System (GPS) is used for positioning. The Marginal Ice Zone (MIZ) is of particular interest in the Southern Ocean (SO).

1.4 Scope, limitations and assumptions

1.5 Plan of development

Chapter 2

Literature review

2.1 Biology

Pillay and Branch [2] describe how sand prawn species are prominent members in the benthic ecosystem. The sand prawns continuously change the sediment properties such as surface topography porosity and biochemistry, through actions such as burrowing and water pumping. These actions redistribute buried sediment to the surface which results in surface mounds which increase the effective surface area of the sediment by up to 15 times. Kristensen et al [5] further explains how thalassinid shrimp populations can turn up to 12kg of sediment per square meter per day. These mounds can increase the effective hydrodynamic resistance which can help to stagnate the water flow close to the sediment. While the paper mentions measurements specific to sand prawn species located in other regions, they provide an insight into approximate feature sizes that can be expected, for example *Callianassa subterranea* present in the North sea can produce mounds 5.4cm tall and 11.7cm wide, and *Pestarella candida*, located in the Mediterranean and Adriatic Seas, which can produce mounds 3cm tall and 15cm wide.

Venter et al [3] further explains how sand prawns, *Callianassa kraussi*, present in the South African coastlines and at the Langebaan Lagoon, function as prominent water filterers. Within the burrows they actively irrigate the sediment through bi-directional pumping of water to replenish oxygen levels within the burrows, and to remove waste products. This irrigation has a secondary effect where phytoplankton are filtered out of the water by the burrow walls. This filtering can help improve ecosystem resilience to harmful conditions such as nutrient pollution which promote phytoplankton blooms. In relation to this project, the papers methodology entails manual visual counting of burrow openings.

The next question with regards to biological activity, is weather surface roughness, specifically rugosity, is a valid measurement to gauge the underlying biological activity within the benthic ecosystem. To this, Schenone et al [1, 6] determined experimentally that analysing two dimensional surface rugosity, they were able to correlate surface roughness measurements with sub-surface activity. One paper [1] described the testing of a controlled laboratory environment to simulate the benthic ecosystem, which can be seen in Figure 2.1. They found that the simple two dimensional rugosity estimate managed to account for up to 61% of nutrient variability. The second paper[6] describes a more complex implementation, with the use of a single diver operated camera recording video of the sediment surface, and the use of visual based structure from motion (SFM) photogrammetry to obtain a three dimensional representation of the surface. similarly, the find that sediment microtopography explains up to 55% of species richness variance, which is of particular interest with regards to this project. These papers show that surface rugosity is a useful measurement when gauging biological activity, and they provide a useful insight into the computational pipeline which is used to convert from spatial meshes to elevation matrices, however there are considerations that need to be considered, these papers discuss fully submerged environments, thus the requirement of under water cameras and SFM, whilst this project focuses on tidally exposed surfaces, where fixed clear observations can be made with the use of a rig.

2.2 Technology

With the understanding that surface roughness statistics can provide a feasible proxy for benthic activity, there is a need for a method to scan sediment surfaces and extract topographical information. The current project, under the biological side <**PLEASE CHANGE THIS, SOMEHOW TIE IN THE EXISTING PROJECT**>, takes surface scans of static, exposed sediment, within the inter-tidal region in the Langebaan lagoon. These conditions mean that there is no need for submersion rated hardware, operated under dynamic conditions.




Figure 2.1: Custom incubation and flow through chamber by Schenone et al [1]

The current project is based off the work developed by Azhar et al [4]. The authors develop a framework that allows for extracting a millimetre scale 3D surface scans non-invasively, and at lower price scales than expensive LiDAR based systems. The system makes use of an off the shelf Asus Xiton pro Live 3D RGB-D structured light camera, mounted within an opaque plastic container to block infrared interference from sunlight. They then tested four polynomial de-trending algorithms to remove natural sediment gradients, and isolate the bioturbation. Due to the nature of the Asus Xiton, which makes use of structured light projection and a single IR camera, the system suffers when the surface is exposed to sunlight. Therefore the authors made use of an opaque plastic enclosure which is set on the sediment for scanning, however this limits the spatial resolution to only $0.3m^2$ with an effective camera resolution of only 240 x 200 pixels. Additionally the authors found that the higher order polynomial de-trending algorithms were sensitive to elevations at the edges of the image frame, requiring them to use a planar model which proved to be more robust.

Chapter 3

Theoretical development

Chapter 4

Design description

Chapter 5

Experimental design

Chapter 6

Results

Chapter 7

Discussion

Chapter 8

Conclusions

Chapter 9

Further work

Appendix A

GitHub repository

This appendix provides a link to the GitHub repository where the code and additional materials for this final year project report can be found.

A.1 Title for appendix A.1

- GitHub Repository: Repo name

Appendix B: Ethics approval

Appendix C: Topic description

The original topic description used during the project selection process is included below. This topic has since been further refined, and the exact problem that was investigated is described in Section 1.2.

Appendix D: ECSA Graduate Attributes

Project mapping to ECSA Graduate Attributes

The table below provides a structured reflection on where each of the required Graduate Attributes (GAs) as defined by Engineering Council of South Africa (ECSA) associated with the final year project is met. This includes reference to the specific Chapter/Section (where applicable).

GA	Graduate Attribute Description	Student reflection: How this outcome was achieved during the project
GA1	*Problem solving*	
GA4	*Investigations, experiments and data analysis*	
GA5	*Use of engineering tools*	
GA6	*Professional and technical communication*	
GA8	*Individual and collaborative teamwork*	
GA9	*Independent learning ability*	

Bibliography

[1] S. Schenone, I. Bartl, and S. F. Thrush, "Sediment microtopography predicts localised benthic ecosystem functioning and the effect of species interactions," *Marine Ecology Progress Series*, vol. 749, pp. 47–55, Nov. 2024.

[2] D. Pillay and G. Branch, "Bioengineering effects of burrowing thalassinidean shrimps on marine soft-bottom ecosystems," in *Oceanography and Marine Biology: An Annual Review*, Jun. 2011, vol. 49, pp. 137–192.

[3] O. Venter, D. Pillay, and K. Prayag, "Water filtration by burrowing sandprawns provides novel insights on endobenthic engineering and solutions for eutrophication," *Scientific Reports*, vol. 10, no. 1, p. 1913, Feb. 2020.

[4] M. Azhar, J. R. Hillman, T. Gee, S. Schenone, W. van der Mark, S. F. Thrush, and P. Delmas, "An RGB-D framework for capturing soft-sediment microtopography," *Methods in Ecology and Evolution*, vol. 13, no. 8, pp. 1730–1745, 2022.

[5] E. Kristensen, G. Penha-Lopes, M. Delefosse, T. Valdemarsen, C. O. Quintana, and G. T. Banta, "What is bioturbation? The need for a precise definition for fauna in aquatic sciences," *Marine Ecology Progress Series*, vol. 446, pp. 285–302, Feb. 2012.

[6] S. Schenone, J. E. Hewitt, J. Hillman, R. Gladstone-Gallagher, J. Gammal, C. Pilditch, A. M. Lohrer, E. Ferretti, M. Azhar, P. Delmas, and S. F. Thrush, "Seafloor sediment microtopography as a surrogate for biodiversity and ecosystem functioning," *Ecological Applications*, vol. 35, no. 1, p. e3069, 2025.