

# Supporting Information: The Transport Mechanisms of Polar Solutes in a Cross-linked H<sub>II</sub> Phase Lyotropic Liquid Crystal Membrane

Benjamin J. Coscia

Michael R. Shirts

March 11, 2019

## S1 Setup and analysis scripts

All python and bash scripts used to set up systems and conduct post-simulation trajectory analysis are available online at [https://github.com/shirtsgroup/LLC\\_Membranes](https://github.com/shirtsgroup/LLC_Membranes). Documentation for the LLC\_Membranes repository is available at <https://llc-membranes.readthedocs.io/en/latest/>. Table S1 provides more detail about specific scripts used for each type of analysis performed in the main text.

## S2 Water content equilibration

We initially attempted to equilibrate our system with water by allowing water molecules to naturally penetrate the membrane from a water bath separating periodic images of the system in the  $z$ -direction (See Figure S1a). We allowed a dry, previously equilibrated system to further equilibrate in coexistence with a 3 nm-thick (in the  $z$ -direction) layer of water. Water readily enters the tail region where the density of monomers is low. About 3 times more water molecules occupy the tail region after 1000 ns of equilibration (See Figure S1b). Although the water level in the pore appears to plateau in this system, it is clear that equilibration of this system is kinetically limited since water does not fill the pores uniformly. The density of water along the pore axis, averaged over the last 50 nanoseconds of simulation, is close to zero at the membrane center. Therefore, we required a different equilibration technique in order to overcome the kinetic limitation.

We equilibrated 4 systems where we initially placed water in the pores and in the tails in addition to a water reservoir between periodic images. We filled the pores with water by running `gmx solvate` on our initial configuration and removing any water molecules placed outside of the pore region. The GROMACS command `gmx solvate` places water molecules in proximity to other atoms based on their van der waals radii and therefore does a decent job of preventing equilibration issues. This also means that the initial pore radius dictates the water content of the pore. We can however, put arbitrary amounts of water into the tail region. We chose to test systems with initial pore radii of 5, 6, 7 and 8 Å with tail and pore water compositions given in Table S2.

Systems appear to be most stable when there is more water in the pores than tails. In systems started with more water in the tails (Figures S2a and S2b), the pore water content tends to increase over time, while that of the tails decreases or stays stable. Filling the pores with water is likely a very long process since it requires monomers to make space for water molecules. Systems started with a higher pore water content (Figures S2c and S2d) tend to plateau relatively quickly, with about one third of the water staying in the tails. We used this ratio in order to construct the initial configurations used for the studies in the main text. Clearly, a more complicated methodology is needed in order to predict the equilibrium water content of a given LLC membrane. However, getting the value exactly right is not required for our study. Instead, we can observe mechanisms as a function of water content in each region.

Since the equilibrium water content is unclear based on the previous simulations, we elected to choose and study systems with two different water contents. We removed the water reservoir and allowed the pore and tail water contents to equilibrate with 5 and 10 wt % total water. We placed one third of the total water

Script Name	Section	Description
/setup/param.sh	TBD	Parameterize liquid crystal monomers and solutes with GAFF
/setup/solvation.equilibration.py	TBD	Add water to the pores and tails in order to achieve a specific total water content and ratio of water molecules in each region, then equilibrate the solvated system.
/setup/input.py	TBD	Create GROMACS topology and .mdp files
/setup/xlink.py	TBD	Iteratively cross-link a configuration
/setup/place_solutes.pores.py	TBD	Place a desired number of solutes in the pore center, equally space in the $z$ -direction.
/analysis/msd.py	TBD	Calculate the mean squared displacement of residues
/analysis/rdf.py	TBD	Calculate the cylindrical radial distribution function of a solute or membrane component with respect to the pore centers.
/analysis/hbonds.py	TBD	Identify hydrogen bonds based on geometric criteria
/timeseries/forecast_ctrw.py	TBD	Construct dwell time and hop length distributions
/analysis/coordination_number.py	TBD	Calculate number of molecules or atoms within a cutoff distance of another type of molecule or atom.
/analysis/ztrace.py	TBD	Plot the center of mass $z$ -coordinate of an atom or molecule versus time which is colored according to its distance from the pore center.

Table S1: The first column provides the names of the python scripts available in the **LLC.Membranes** GitHub repository that were used for system setup and post-simulation trajectory analysis. Paths preceding script names are relative to the **LLC.Membranes/LLC.Membranes** directory. The second columns lists the section in the main text where the output or usage of the script is first described. The third column gives a brief description of the purpose of each script.

Pore Radius (Å)	wt % water tails	wt % water pores
5	5.67	1.09
6	2.88	2.38
7	1.91	4.12
8	2.78	6.00

Table S2: We chose a diverse combination of initial pore and tail water contents in order to study its effect on equilibrium water content.

needed in the tails, based on the intuition gained in Figure S2. We considered the water content equilibrated once the water contents plateaued. The 5 wt% system did not plateau until  $\sim 600$  ns (Figure S3a) while the 10 wt % water system equilibrated within the first 100 ns of simulation (Figure S3b). The pores contain 72 % and 69 % of the total water in the 5 and 10 wt % systems respectively.

We cross-linked the equilibrated solvated systems, then allowed them to equilibrate further for 100 ns. The water contents in each region does not change significantly in either case.




Figure S1: (a) Using an equilibrated dry configuration, we inserted a layer of water between periodic copies of the system in the  $z$ -direction. (b) Water slowly enters the membrane. Most water enters the tail region where the density of monomers is lowest. Water entering the pore plateaus after 500 ns. (c) Although the water content of the pore appears equilibrated in (b), the density of water throughout the pores is not uniform, with almost no water close to the pore center. Note that the density is not shown below 3 nm or above 9.5 nm, because the the system enters the water layer at those points.

## S2.1 Solute Interaction

We chose to model 6 solutes in each pore because there was a low degree of interaction between solutes which gave us a sufficient number of independent trajectories to observe and analyze. There are a negligible number of occurrences where the center of a given solute came within 3.5 Å of another same-solute center of mass.




Figure S2: There is likely more water in the pore region than in the tail region. When we create configurations with more water in the tails, equilibration is slow. When configurations start with more water in the pores, the water content in each region equilibrates quickly.




Figure S3: We created solvated systems with one third of the total water initially placed in the tail region. (a) With 5 wt % total water, the water content equilibrates after 600 ns, with  $\sim 72 \%$  of the total water in the pores. (b) With 10 wt % total water, the water content equilibrates after 100 ns, with  $\sim 69 \%$  of the total water in the pores.




Figure S4: The water content in the tail and pore region is not affected by cross-linking




Figure S5: Interactions between same-solutes are negligible when systems are built with 6 solutes per pore. Even methanol, which interacts with other same-solute molecules the most, does so less than 0.05 % of the time.

## References

- [1] Y. Meroz and I. M. Sokolov, “A Toolbox for Determining Subdiffusive Mechanisms,” *Physics Reports*, vol. 573, pp. 1–29, Apr. 2015.