

Inference in Bayes Nets

- ▶ What is $P(G|W)$? (i.e., the four probability values $P(g|w)$, $P(-g|w)$, $P(g|-w)$, and $P(-g|-w)$).
- ▶ Query variable is G .
- ▶ First run of VE, evidence is $W = w$.
- ▶ Second run of VE, evidence is $W = -w$.
- ▶ Use same ordering for both runs of VE: E, B, S, G .
- ▶ With same ordering some factors can be reused between the two runs of VE.

Inference in Bayes Nets

- ▶ What is $P(G|W)$? (i.e., the four probability values $P(g|w)$, $P(-g|w)$, $P(g|-w)$, and $P(-g|-w)$).
 1. E : $P(E)$, $P(S|E, B)$
 2. B : $P(B)$,
 3. S : $P(w|S)$, $P(S|G)$
 4. G :

Inference in Bayes Nets

- What is $P(G|W)$? (i.e., the four probability values $P(g|w)$, $P(-g|w)$, $P(g|-w)$, and $P(-g|-w)$).

1. E : $P(E)$, $P(S|E, B)$

2. B : $P(B)$,

3. S : $P(w|S)$, $P(S|G)$

4. G :

$$\begin{aligned} F_1(S, B) &= \sum_E P(E) \times P(S|E, B) \\ &= P(e) \times P(S|e, B) + P(-e) \times P(S|-e, B) \end{aligned}$$

$$\begin{aligned} F_1(-s, -b) &= P(e)P(-s|e, -b) + P(-e)P(-s|-e, -b) \\ &= 0.1 \times 0.8 + 0.9 \times 1 = 0.98 \end{aligned}$$

$$\begin{aligned} F_1(-s, b) &= P(e)P(-s|e, b) + P(-e)P(-s|-e, b) \\ &= 0.1 \times 0.1 + 0.9 \times 0.2 = 0.19 \end{aligned}$$

$$\begin{aligned} F_1(s, -b) &= P(e)P(s|e, -b) + P(-e)P(s|-e, -b) \\ &= 0.1 \times 0.2 + 0.9 \times 0 = 0.02 \end{aligned}$$

$$\begin{aligned} F_1(s, b) &= P(e)P(s|e, b) + P(-e)P(s|-e, b) \\ &= 0.1 \times 0.9 + 0.9 \times 0.8 = 0.81 \end{aligned}$$

Inference in Bayes Nets

1. E : $P(E), P(S|E, B)$
2. B : $P(B), F_1(S, B)$
3. S : $P(w|S), P(S|G)$
4. G :

$$\begin{aligned} F_2(S) &= \sum_B P(B) \times F_1(S, B) \\ &= P(b)F_1(S, b) + P(-b)F_1(S, -b) \end{aligned}$$

$$\begin{aligned} F_2(-s) &= P(b)F_1(-s, b) + P(-b)F_1(-s, -b) \\ &= 0.1 \times 0.19 + 0.9 \times 0.98 = \mathbf{0.901} \end{aligned}$$

$$\begin{aligned} F_2(s) &= P(b)F_1(s, b) + P(-b)F_1(s, -b) \\ &= 0.1 \times 0.81 + 0.9 \times 0.02 = \mathbf{0.099} \end{aligned}$$

Inference in Bayes Nets

1. E : $P(E), P(S|E, B)$
2. B : $P(B), F_1(S, B)$
3. S : $P(w|S), P(S|G), F_2(S)$
4. G :

$$\begin{aligned} F_3(G) &= \sum_S P(w|S) \times P(S|G) \times F_2(S) \\ &= P(w|s)P(s|G)F_2(s) + P(w|-s)P(-s|G)F_2(-s) \end{aligned}$$

$$\begin{aligned} F_3(-g) &= P(w|s)P(s|-g)F_2(s) + P(w|-s)P(-s|-g)F_2(-s) \\ &= 0.8 \times 0.5 \times 0.099 + 0.2 \times 1 \times 0.901 = 0.2198 \end{aligned}$$

$$\begin{aligned} F_3(g) &= P(w|s)P(s|g)F_2(s) + P(w|-s)P(-s|g)F_2(-s) \\ &= 0.8 \times 0.5 \times 0.099 + 0.2 \times 0 \times 0.901 = 0.0396 \end{aligned}$$

Inference in Bayes Nets

1. E : $P(E), P(S|E, B)$
2. B : $P(B), F_1(S, B)$
3. S : $P(w|S), P(S|G), F_2(S)$
4. G : $F_3(G)$

Normalize $F_3(G)$:

$$P(-g|w) = \frac{0.2198}{0.2198+0.0396} = 0.8473$$

$$P(g|w) = \frac{0.0396}{0.2198+0.0396} = 0.1527$$

Inference in Bayes Nets

- ▶ Now $P(G \mid -w)$?
 1. E : $P(E), P(S|E, B)$
 2. B : $P(B),$
 3. S : $P(-w|S), P(S|G)$
 4. G :

Already computed as $F_1(S, B)$

Inference in Bayes Nets

1. E : $P(E), P(S|E, B)$
2. B : $P(B), F_1(S, B)$
3. S : $P(-w|S), P(S|G)$
4. G :

Already computed as $F_2(S)$

Inference in Bayes Nets

1. E : $P(E), P(S|E, B)$
2. B : $P(B), F_1(S, B)$
3. S : $P(-w|S), P(S|G), F_2(S)$
4. G :

$$\begin{aligned} F_3(G) &= \sum_S P(-w|S) \times P(S|G) \times F_2(S) \\ &= P(-w|s)P(s|G)F_2(s) + P(-w|-s)P(-s|G)F_2(-s) \end{aligned}$$

$$\begin{aligned} F_3(-g) &= P(-w|s)P(s|-g)F_2(s) + P(-w|-s)P(-s|-g)F_2(-s) \\ &= 0.2 \times 0.5 \times 0.099 + 0.8 \times 1 \times 0.901 = \mathbf{0.7307} \end{aligned}$$

$$\begin{aligned} F_3(g) &= P(-w|s)P(s|g)F_2(s) + P(-w|-s)P(-s|g)F_2(-s) \\ &= 0.2 \times 0.5 \times 0.099 + 0.8 \times 0 \times 0.901 = \mathbf{0.0099} \end{aligned}$$

Inference in Bayes Nets

1. E : $P(E), P(S|E, B)$
2. B : $P(B), F_1(S, B)$
3. S : $P(-w|S), P(S|G), F_2(S)$
4. G : $F_3(G)$

Normalize $F_3(G)$:

$$P(-g|w) = \frac{0.7307}{0.7307 + 0.0099} = 0.9866$$

$$P(g|w) = \frac{0.0099}{0.7307 + 0.0099} = 0.0134$$

Inference in Bayes Nets

- ▶ What do these values tell us about the relationship between G and W , and why does this relationship differ when we know S ?

Inference in Bayes Nets

- ▶ What do these values tell us about the relationship between G and W , and why does this relationship differ when we know S ?

G and W are not independent of each other. But when S is known they become independent.