

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB using the details at the end of this handout.

What GHK-Cu is

GHK-Cu is a copper-binding peptide that occurs naturally in the human body. It is found in blood plasma, saliva and urine, and its levels decline with age. It is being studied for possible support of tissue health, collagen support, wound healing and anti-inflammatory processes.

GHK-Cu contains copper as part of its structure. Copper, zinc and ceruloplasmin are part of your standard lab panel, and your provider will pay particular attention to these results while you are on GHK-Cu.

What it may support

This medication is one possible adjunctive support option for patients with goals related to:

- Skin quality, collagen support and tissue healing
- Anti-inflammatory support in the context of a structured program
- Wound healing and cellular tissue wellness
- Recovery physiology as an adjunctive support option

Response to GHK-Cu varies from patient to patient. It is not a cosmetic treatment or a guaranteed skin improvement product. Your provider will assess whether therapy is appropriate to continue at your follow-up visit.

How to use it

Your prescription label, or the instructions given to you by your KORB clinical provider, is the authoritative source for your dose and directions. Some pharmacies label vials "as directed by provider" rather than printing exact instructions. What follows is a general reference. Do not adjust anything without speaking to your provider first.

How to inject	Subcutaneous (SQ) injection — fatty tissue under the skin
When to inject	Evening • Choose consistent days (e.g. Mon / Wed / Fri)
Schedule	3 times per week (evenings)

Gateway / Peak add-on — active weeks	Weeks 5–8
--------------------------------------	-----------

GHK-Cu is an optional add-on. It begins two weeks after you start BPC-157 and runs for four weeks.

Choosing your three days

Choose three consistent days each week, for example Monday, Wednesday and Friday, and keep the same days each week to build a reliable routine.

Copper, zinc and ceruloplasmin are watched closely

These are part of your standard lab panel. While you are on GHK-Cu your provider reviews them closely at baseline and at each follow-up visit, and therapy may be discontinued if copper or ceruloplasmin levels trend upward.

If you miss a dose

If you miss a dose on a scheduled day, skip it. Do not double up the next day.

Storage and handling

What to do	Detail
Refrigerate	36°F – 46°F (2°C – 8°C)
Do not freeze	Freezing damages medication
Protect from light	Keep away from direct sunlight
Discard at 28 days	Write the open date on the vial

Before first use: refrigerate, protect from light, do not freeze or shake.

After opening: write the open date on your vial and discard 28 days after first use, even if medication remains. If your label shows an earlier beyond-use date, follow the earlier one.

Do not use if the solution appears cloudy or discolored, or contains particles.

Traveling with your medication

Traveling does not have to interrupt your program. Keep your medication with you, in your carry-on or personal item rather than checked baggage, in its original labeled vial or packaging. It does not need to be packed in ice during travel; keep it away from excessive heat (above 86°F / 30°C) and from freezing, and refrigerate it again when you arrive. Medication and syringes are permitted in carry-on luggage, and TSA recommends keeping the original pharmacy label visible. Crossing time zones can shift your injection time by a few hours without a problem; if you will be off your usual schedule for several days, contact KORB.

What to expect

Timeline	Phase	What to expect
First 2 weeks	Active course begins	Most patients notice nothing initially. This is normal; the therapy works gradually.
Weeks 3–4	Active course continues	Skin quality or tissue changes may begin to emerge. Individual response varies.
After your course	Washout — labs and lifestyle	No GHK-Cu injections. Your copper, zinc and ceruloplasmin results will be reviewed at your 16-week follow-up visit, and this is a good window to focus on nutrition, exercise and sleep before your next course.

Your provider will review copper, zinc and ceruloplasmin closely at baseline and at your follow-up visit while GHK-Cu is active. Therapy may be discontinued if copper or ceruloplasmin levels rise.

Side effects and what to watch for

What you may notice

What you may notice	What to do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell your KORB provider if it persists.
Mild skin changes at the injection site	Occasional mild discoloration or sensitivity. Usually self-limiting.
Fatigue	Uncommon. Tell your KORB provider if it persists.

Tell your KORB provider at your next visit

What you may notice	What to do
Copper or ceruloplasmin trending up on labs	Your provider will review this at your follow-up visit. Therapy may be discontinued if levels rise.
Unusual rash, hives or skin reaction	Tell your KORB provider promptly. Copper-containing compounds can occasionally cause hypersensitivity.
Any systemic symptom	Tell your KORB provider. Do not continue without guidance if you have a new unexplained systemic symptom.

Stop injecting and seek emergency care immediately if you have trouble breathing, swelling of the face, lips, tongue or throat, a severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab monitoring

Lab monitoring is a required part of your program, not optional. Your provider uses labs to assess your response and your safety.

- Standard longevity panel: CBC, CMP, lipid panel, HbA1c, fasting glucose, fasting insulin, IGF-1, TSH, free T4, free T3, copper, zinc and ceruloplasmin, at baseline and every 16-week follow-up.
- PSA, added for men aged 45 and older at every draw.
- Copper, zinc and ceruloplasmin get particular attention while you are on GHK-Cu, and therapy may be discontinued if copper or ceruloplasmin trend upward.

Your lab draw should be completed between weeks 12 and 14 of your active cycle, so results are back in time for your 16-week follow-up visit. Some results, such as IGF-1, can take about a week. Unless your lab order says otherwise, plan to fast overnight and have blood drawn first thing in the morning.

Safety reminders

- Use GHK-Cu only as prescribed. Do not change your dose, schedule or course length without provider direction.
- GHK-Cu is absolutely contraindicated if you have Wilson's disease, a copper metabolism disorder.
- Copper, zinc and ceruloplasmin are part of your standard labs. Do not skip your scheduled draw between weeks 12 and 14.
- Tell your KORB provider if you develop a new medical condition or become pregnant.
- Use a new insulin needle and syringe for every injection. Do not reuse.
- Never mix peptides. Use a separate needle and syringe for each peptide and each injection; do not combine them into one syringe.
- Follow your state or local rules for sharps disposal. Use an FDA-cleared sharps container, and do not put used needles or syringes in household trash unless your local guidance specifically allows it.

When to contact KORB

Contact KORB Operations	Patient Portal — message your provider	Seek emergency care immediately
- Questions about timing, storage or administration - Scheduling, billing and shipping Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	- Injection-site irritation or redness that does not resolve - Mild but persistent side effects - Questions about whether to continue therapy	- Trouble breathing, or swelling of the face, lips, tongue or throat - Severe rash or widespread hives - Chest pain or fainting - Severe or rapidly worsening symptoms - Confusion or severe weakness - Any symptom that feels urgent or unsafe

The Patient Portal is the only HIPAA-compliant way to message your provider directly. Providers check portal messages once per day on their clinic days. For anything that does not need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key reminders

- GHK-Cu is taken three times a week in the evening, on consistent days.
- It is absolutely contraindicated in Wilson's disease.
- Copper, zinc and ceruloplasmin are watched closely and may end therapy if they rise.
- Write the open date on your vial and discard 28 days after first use.
- Contact KORB Operations for non-urgent questions, your provider through the Patient Portal for medical questions, and emergency care for severe symptoms.