

Injection, Storage & Safety Guide

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This guide is for general educational reference only. It does not replace your prescription label, pharmacy-specific instructions, or your KORB provider's guidance. If your pharmacy or provider instructions differ from anything in this guide, always follow your pharmacy and provider instructions.

This guide applies to your injectable Functional Health & Longevity Program therapy — Sermorelin, BPC-157, CJC-1295/Ipamorelin, GHK-Cu, or Tesamorelin — and any other injectable therapy your KORB provider prescribes as part of the program.

What This Guide Covers

This guide supports safe medication handling, storage, injection preparation, site rotation, sharps disposal, and general safety awareness for injectable Longevity Program therapies.

It does not replace your prescription label, pharmacy instructions, your KORB provider's guidance, medication-specific patient education materials, or emergency medical care.

Important Safety Reminder

Injectable therapies must be used exactly as prescribed. Unless your KORB provider specifically directs otherwise:

- Do not change your dose, frequency, or injection timing
- Do not change your injection route or injection location
- Do not change your treatment schedule or combine it with other therapies

Before You Inject

Before each injection, make sure you have:

- Your medication
- A new insulin needle and syringe
- Alcohol swabs
- A sharps container
- A clean surface and good lighting
- Your prescription or pharmacy instructions

Wash your hands with soap and water before preparing your injection. If you're unsure about the dose, medication, route, or instructions, contact KORB before proceeding.

Check Your Medication First

Don't use your medication if:

- It looks cloudy when it should be clear, or has particles or a color change
- The vial appears cracked or damaged
- It has expired or you're unsure it was stored correctly
- You're unsure whether it's safe to use for any reason

If you have any concern about the medication's appearance, storage, or safety, contact the pharmacy or KORB before using it.

Storage Basics

Refrigerate 36°F – 46°F (2°C – 8°C)	Do not freeze Freezing damages medication	Protect from light Keep away from direct sunlight	Discard at 28 days Write open date on vial when opened
---	---	---	--

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

Always follow the storage instructions on your pharmacy label if they differ from this general guidance. Keep all medication out of reach of children and pets.

Injection Route & Technique

Longevity Program injectables are typically given subcutaneously — into the fatty tissue under the skin. Never inject into a vein, muscle, joint, tendon, ligament, wound, rash, infected area, or broken or irritated skin unless your provider has specifically instructed otherwise.

Common injection sites: abdomen (avoiding the belly button), outer thigh, back of the upper arm (if someone else is injecting you), or upper buttock/hip if instructed. Use only the sites your provider or pharmacy recommends, and avoid any area that's red, swollen, hot, painful, bruised, scarred, hard, lumpy, or recently used.

Rotate sites every time to reduce irritation and tissue changes — stay at least 1–2 inches from your last injection site and keep track if you're injecting frequently.

- Wash your hands and clean the injection site with an alcohol swab; let it dry before injecting
- Prepare the correct dose exactly as instructed
- Dispose of the needle and syringe immediately in a sharps container
- Use a new insulin needle with syringe for every injection — do not reuse.
- Never mix peptides — use one insulin needle with syringe for every peptide, every injection. Do not combine into one injection or syringe.

Sharps Disposal

Sharps disposal is your responsibility, and rules vary by state and even by city — check your local regulations.

- Use an FDA-cleared sharps container, or a heavy-duty plastic container with a tight-fitting lid if one isn't available
- Ask your pharmacy about sharps container options or local drop-off/mail-back programs
- For state-specific disposal rules and drop-off locations, visit safeneedledisposal.org or call 1-800-643-1643 (Coalition for Safe Community Needle Disposal, the resource recommended by the FDA)

Missed a Dose?

- If you miss a dose on a scheduled day: skip it — do not double up the next day

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep your medication with you — in your carry-on or personal item, not checked baggage — in its original labeled vial or packaging. It doesn't need to be packed in ice or kept refrigerated during travel; just keep it away from excessive heat (above 86°F / 30°C) and from freezing. Once you reach your destination, refrigerate it again. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Common Mild Injection-Site Reactions

Mild redness, soreness, swelling, bruising, itching, or temporary tenderness at the injection site are common and usually improve on their own. Contact KORB if a reaction is worsening, spreading, persistent, painful, warm, or draining.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations

- Storage questions
- Sharps disposal questions
- General medication questions

Email: info@korbhealth.com

Phone: +1 (888) 959-7299

Hours: Mon–Fri, 9 AM–6 PM CT

Patient Portal — Message Your Provider

- Anything requiring your provider's direct medical judgment
- Worsening or persistent injection-site reactions
- Questions about whether to continue therapy

Seek Emergency Care Immediately

- Trouble breathing or swelling of face, lips, tongue, or throat
- Severe rash or widespread hives
- Chest pain or fainting
- Severe weakness or confusion
- New neurologic symptoms
- Signs of serious infection

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Seek urgent or emergency care right away for: trouble breathing; swelling of the face, lips, tongue, or throat; severe rash or widespread hives; chest pain or fainting; severe weakness or confusion; severe or rapidly worsening pain; new neurologic symptoms (weakness, numbness, confusion, loss of bladder/bowel control); or signs of serious infection (fever, spreading redness, severe warmth, drainage, or red streaking from an injection site). Phone and email are not appropriate for emergencies.

Key Reminders

Use injectable therapies only as prescribed — never change dose, frequency, route, or schedule without provider direction.

Never mix peptides — use one insulin needle with syringe for every peptide, every injection. Do not combine into one injection or syringe.

Check your medication before every injection — don't use it if it looks cloudy, discolored, has particles, or the vial is damaged.

Rotate injection sites and never inject into irritated, infected, broken, or abnormal skin.

Use a new insulin needle with syringe every time. Never recap, reuse, or share needles.

Sharps disposal is your responsibility, and rules vary by location — follow your state or local regulations.

Use the Patient Portal only for provider-directed messages. Contact KORB Operations by phone or email for everything else.

Seek emergency care immediately for severe or unsafe symptoms.