

Injection, Storage & Safety Guide

Patient Education Handout

KORB Health Group · For Patient Reference

This guide is for general educational reference only. It does not replace your prescription label, pharmacy-specific instructions, or your KORB provider's guidance. If your pharmacy or provider instructions differ from anything in this guide, always follow your pharmacy and provider instructions.

This guide applies to any injectable medication your KORB provider prescribes — including GLP-1 medications (Semaglutide, Tirzepatide), testosterone therapy, and Functional Health & Longevity peptides. If your program uses a patch, cream, or oral medication instead of an injection, ask your provider or pharmacy for storage and use instructions specific to that formulation.

What This Guide Covers

This guide supports safe medication handling, storage, injection preparation, site rotation, sharps disposal, and general safety awareness for injectable KORB therapies. It does not replace your prescription label, pharmacy instructions, your KORB provider's guidance, medication-specific patient education materials, or emergency medical care.

Important Safety Reminder

Injectable therapies must be used exactly as prescribed. Unless your KORB provider specifically directs otherwise:

- Do not change your dose, frequency, or injection timing
- Do not change your injection route or injection location
- Do not change your treatment schedule or combine it with other therapies

Before You Inject

Before each injection, make sure you have:

- Your medication
- Alcohol swabs
- A sharps container
- A clean surface and good lighting
- Your pharmacy label or the written instructions that came with your medication

Your syringe and needles depend on which medication you are using.

Medication	What you use
GLP-1 and peptides	A new insulin syringe with the needle already attached, one per injection
Testosterone	A Luer lock syringe with two separate needles: a wider one to draw the oil out of the vial, and a finer one to inject with

KORB does not supply sharps containers. Obtaining one and disposing of it correctly is your responsibility, and the rules are set by your state and sometimes your city. Look up what applies where you live at safeneedledisposal.org before your first injection.

Wash your hands with soap and water before preparing your injection. If you're unsure about the dose, medication, route, or instructions, contact KORB before proceeding.

Check Your Medication First

Don't use your medication if:

- It looks cloudy when it should be clear, or has particles or a color change
- The vial appears cracked or damaged
- It has expired or you're unsure it was stored correctly
- You're unsure whether it's safe to use for any reason

If you have any concern about the medication's appearance, storage, or safety, contact the pharmacy or KORB before using it.

Storage Basics

Refrigerate	Do not freeze	Protect from light	Discard at 28 days
36°F – 46°F (2°C – 8°C)	Freezing damages medication	Keep away from direct sunlight	Write open date on vial

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles. Always follow the storage instructions on your pharmacy label if they differ from this general guidance. Keep all medication out of reach of children and pets.

Injection Route & Technique

Most KORB injectable therapies are given subcutaneously — into the fatty tissue under the skin. Never inject into a vein, muscle, joint, tendon, ligament, wound, rash, infected area, or broken or irritated skin unless your provider has specifically instructed otherwise.

Common injection sites: abdomen (avoiding the belly button), outer thigh, back of the upper arm (if someone else is injecting you), or upper buttock/hip if instructed. Use only the sites your provider or

pharmacy recommends, and avoid any area that's red, swollen, hot, painful, bruised, scarred, hard, lumpy, or recently used.

Rotate sites every time to reduce irritation and tissue changes — stay at least 1–2 inches from your last injection site and keep track if you're injecting frequently.

Wash your hands and clean the injection site with an alcohol swab; let it dry before injecting

Prepare the correct dose exactly as instructed

Dispose of the needle and syringe immediately in a sharps container

Use new equipment for every injection. Never reuse a syringe or needle.

If you are using testosterone: draw the medication up with the wider needle, then change to the finer needle before injecting. The oil is thick and will not draw properly through the fine needle, and drawing with the injecting needle dulls it and makes the injection hurt more.

If you are prescribed more than one injectable medication, use a separate needle and syringe for each — never combine different medications into one injection or syringe.

Sharps Disposal

Sharps disposal is your responsibility, and rules vary by state and even by city — check your local regulations.

Use an FDA-cleared sharps container, or a heavy-duty plastic container with a tight-fitting lid if one isn't available

Ask your pharmacy about sharps container options or local drop-off/mail-back programs

For state-specific disposal rules and drop-off locations, visit safeneedledisposal.org or call 1-800-643-1643 (Coalition for Safe Community Needle Disposal, the resource recommended by the FDA)

Missed a Dose?

If you miss a dose on a scheduled day, contact KORB or check your medication-specific guide — timing varies by medication. Don't guess, and don't double up without guidance.

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep your medication with you — in your carry-on or personal item, not checked baggage — in its original labeled vial or packaging. It doesn't need to be packed in ice or kept refrigerated during travel; just keep it away from excessive heat (above 86°F / 30°C) and from freezing. Once you reach your destination, refrigerate it again. If flying, medication and syringes are permitted in carry-on luggage. TSA asks that you tell the officer at the start of screening that you are carrying medically necessary liquids, and they may need to be inspected separately. You can read the current rules yourself at tsa.gov/travel/travel-tips. TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Mild Injection-Site Reactions

Some patients notice mild redness, soreness, swelling, bruising, itching, or temporary tenderness at the injection site. These usually settle on their own. Contact KORB if a reaction is worsening, spreading, persistent, painful, warm, or draining.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations	Patient Portal — Message Your Provider	Seek Emergency Care Immediately
Questions about timing, storage, or administration Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	- Mild but persistent side effects - Questions about whether to continue therapy - Worsening or persistent injection-site reactions	- Trouble breathing or throat / face swelling - Severe rash or widespread hives - Chest pain or fainting - Severe or rapidly worsening symptoms - Confusion or severe weakness - Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on clinic days. Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Seek urgent or emergency care right away for: trouble breathing; swelling of the face, lips, tongue, or throat; severe rash or widespread hives; chest pain or fainting; severe weakness or confusion; severe or rapidly worsening pain; new neurologic symptoms (weakness, numbness, confusion, loss of bladder/bowel control); or signs of serious infection (fever, spreading redness, severe warmth, drainage, or red streaking from an injection site). Phone and email are not appropriate for emergencies.

Key Reminders

Use injectable therapies only as prescribed — never change dose, frequency, route, or schedule without provider direction.

If prescribed more than one injectable medication, use a separate needle and syringe for each — never combine into one injection or syringe.

Check your medication before every injection — don't use it if it looks cloudy, discolored, has particles, or the vial is damaged.

Rotate injection sites and never inject into irritated, infected, broken, or abnormal skin.

Use new equipment every time. Never recap, reuse, or share needles.

Sharps disposal is your responsibility. Look up the rules where you live at safeneedledisposal.org.

For medical questions, message your provider through the [Patient Portal](#). For anything else, call KORB Operations at +1 (888) 959-7299.

Use the Patient Portal only for provider-directed messages. Contact KORB Operations by phone or email for everything else.

Seek emergency care immediately for severe or unsafe symptoms.