

Hair Loss Treatment

Patient Education Handout

Hair Loss Add-On · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label and your KORB provider's guidance. If your label differs from anything here, follow your label and contact KORB.

What You May Be Prescribed

Hair loss treatment comes as a once-daily pill, a once-daily topical you apply to your scalp, or both together. Your provider chooses based on your pattern of loss, your health history, and your preference.

	Pill	Topical
Men	Finasteride	Minoxidil with finasteride and tretinoin
Women	Spironolactone	Minoxidil with spironolactone and latanoprost

These are not interchangeable between people. The men's pill, finasteride, must never be taken by a woman who is or could become pregnant, because it can cause serious birth defects. Never take, share, or borrow someone else's prescription, including a partner's.

Give It Three to Six Months

This is the single most important thing to understand before you start. Hair grows slowly, and these treatments work by protecting the hair you have and improving the quality of new growth. Most people see no visible change for the first three months, and the clearest results appear between three and six months.

Some people notice increased shedding in the first few weeks. That is usually a normal part of the cycle resetting and not a sign the treatment is failing. Stopping early is the most common reason treatment does not work. If you stop, any gains will gradually reverse.

How to Use the Topical

Apply once daily to a dry scalp, not damp or freshly washed hair.

Use one to two applications depending on how much area you are covering. Your provider will tell you which.

Work it into the scalp where you are thinning, not onto the hair itself.

Wash your hands afterward.

Leave it on for at least four hours, or overnight if that suits your routine better.

How long your bottle lasts depends on how much you use. If you apply two doses a day it will run out roughly twice as fast as one. Keep this in mind so you are not caught short, and tell us early if you are running low rather than at the last minute.

Side Effects

What you may notice	What to do
Scalp redness, itching, or dryness (topical)	Common early on. Tell your provider if it is severe or does not settle.
Increased shedding in the first few weeks	Usually normal. Contact KORB if it is heavy or lasts beyond about six weeks.
Reduced sex drive or difficulty with erections (finasteride)	Uncommon but real. Tell your provider rather than stopping on your own.
Breast tenderness or swelling (either pill)	Tell your provider.
Dizziness or feeling lightheaded (spironolactone)	Stand up slowly. Tell your provider if it continues.
Unwanted hair growth on the face where the topical has run	Apply more carefully and wash your hands. Tell your provider if it persists.

Tell your provider before continuing if you become pregnant, are trying to conceive, or are breastfeeding. Both pills require a conversation in those situations, and finasteride must be stopped and kept away from anyone who is pregnant.

Key Reminders

Results take three to six months. Do not judge it early.

Some shedding at the start is normal.

Apply the topical to a dry scalp once daily and leave it on at least four hours.

Never share your prescription. Finasteride is dangerous in pregnancy.

Gains reverse if you stop, so tell your provider before discontinuing.

Tell us early if you are running low.

How to reach us

Operations, non-medical: +1 (888) 959-7299 · info@korbhealth.com · Mon–Fri, 9 AM–6 PM CT

Medical questions: message your provider at portal.kareo.com

Emergency: call 911 or go to the nearest emergency room.