

Foundation

Functional Health & Longevity • Provider Reference

KORB Health Medical Texas PA • korb-dosing-data.js v2.7 • built 2026-09-14

Everything needed to prescribe the Foundation Program, complete on its own. The provider tool covers the same ground; this document exists so it is not required. Values are copied literally into the Tebra Compound section.

At a glance

Program	Foundation Program
Structure	Single agent, provider selects one. No stagger, no optional add-on.
Cycle length	16 weeks
Agents in this document	Sermorelin 200 mcg • Sermorelin 300 mcg • Sermorelin 400 mcg • CJC-1295 / Ipamorelin 100 mcg CJC-1295 / 100 mcg Ipamorelin • CJC-1295 / Ipamorelin 150 mcg CJC-1295 / 150 mcg Ipamorelin • CJC-1295 / Ipamorelin 200 mcg CJC-1295 / 200 mcg Ipamorelin • BPC-157 500 mcg
Pharmacies	Premier Pharmacy and Greenwich Pharmacy — routed by patient state

The 16-week cycle

Agent	Role	Active weeks	Timing
Sermorelin 200 mcg	Standalone course	Weeks 1–12	Bedtime • Empty stomach (2+ hrs after eating)
Sermorelin 300 mcg	Standalone course	Weeks 1–12	Bedtime • Empty stomach (2+ hrs after eating)
Sermorelin 400 mcg	Standalone course	Weeks 1–12	Bedtime • Empty stomach (2+ hrs after eating)
CJC-1295 / Ipamorelin 100 mcg CJC-1295 / 100 mcg Ipamorelin	Standalone course	Weeks 1–12	Bedtime • Empty stomach (2+ hrs after eating)

Agent	Role	Active weeks	Timing
CJC-1295 / Ipamorelin 150 mcg CJC-1295 / 150 mcg Ipamorelin	Standalone course	Weeks 1–12	Bedtime · Empty stomach (2+ hrs after eating)
CJC-1295 / Ipamorelin 200 mcg CJC-1295 / 200 mcg Ipamorelin	Standalone course	Weeks 1–12	Bedtime · Empty stomach (2+ hrs after eating)
BPC-157 500 mcg	Standalone course	Weeks 1–8	Any consistent time · No food restriction

Foundation is a single agent. The provider selects one of the three; there is no stagger and GHK-Cu is never a Foundation option.

Dose ladder — Sermorelin

Escalation is lab-based. Increase to the next step only if IGF-1 remains below target at the 16-week recheck.

Titrate 200 → 300 → 400 mcg at 16-week visits, based on IGF-1. 500 mcg is NOT a Foundation dose - a patient who needs it moves to Gateway rather than being held at 400. See [programs.foundation.sermorelin500ExclusionReason](#).

500 mcg is not a Foundation dose

A correctly supplied 500 mcg is 45 ml a cycle against 36 ml at 400 mcg, which requires an additional vial each month. Foundation margin does not justify it. Gateway retains 500 mcg. This is a commercial ceiling, not a clinical one - escalate the patient to Gateway rather than holding at 400.

Dose	Premier — units	Greenwich — units	Active weeks	Schedule
200 mcg	20 units	7 units	Weeks 1–12	6 days on, 1 day off
300 mcg	30 units	10 units	Weeks 1–12	6 days on, 1 day off
400 mcg	40 units	13 units	Weeks 1–12	6 days on, 1 day off

The milligram dose is what is prescribed. The unit count is what the patient draws, and it differs by pharmacy.

Dose ladder — CJC-1295 / Ipamorelin

Escalation is lab-based. Increase to the next step only if IGF-1 remains below target at the 16-week recheck.

Titrate 100 → 150 → 200 mcg at 16-week visits. All three doses dispense the same quantity, so titration changes no vial and no cost.

Dose	Premier — units	Greenwich — units	Active weeks	Schedule
100 mcg CJC-1295 / 100 mcg Ipamorelin	5 units	5 units	Weeks 1–12	6 days on, 1 day off
150 mcg CJC-1295 / 150 mcg Ipamorelin	7.5 units	7.5 units	Weeks 1–12	6 days on, 1 day off
200 mcg CJC-1295 / 200 mcg Ipamorelin	10 units	10 units	Weeks 1–12	6 days on, 1 day off

The milligram dose is what is prescribed. The unit count is what the patient draws, and it differs by pharmacy.

Directions given to the patient

Agent	Pharmacy	Direction
Sermorelin 200 mcg	Premier Pharmacy	Inject 20 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
Sermorelin 200 mcg	Greenwich Pharmacy	Inject 7 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
Sermorelin 300 mcg	Premier Pharmacy	Inject 30 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
Sermorelin 300 mcg	Greenwich Pharmacy	Inject 10 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
Sermorelin 400 mcg	Premier Pharmacy	Inject 40 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
Sermorelin 400 mcg	Greenwich Pharmacy	Inject 13 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
CJC-1295 / Ipamorelin 100 mcg CJC-1295 / 100 mcg Ipamorelin	Premier Pharmacy	Inject 5 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
CJC-1295 / Ipamorelin 100 mcg CJC-1295 / 100 mcg Ipamorelin	Greenwich Pharmacy	Inject 5 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
CJC-1295 / Ipamorelin 150 mcg CJC-1295 / 150 mcg Ipamorelin	Premier Pharmacy	Inject 7.5 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.

Agent	Pharmacy	Direction
CJC-1295 / Ipamorelin 150 mcg CJC-1295 / 150 mcg Ipamorelin	Greenwich Pharmacy	Inject 7.5 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
CJC-1295 / Ipamorelin 200 mcg CJC-1295 / 200 mcg Ipamorelin	Premier Pharmacy	Inject 10 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
CJC-1295 / Ipamorelin 200 mcg CJC-1295 / 200 mcg Ipamorelin	Greenwich Pharmacy	Inject 10 units subcutaneously at bedtime, 6 days on, 1 day off, on an empty stomach.
BPC-157 500 mcg	Premier Pharmacy	Inject 13 units subcutaneously once daily.
BPC-157 500 mcg	Greenwich Pharmacy	Inject 17 units subcutaneously once daily.

The rest day is not a fixed weekday

The protocol is 6 days on, 1 day off. The specific weekday is a recommendation for ease of patient recall, not a clinical requirement. The patient's tracker derives their rest day from their own start date, so a patient starting on a Wednesday rests Tuesday. Do not tell a patient their rest day is Sunday.

Syringes

Pharmacy	Standard syringe	When a 100-unit syringe is required instead
Premier Pharmacy	100-unit (1 mL)	Not applicable — Premier ships 100-unit syringes across the board.
Greenwich Pharmacy	50-unit (0.5 mL)	Any dose at or above 50 units. Currently Tesamorelin 1.5 mg, dosed at 50 units; Tesamorelin 2 mg, dosed at 67 units.

Which pharmacy, and where you can prescribe

Defaults to Premier	AZ, CO, CT, DE, FL, IL, KS, KY, LA, MD, ME, MI, MO, MT, NC, ND, NE, NM, NV, OH, OK, OR, PA, SD, TN, TX, UT, VA, VT, WI, WY
Everywhere else	Greenwich Pharmacy, for every state not listed as closed
Functional Health & Longevity is not offered in these states	AL, AK, AR, CA, DC, GA, HI, IN, MA, MN, MS, NH, NJ, NY, RI, SC, WA, WV

Paused — no pharmacy can fill — AR, CA, IN, NH, WA

Greenwich stopped relying on central-fill arrangements on 11 September 2026 and now dispenses only into the 23 states where it is directly licensed. Premier is not licensed in any of these, so there is no peptide source while the restriction holds. No visit, no prescription, no shipment.

This is a pause, not a withdrawal. Greenwich expects to restore coverage in 30 to 60 days from 11 September 2026. Tell a patient the program is temporarily unavailable in their state and that coverage is expected back. Do not tell them it has been discontinued. Operations reviews this by 10 November 2026.

AL, MN, SC also lost Greenwich, but each was already out of the peptide program before this change and stays closed regardless of the restore. They are not listed here.

Single-source states — IA, ID

These states are open and taking patients. Greenwich is the only pharmacy licensed to fill a peptide there and Premier cannot cover them, so there is no fallback if Greenwich restricts again. Greenwich is mid-restriction, so confirm the fill before promising a patient a start date.

Excluded from the offering — AL, MS, SC

No peptide or other Functional Health & Longevity product may be shipped to these states. They are excluded from the offering, not pending launch. Confirm with Operations before any exception.

Held for lab workflow — NY, NJ, RI

These are held for a distinct reason: a different lab workflow required by state-specific legislation, not provider coverage. Do not proceed until Operations confirms.

Unit counts differ between the pharmacies for the same milligram dose. Always read the units off the row for the patient's pharmacy, never from the other column.

Tebra Compounded Drug Favorite — exact fields

Copy these values literally. Do not paraphrase, and do not adjust quantity, refill or days supply. Greenwich requires “as directed by provider” in Patient Instructions because KORB does not follow their protocols — the full direction still goes to the patient through their treatment schedule.

Sermorelin — 200 mcg

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – Sermorelin 200 mcg SQ Injection

Drug Formulation	Sermorelin 1 mg/ml inj
Name	PREMIER – Sermorelin 200 mcg SQ Injection
Allow Substitution	Yes — select Allow Substitution
Quantity	18
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	INJECT SERMORELIN 20 UNITS SUBCUTANEOUSLY AT BEDTIME, 6 DAYS ON 1 DAY OFF, ON AN EMPTY STOMACH
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 80 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – Sermorelin 0.6 mg SQ Injection

Drug Formulation	KBH Sermorelin 3mg/mL
Name	GREENWICH – Sermorelin 0.6 mg SQ Injection
Allow Substitution	Yes — select Allow Substitution
Quantity	15
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

Sermorelin 300mcg — 300 mcg

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – Sermorelin 300 mcg SQ Injection	
Drug Formulation	Sermorelin 1 mg/ml inj
Name	PREMIER – Sermorelin 300 mcg SQ Injection
Allow Substitution	Yes — select Allow Substitution
Quantity	27
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	INJECT SERMORELIN 30 UNITS SUBCUTANEOUSLY AT BEDTIME, 6 DAYS ON 1 DAY OFF, ON AN EMPTY STOMACH
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 80 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – Sermorelin 0.9 mg SQ Injection	
Drug Formulation	KBH Sermorelin 3mg/mL
Name	GREENWICH – Sermorelin 0.9 mg SQ Injection
Allow Substitution	Yes — select Allow Substitution
Quantity	15
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

Sermorelin 400mcg — 400 mcg

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – Sermorelin 400 mcg SQ Injection

Drug Formulation	Sermorelin 1 mg/ml inj
Name	PREMIER – Sermorelin 400 mcg SQ Injection
Allow Substitution	Yes — select Allow Substitution
Quantity	36
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	INJECT SERMORELIN 40 UNITS SUBCUTANEOUSLY AT BEDTIME, 6 DAYS ON 1 DAY OFF, ON AN EMPTY STOMACH
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 80 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – Sermorelin 1.2 mg SQ Injection

Drug Formulation	KBH Sermorelin 3mg/mL
Name	GREENWICH – Sermorelin 1.2 mg SQ Injection
Allow Substitution	Yes — select Allow Substitution
Quantity	15
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

CJC-1295 / Ipamorelin 100mcg/100mcg — 100 mcg CJC-1295 / 100 mcg Ipamorelin

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – CJC/Ipam 100 mcg/100 mcg SQ Inj	
Drug Formulation	CJC-1295 / Ipamorelin 2 mg/2 mg per 1mL inj
Name	PREMIER – CJC/Ipam 100 mcg/100 mcg SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	9
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	INJECT 5 UNITS SUBCUTANEOUSLY AT BEDTIME, 6 DAYS ON 1 DAY OFF, ON AN EMPTY STOMACH
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 80 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – CJC/Ipam 100 mcg/100 mcg SQ Inj	
Drug Formulation	KBH CJC/Ipamorelin 2mg/2mg/mL
Name	GREENWICH – CJC/Ipam 100 mcg/100 mcg SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	15
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

CJC-1295 / Ipamorelin 150mcg/150mcg — 150 mcg CJC-1295 / 150 mcg Ipamorelin

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – CJC/Ipam 150 mcg/150 mcg SQ Inj

Drug Formulation	CJC-1295 / Ipamorelin 2 mg/2 mg per 1mL inj
Name	PREMIER – CJC/Ipam 150 mcg/150 mcg SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	9
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	INJECT 7.5 UNITS SUBCUTANEOUSLY AT BEDTIME, 6 DAYS ON 1 DAY OFF, ON AN EMPTY STOMACH
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 80 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – CJC/Ipam 150 mcg/150 mcg SQ Inj

Drug Formulation	KBH CJC/Ipamorelin 2mg/2mg/mL
Name	GREENWICH – CJC/Ipam 150 mcg/150 mcg SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	15
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

CJC-1295 / Ipamorelin 200mcg/200mcg — 200 mcg CJC-1295 / 200 mcg Ipamorelin

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – CJC/Ipam 200 mcg/200 mcg SQ Inj	
Drug Formulation	CJC-1295 / Ipamorelin 2 mg/2 mg per 1mL inj
Name	PREMIER – CJC/Ipam 200 mcg/200 mcg SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	9
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	INJECT 10 UNITS SUBCUTANEOUSLY AT BEDTIME, 6 DAYS ON 1 DAY OFF, ON AN EMPTY STOMACH
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 80 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – CJC/Ipam 200 mcg/200 mcg SQ Inj	
Drug Formulation	KBH CJC/Ipamorelin 2mg/2mg/mL
Name	GREENWICH – CJC/Ipam 200 mcg/200 mcg SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	15
Unit	ml
Refill	0
Days Supply	84
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

BPC-157 — 500 mcg

Premier Pharmacy • Compounded Drug Favorite Entry • PREMIER – BPC-157 500 mcg SQ Inj Daily	
Drug Formulation	BPC-157, 4mg/ml inj
Name	PREMIER – BPC-157 500mcg SQ Inj Daily
Allow Substitution	Yes — select Allow Substitution
Quantity	12
Unit	ml
Refill	0
Days Supply	56
Patient Instructions	INJECT 13 UNITS SUBCUTANEOUSLY ONCE DAILY
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing. Dispense 60 insulin syringes.

Greenwich Pharmacy • Compounded Drug Favorite Entry • GREENWICH – BPC-157 0.6 MG SQ Inj	
Drug Formulation	KBH BPC-157 3mg/mL
Name	GREENWICH – BPC-157 0.6 MG SQ Inj
Allow Substitution	Yes — select Allow Substitution
Quantity	10
Unit	ml
Refill	0
Days Supply	56
Patient Instructions	Inject as directed by provider. Follow your KORB treatment schedule.
Reason for Compounding	Customized peptide dosing
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Customized peptide dosing

Storage: Refrigerate (36–46 °F). Do not freeze. Protect from light. Discard 28 days after initial puncture.

Counseling and monitoring

Sermorelin — counseling points

- Not GH replacement — stimulates body's own GH signaling pathway
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

Sermorelin — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention
- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions
- Adherence to 6-on/1-off schedule

CJC-1295 / Ipamorelin — counseling points

- Dual-peptide GH secretagogue combo — not GH replacement
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

Sermorelin — counseling points

- Not GH replacement — stimulates body's own GH signaling pathway
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval

- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

Sermorelin — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention
- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions
- Adherence to 6-on/1-off schedule

Sermorelin 300mcg — counseling points

- Not GH replacement — stimulates body's own GH signaling pathway
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

Sermorelin 300mcg — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention
- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions

- Adherence to 6-on/1-off schedule

Sermorelin 400mcg — counseling points

- Not GH replacement — stimulates body's own GH signaling pathway
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

Sermorelin 400mcg — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention
- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions
- Adherence to 6-on/1-off schedule

CJC-1295 / Ipamorelin 100mcg/100mcg — counseling points

- Dual-peptide GH secretagogue combo — not GH replacement
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

CJC-1295 / Ipamorelin 100mcg/100mcg — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention

- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions
- Adherence to 6-on/1-off schedule

CJC-1295 / Ipamorelin 150mcg/150mcg — counseling points

- Dual-peptide GH secretagogue combo — not GH replacement
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

CJC-1295 / Ipamorelin 150mcg/150mcg — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention
- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions
- Adherence to 6-on/1-off schedule

CJC-1295 / Ipamorelin 200mcg/200mcg — counseling points

- Dual-peptide GH secretagogue combo — not GH replacement
- Lifestyle (nutrition, resistance training, sleep) remains critical — not a substitute
- Response varies between patients; benefit not guaranteed
- Do not combine with other GH-axis therapies without provider approval
- 6 days ON / 1 day OFF pattern must be followed — your rest day is shown in your KORB treatment schedule
- Take at bedtime on an empty stomach — timing is clinically relevant
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

CJC-1295 / Ipamorelin 200mcg/200mcg — monitoring

- Sleep quality and recovery quality
- Energy and wellness trend
- Body composition / waist circumference trend
- Exercise tolerance and lean mass preservation
- Edema / fluid retention
- Headache
- Joint discomfort
- Paresthesias / tingling
- Glucose-related changes (if at-risk)
- IGF-1 (per lab protocol)
- Injection site reactions
- Adherence to 6-on/1-off schedule

BPC-157 — counseling points

- Investigational recovery-support adjunct — not a cure or injury reversal
- Does not replace appropriate medical evaluation, imaging, or physical therapy when indicated
- Course-based cycling — do not continue indefinitely without provider reassessment
- Report injection site reactions, rash, swelling, or difficulty breathing immediately
- Lifestyle foundations (sleep, nutrition, structured rehab) remain critical
- Discard vial 28 days after initial puncture — do not use beyond this date regardless of remaining volume

BPC-157 — monitoring

- Recovery progress toward documented goal
- Soft tissue / tendon / musculoskeletal symptom trend
- Exercise recovery quality
- Injection site reactions
- Any systemic symptoms (rash, swelling, breathing change)
- Need for specialist evaluation
- Course completion vs. discontinuation
- Lifestyle: sleep, nutrition, structured rehab adherence

Lab protocol

Baseline before medication start, then every 16 weeks. Order in Tebra via the Quest integration with a future collection date set 12 weeks out. No STAT designation is used.

Test	Code	Test	Code
CBC w/ Differential	6399	TSH	899
CMP	10231	Free T4	866

Test	Code	Test	Code
Lipid Panel	7600	T3, Free	34429
HbA1c	496	Copper, RBC	3481
Fasting Glucose	483	Zinc, RBC	6354
Fasting Insulin	561	Ceruloplasmin	326
IGF-1 (LC/MS)	16293		

Add for men aged 45 and over: PSA, Total (Men 45+) — code 5363. Men age 45 and older only

Copper RBC, Zinc RBC and Ceruloplasmin are part of the standard panel for every patient, regardless of which agent is prescribed.

Pricing and charge codes

Provider and internal only. Never quote a price to a patient from this document without confirming with Operations.

Item	Website	Partner	Charge code
Baseline Labs (one-time, before program selection)	\$99.00	\$99.00	BASEPeptideLab
Foundation Program — BPC-157	\$249.00 / mo	\$199.00 / mo	FndnPeptide001 FndnPeptideP01
Foundation Program — Sermorelin	\$249.00 / mo	\$199.00 / mo	FndnPeptide002 FndnPeptideP02
Foundation Program — CJC-1295 / Ipamorelin	\$249.00 / mo	\$199.00 / mo	FndnPeptide003 FndnPeptideP03

Monthly figures are the recurring payment. Full 16-week cycle: \$996.00 website, \$796.00 partner. Charge codes are shown website first, partner second.

Questions and escalation

- Pharmacy or shipping issues — Operations
- Charge codes and billing — Operations
- Clinical protocol questions — Clinical Director
- Corrections to this document — Clinical Operations

Do not improvise

- Do not alter quantity, refill or days supply
- Do not paraphrase patient instructions
- Do not prescribe into a state listed as closed
- Do not quote pricing to a patient without confirming with Operations

For KORB provider use only. Not for patient distribution. KORB Health Group LLC is a management services organisation. Clinical care is delivered by the affiliated medical practice and its licensed providers.