

GHK-Cu

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB by phone or email (see contact information below).

What Is GHK-Cu

GHK-Cu is a copper-binding peptide that occurs naturally in the human body. It is found in blood plasma, saliva, and urine, and its levels decline with age. GHK-Cu is being studied for possible support of tissue health, collagen support, wound healing, and anti-inflammatory processes.

GHK-Cu contains copper as part of its structure. Copper, zinc, and ceruloplasmin are part of your standard lab panel — your provider will pay particular attention to these results while you're on GHK-Cu.

Possible Support From GHK-Cu

This medication is one possible adjunctive support option for patients with goals related to:

- Skin quality, collagen support, and tissue healing
- Anti-inflammatory support in the context of a structured program
- Wound healing and cellular tissue wellness
- Recovery physiology as an adjunctive support option

Response to GHK-Cu varies from patient to patient. It is not a cosmetic treatment or a guaranteed skin improvement product. Your provider will assess whether therapy is appropriate to continue at your follow-up visit.

How to Use GHK-Cu

Your prescription label — or the instructions given to you by your KORB clinical provider — is the authoritative source for your dose and directions. (Some pharmacies label vials “as directed by provider” rather than printing exact instructions on the label.) What follows is a general reference. **Do not adjust anything without speaking to your provider first.**

How to inject	Schedule	Timing	Gateway / Peak (add-on)
Subcutaneous (SQ) injection — fatty tissue under the skin	3 times per week — evenings (e.g. Mon / Wed / Fri)	Evening — choose consistent days	Begins two weeks after you start BPC-157, active for 4 weeks

Copper / Zinc Lab Monitoring

Copper, zinc, and ceruloplasmin are part of your standard lab panel. While you're on GHK-Cu, your provider reviews these results closely at baseline and at each follow-up visit. Therapy may be discontinued if copper or ceruloplasmin levels trend upward.

Choosing Your 3 Days

Choose 3 consistent days each week — for example, Monday, Wednesday, and Friday. Try to keep the same days each week to build a reliable routine.

Missed a Dose?

- If you miss a dose on a scheduled day: skip it — do not double up the next day

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep your medication with you — in your carry-on or personal item, not checked baggage — in its original labeled vial or packaging. It doesn't need to be packed in ice or kept refrigerated during travel; just keep it away from excessive heat (above 86°F / 30°C) and from freezing. Once you reach your destination, refrigerate it again. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Storage and Handling

Refrigerate 36°F – 46°F (2°C – 8°C)	Do not freeze Freezing damages medication	Protect from light Keep away from direct sunlight	Discard at 28 days Write open date on vial when opened
---	---	---	--

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

What to Expect — Realistic Timeline

Timeline	Phase	What to Expect
First 2 weeks of GHK-Cu	Active course begins	Most patients notice nothing initially. This is normal. The therapy works gradually.
Weeks 3–4 of GHK-Cu	Active course continues	Skin quality or tissue changes may begin to emerge. Individual response varies from patient to patient.
After your GHK-Cu course	Washout — Labs & Lifestyle Focus	No GHK-Cu injections. Your copper, zinc, and ceruloplasmin results will be reviewed at your 16-week follow-up visit, and this is a good window to focus on nutrition, exercise, sleep, and other lifestyle habits before your next course.

Copper, zinc, and ceruloplasmin are part of your standard lab panel. Your provider will review these values closely at baseline and at your follow-up visit while GHK-Cu is active. Therapy may be discontinued if copper or ceruloplasmin levels rise.

Side Effects and What to Watch For

What You May Notice — Common	What to Do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell KORB provider if persistent.
Mild skin changes at injection site	Occasional mild discoloration or sensitivity. Usually self-limiting.
Fatigue	Uncommon; tell KORB provider if persistent.

Worth Monitoring — Tell KORB Provider at Your Next Visit

Copper or ceruloplasmin trending up on labs	Your provider will review at your follow-up visit. Therapy may be discontinued if levels rise.
Unusual rash, hives, or skin reaction	Tell KORB provider promptly — copper-containing compounds can occasionally cause hypersensitivity.
Any systemic symptom	Tell KORB provider. Do not continue without guidance if you have a new unexplained systemic symptom.

Stop injecting and seek emergency care immediately if you experience: trouble breathing, swelling of the face, lips, tongue, or throat, severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab Monitoring

Lab monitoring is a required part of your program — not optional. Your provider uses labs to assess your response and safety.

- Standard longevity panel (CBC, CMP, lipid panel, HbA1c, fasting glucose, fasting insulin, IGF-1, TSH, free T4, free T3, copper, zinc, and ceruloplasmin) — baseline and every 16-week follow-up.
- PSA — added for men aged 45 and older at every draw.

While you're on GHK-Cu, your provider pays particular attention to your copper, zinc, and ceruloplasmin results — therapy may be discontinued if copper or ceruloplasmin levels trend upward.

Your lab draw should be completed between weeks 12 and 14 of your active cycle, so results are back in time for your 16-week follow-up visit — some results, like IGF-1, can take about a week to come back. Unless your lab order says otherwise, plan to fast overnight and have your blood drawn first thing in the morning.

Safety Reminders

- Use GHK-Cu only as prescribed. Do not change your dose, schedule, or course length without provider direction.
- GHK-Cu is absolutely contraindicated in patients with Wilson's disease (copper metabolism disorder).
- Copper, zinc, and ceruloplasmin are part of your standard labs — don't skip your scheduled draw between weeks 12 and 14.
- Tell KORB provider if you develop any new medical condition or become pregnant.
- Your prescription label is the authoritative source for your dose and instructions.
- Use a new insulin needle with syringe for every injection — do not reuse.
- Never mix peptides — use one insulin needle with syringe for every peptide, every injection. Do not combine into one injection or syringe.
- Follow your state or local regulations for sharps disposal — use an FDA-cleared sharps container; do not place used needles or syringes in household trash unless your local guidelines specifically allow it.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations

- Questions about timing, storage, or administration

Email: info@korbhealth.com

Phone: +1 (888) 959-7299

Hours: Mon–Fri, 9 AM–6 PM CT

Patient Portal — Message Your Provider

- Injection-site irritation or redness that does not resolve
- Mild but persistent side effects
- Questions about whether to continue therapy

Seek Emergency Care Immediately

- Trouble breathing or throat / face swelling
- Severe rash or widespread hives
- Chest pain or fainting
- Severe or rapidly worsening symptoms
- Confusion or severe weakness
- Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key Reminders

Copper, zinc, and ceruloplasmin are part of your standard labs, and your provider watches them closely while you're on GHK-Cu.

Inject 3 times weekly in the evening. Choose consistent days (e.g. Mon / Wed / Fri).

Absolute contraindication: Wilson's disease.

Write the open date on your vial — discard 28 days after first use.

Short-course therapy — 4 weeks, offered as a Gateway / Peak add-on, beginning two weeks after you start BPC-157.

Contact KORB Operations by phone or email for non-urgent questions, or your provider via the Patient Portal for medical questions. Seek emergency care for severe symptoms.