

Wegovy — brand semaglutide

Weight Loss & Metabolic Health · Provider Reference

KORB Health Medical Texas PA · korb-glp1-data.js v2.20 · built 2026-09-15

Everything needed to prescribe Wegovy brand semaglutide, complete on its own. Values are copied literally into Tebra — do not paraphrase, and do not adjust quantity, refill or days supply.

At a glance

Type	Brand product — dispensed by the manufacturer, not compounded by KORB
Fulfilment	Every brand-name pathway — LillyDirect, NovoCare and the patient's local pharmacy. One fee, one code, no variation by product or channel.
Order via	Tebra standard prescription
Visit fee	\$79 · FITGLP1001
Medication pricing	Not held by KORB — see below

Brand pathway — where it is available and how to order

Available in 4 states only

TX · CA · AL · WI

Brand-name GLP-1 through the manufacturer programs is only available in Texas, California, Alabama and Wisconsin. Source: the brand prescribing document. This is a program limit, not a shipping limit — do not offer a brand pathway to a patient outside these four states.

Order via	Tebra Standard prescription (NOT Tebra Compound)
Dispensed by	NovoCare Pharmacy
Billing	Patient pays the manufacturer program or the local pharmacy directly.
Prior authorisation	KORB does not complete insurance prior authorisations

Coupons and savings cards	Not handled by KORB
Local pharmacy	A brand prescription can be sent to the patient's local pharmacy on request

Dispensing pharmacy

NovoCare Pharmacy
2400 Sand Lake Road, Suite 200B
Orlando, FL, 32809
(833) 949-5527

KORB does not complete insurance prior authorizations and does not handle coupons or manufacturer savings cards. The patient is responsible for checking pricing, coverage and every other aspect of the medication.

A brand prescription can be sent to the patient's local pharmacy on request, under the same rules. KORB will not re-send a prescription between pharmacies to chase a lower price. The patient does that comparison before ordering.

Pricing and charge codes

Wegovy pen (semaglutide) — brand name — 4-week

Program	Price	Charge code
Prescription visit fee	\$79	FITGLP1001

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Wegovy pen (semaglutide) — brand name — 8-week

Program	Price	Charge code
Prescription visit fee	\$79	FITGLP1001

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Wegovy oral tablet (semaglutide) — brand name — 30-day

Program	Price	Charge code
Prescription visit fee	\$79	FITGLP1001

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Wegovy oral tablet (semaglutide) — brand name — 60-day

Program	Price	Charge code
Prescription visit fee	\$79	FITGLP1001

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Includes: Telehealth visit, medication, shipping and supplies. Never quote pricing to a patient without confirming with Operations.

Brand product

Brand product dispensed through the manufacturer programme. Not compounded. Fulfilment, payment and shipping are handled by the manufacturer, not KORB.

Program options

Wegovy pen (semaglutide) — brand name

Program	Quantity	Refill	Days supply	When to use
4-week (28-day)	4	0	28	New prescription or any dose change. Use until the dose is stable.
8-week (56-day) — 4 pens + 1 refill	4	1	28	Clinically stable patients only.
12-week (84-day) — 4 pens + 2 refills	4	2	28	Clinically stable patients on a settled dose. Three fills of four pens, four weeks each.

Wegovy oral tablet (semaglutide) — brand name

Program	Quantity	Refill	Days supply	When to use
30-day supply	30	0	30	New prescription or any dose change. Use until the dose is stable.
60-day supply	60	0	60	Clinically stable patients only. Quantity 60, no refill — not 30 with a refill.
90-day supply	90	0	90	Clinically stable patients on a settled dose. Quantity 90, no refill.

Why the longer injectable programs carry refills

Brand pharmacies will not ship more than a 4-week supply of an injectable at a time. The 8-week and 12-week programs are therefore the SAME 4-week fill with refills, not a larger shipment - which is why days supply stays 28 on all three. It describes the fill, not the program.

Orals dispense as one fill

Orals are not subject to that ceiling and dispense as a single fill, so the 30, 60 and 90 day supplies are quantity changes with NO refill.

KORB does not control brand fulfilment

KORB does not control brand fulfilment. The prescription is sent like any other, and the patient pays the pharmacy directly. Price and coverage are between the patient and the manufacturer programme.

Tebra — copy these values literally

Do not paraphrase, and do not adjust quantity, refill or days supply.

Wegovy pen (semaglutide) — brand name

Tebra Standard prescription

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 0.25 mg

4-WEEK

Drug	Wegovy 0.25 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 0.25 mg SQ 4-week
Quantity	4
Refill	0
Days Supply	28
Patient Instructions	Inject 0.25 mg subcutaneously once weekly for 4 weeks.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 0.25 mg
8-WEEK

Drug	Wegovy 0.25 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 0.25 mg SQ 8-week
Quantity	4
Refill	1
Days Supply	28
Patient Instructions	Inject 0.25 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 0.25 mg
12-WEEK

Drug	Wegovy 0.25 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 0.25 mg SQ 12-week
Quantity	4
Refill	2
Days Supply	28
Patient Instructions	Inject 0.25 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 0.5 mg
4-WEEK

Drug	Wegovy 0.5 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 0.5 mg SQ 4-week
Quantity	4
Refill	0
Days Supply	28
Patient Instructions	Inject 0.5 mg subcutaneously once weekly for 4 weeks.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 0.5 mg
8-WEEK

Drug	Wegovy 0.5 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 0.5 mg SQ 8-week
Quantity	4
Refill	1
Days Supply	28
Patient Instructions	Inject 0.5 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 0.5 mg
12-WEEK

Drug	Wegovy 0.5 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 0.5 mg SQ 12-week
Quantity	4
Refill	2
Days Supply	28
Patient Instructions	Inject 0.5 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 1 mg
4-WEEK

Drug	Wegovy 1 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1 mg SQ 4-week
Quantity	4
Refill	0
Days Supply	28
Patient Instructions	Inject 1 mg subcutaneously once weekly for 4 weeks.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 1 mg
8-WEEK

Drug	Wegovy 1 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1 mg SQ 8-week
Quantity	4
Refill	1
Days Supply	28
Patient Instructions	Inject 1 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 1 mg
12-WEEK

Drug	Wegovy 1 mg/0.5 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1 mg SQ 12-week
Quantity	4
Refill	2
Days Supply	28
Patient Instructions	Inject 1 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 1.7 mg
4-WEEK

Drug	Wegovy 1.7 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1.7 mg SQ 4-week
Quantity	4
Refill	0
Days Supply	28
Patient Instructions	Inject 1.7 mg subcutaneously once weekly for 4 weeks.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 1.7 mg
8-WEEK

Drug	Wegovy 1.7 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1.7 mg SQ 8-week
Quantity	4
Refill	1
Days Supply	28
Patient Instructions	Inject 1.7 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 1.7 mg
12-WEEK

Drug	Wegovy 1.7 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1.7 mg SQ 12-week
Quantity	4
Refill	2
Days Supply	28
Patient Instructions	Inject 1.7 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 2.4 mg
4-WEEK

Drug	Wegovy 2.4 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 2.4 mg SQ 4-week
Quantity	4
Refill	0
Days Supply	28
Patient Instructions	Inject 2.4 mg subcutaneously once weekly for 4 weeks.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 2.4 mg
8-WEEK

Drug	Wegovy 2.4 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 2.4 mg SQ 8-week
Quantity	4
Refill	1
Days Supply	28
Patient Instructions	Inject 2.4 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 2.4 mg
12-WEEK

Drug	Wegovy 2.4 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 2.4 mg SQ 12-week
Quantity	4
Refill	2
Days Supply	28
Patient Instructions	Inject 2.4 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 7.2 mg
4-WEEK

Drug	Wegovy hd 7.2 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 7.2 mg SQ 4-week
Quantity	4
Refill	0
Days Supply	28
Patient Instructions	Inject 7.2 mg subcutaneously once weekly for 4 weeks.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 7.2 mg
8-WEEK

Drug	Wegovy hd 7.2 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 7.2 mg SQ 8-week
Quantity	4
Refill	1
Days Supply	28
Patient Instructions	Inject 7.2 mg subcutaneously once weekly.

Wegovy pen (semaglutide) — brand name · Tebra Standard prescription · 7.2 mg
12-WEEK

Drug	Wegovy hd 7.2 mg/0.75 ml subcutaneous pen injector Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 7.2 mg SQ 12-week
Quantity	4
Refill	2
Days Supply	28
Patient Instructions	Inject 7.2 mg subcutaneously once weekly.

Wegovy oral tablet (semaglutide) — brand name

Tebra Standard prescription

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 1.5 mg
30-DAY

Drug	Wegovy 1.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1.5 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 1.5 mg
60-DAY

Drug	Wegovy 1.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1.5 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 1.5 mg
90-DAY

Drug	Wegovy 1.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 1.5 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 4 mg
30-DAY

Drug	Wegovy 4 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 4 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 4 mg
60-DAY

Drug	Wegovy 4 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 4 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 4 mg
90-DAY

Drug	Wegovy 4 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 4 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 9 mg
30-DAY

Drug	Wegovy 9 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 9 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 9 mg
60-DAY

Drug	Wegovy 9 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 9 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 9 mg
90-DAY

Drug	Wegovy 9 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 9 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name · Tebra Standard prescription · 25 mg
30-DAY

Drug	Wegovy 25 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 25 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name • Tebra Standard prescription • 25 mg
60-DAY

Drug	Wegovy 25 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 25 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Wegovy oral tablet (semaglutide) — brand name • Tebra Standard prescription • 25 mg
90-DAY

Drug	Wegovy 25 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Wegovy 25 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.

Clinical reference — Semaglutide — GLP-1 receptor agonist

Indications

- Chronic weight management and long-term maintenance of weight reduction in adults with obesity, or overweight with at least one weight-related comorbid condition
- Improvement of glycemic control in adults with type 2 diabetes mellitus
- Reduction of cardiovascular risk in adults with type 2 diabetes and established cardiovascular disease
- Reduction of kidney disease progression and cardiovascular death in adults with type 2 diabetes and chronic kidney disease
- Treatment of noncirrhotic MASH with moderate-to-advanced liver fibrosis (F2-F3) in adults

What KORB treats

KORB treats weight loss only. Every other indication above stays with the patient's PCP or specialist - KORB does not manage type 2 diabetes, cardiovascular risk, chronic kidney disease or MASH, and does not order or monitor labs for them. A patient carrying one of those diagnoses is acceptable on the weight-loss program provided that care continues elsewhere.

Identifying the appropriate candidate

Age. Adults 18 years and older. KORB does not treat patients under 18.

Bmi

- BMI greater than 25 is overweight
- BMI greater than 30 is obese

Diabetes Note. KORB treats weight loss. Patients who are diabetic or pre-diabetic are acceptable. The patient must continue diabetes management with their PCP — KORB does not manage diabetes and does not order or monitor labs for it.

Definition

Long-acting glucagon-like peptide-1 (GLP-1) receptor agonist. A 31-amino-acid analogue with roughly 94% homology to human GLP-1, acylated with a C18 fatty diacid that binds albumin and extends the half-life to about one week, allowing once-weekly dosing. FDA-approved as Ozempic and Rybelsus for type 2 diabetes and as Wegovy for chronic weight management.

Mechanism

- Agonises the GLP-1 receptor.
- Glucose-dependent insulin secretion from pancreatic beta cells.
- Suppression of inappropriate glucagon release.
- Slowed gastric emptying, which prolongs satiety.
- Direct action on hypothalamic appetite centres, principally the arcuate nucleus, reducing hunger and energy intake.
- Insulin release is glucose-dependent, so hypoglycaemia risk is low as monotherapy.

Clinical evidence

- Strong randomised evidence — for the BRANDED product.
- STEP 1: mean weight loss near 15% of body weight at 68 weeks, against roughly 2.4% on placebo.
- SELECT: significant reduction in major adverse cardiovascular events in adults with overweight or obesity and established cardiovascular disease, without diabetes.
- Weight regain is common after discontinuation. Raise this at the start rather than letting it become a later surprise.

Absolute contraindications

- Personal or family history of medullary thyroid carcinoma (MTC)

- Multiple endocrine neoplasia syndrome type 2 (MEN2)
- Known hypersensitivity to semaglutide or any component of the formulation
- Pregnancy, breastfeeding, or planning pregnancy

Cautions

- History of pancreatitis — evaluate carefully before initiating
- Gastroparesis or other significant gastrointestinal motility disorder
- Active gallbladder disease or prior gallbladder-related surgical complications
- Severe renal impairment (eGFR below 30 mL/min/1.73 m²)
- Diabetic retinopathy — rapid glycaemic improvement can transiently worsen it
- History of suicidal ideation or active depression — no causal link has been established in regulatory review, but monitor and ask

Medication interactions

- Delayed gastric emptying can alter absorption of concomitant oral medication. Use caution with narrow-therapeutic-index drugs such as levothyroxine and warfarin.
- Insulin and sulfonylureas: additive hypoglycaemia risk. Dose reduction of the concomitant agent is often required. Coordinate with the prescribing PCP.
- Alcohol may increase hypoglycaemia risk and worsen gastrointestinal side effects.
- ANAESTHESIA AND PROCEDURES: delayed gastric emptying raises aspiration risk under sedation. HOLD for 7 days (one week) before an elective procedure requiring sedation, or for whatever longer period the anaesthesia team or the surgeon's office requires - their instruction takes precedence over this one. Tell the patient to disclose GLP-1 use to any surgeon, proceduralist or anaesthetist.
- Other GLP-1 receptor agonists: do not combine. Concomitant use is not recommended.

Side effects

Common

- Pain, redness or swelling at the injection site
- Nausea and vomiting
- Diarrhea
- Stomach pain
- Constipation
- Low appetite
- Headaches
- Dizziness

Less Common

- Pancreatitis
- Gastroparesis
- Bowel obstruction
- Gallstone attacks

Monitoring

- Weight and BMI at each visit
- Ask about blood pressure — monitored by the PCP, not measured by KORB
- Ask about recent HbA1c or glucose if diabetic — ordered by the PCP, not by KORB
- Gastrointestinal tolerance, especially through dose escalation
- Abdominal pain that is severe or radiates to the back — assess for pancreatitis
- Right-upper-quadrant pain — assess for gallbladder disease
- Mood and mental health
- Lean mass preservation — protein intake and resistance training
- Injection site and adherence

Patient counseling

Semaglutide works on the same pathway as a hormone your body already makes. It slows how quickly your stomach empties and reduces appetite signalling, so you feel full sooner and stay full longer. Nausea is the most common side effect and is usually worst in the first days after a dose increase. This is a compounded preparation rather than the brand-name product. Weight tends to come back if the medication stops, so we should talk about this as a long-term plan rather than a short course. Tell any surgeon or anaesthetist that you are taking this.

Escalation and discontinuation

Hard Stop

- Suspected pancreatitis — severe persistent abdominal pain, often radiating to the back, with or without vomiting. Stop the medication and evaluate.
- Hypersensitivity or anaphylaxis-type reaction.
- Pregnancy identified or confirmed.
- Medullary thyroid carcinoma or MEN2 identified at any point.
- Persistent severe vomiting with dehydration or acute kidney injury.

Flag And Reassess

- Gastrointestinal side effects that do not settle within two weeks of a dose increase — hold at the current dose or step back rather than pushing forward.
- Right-upper-quadrant pain, or gallstone symptoms.
- Rapid or excessive weight loss, or loss of lean mass.
- New or worsening depression, or any mention of suicidal thoughts.
- Worsening retinopathy in a diabetic patient during rapid glycaemic improvement.
- Plateau with no further response at the maximum tolerated dose — reassess the plan rather than continuing indefinitely.
- Patient scheduled for surgery or a procedure requiring sedation — coordinate holding.

Chart attestation

Patient counseled on semaglutide as a GLP-1 receptor agonist for chronic weight management, including that a compounded preparation is being dispensed rather than the FDA-approved branded product. Mechanism, expected time course, gastrointestinal side effects during titration, pancreatitis and gallbladder warning signs, the need to disclose GLP-1 use before any procedure requiring sedation, likelihood of weight regain on discontinuation, and the monitoring and follow-up plan were reviewed. Patient verbalized understanding and consented to proceed.

ICD-10

Note. DOCUMENTATION ONLY — NOT BILLING. Select the code matching the documented presentation.

Primary

- E66.9 — Obesity, unspecified
- E66.01 — Morbid (severe) obesity due to excess calories
- E66.3 — Overweight

Secondary

- Z68.— — Body mass index (add the matching BMI code)
- Z71.3 — Dietary counseling and surveillance
- Z72.3 — Lack of physical exercise

Clinically reviewed

Reviewed and signed off by Donald Stevenson, PA-C (Director of Clinical Operations and Lead Provider) on 2026-09-06, against korb-glp1-data.js v2.13.

Monograph fingerprint fp-ff64f4c7-5796. This document re-checks the fingerprint every time it is opened: if the monograph is edited after sign-off, this line is replaced by a notice that the sign-off no longer covers the text.

Questions and escalation

- Pharmacy or shipping — Operations
- Charge codes and billing — Operations
- Clinical protocol — Clinical Director
- Corrections to this document — Clinical Operations

Do not improvise

- Do not substitute an unavailable pharmacy
- Do not alter quantity, refill or days supply
- Do not paraphrase patient instructions
- Do not quote pricing without Operations

KORB Health Group LLC is a management services organisation. Clinical care is delivered by the affiliated medical practice and its licensed providers.