

Weight Loss Program Overview

Patient Guide

Weight Loss / GLP-1 Program · KORB Health Group · For Patient Reference

This guide is for educational reference only. It is not medical advice. Your KORB provider determines which medication and dose are right for you. If you have questions, contact KORB by phone or email (see contact information below).

What Is the Weight Loss Program

KORB's Weight Loss Program uses GLP-1 medication — Semaglutide or Tirzepatide — prescribed by your KORB provider and dispensed by a licensed U.S. pharmacy. Your provider selects your medication and starting dose based on your goals and health history, and adjusts your plan at regular follow-up visits.

Both medications are FDA-approved for weight management, and Semaglutide is also FDA-approved for type 2 diabetes and for reducing cardiovascular risk in adults with cardiovascular disease who are overweight or obese. If you are also managing diabetes or pre-diabetes, KORB can continue to provide your GLP-1 medication, but you should continue your diabetes care — including any related lab monitoring — with your primary care provider.

Your Medication Options

Semaglutide

A GLP-1 receptor agonist. Slows digestion and reduces appetite, which supports gradual, sustained weight loss. Injected once weekly. Your provider starts you at a low dose and increases it every 4 weeks as tolerated, based on your response.

Tirzepatide

A dual GIP/GLP-1 receptor agonist. Works through two hormone pathways rather than one, which may support greater appetite reduction for some patients. Injected once weekly, with the same gradual dose increase approach as Semaglutide.

Your provider will recommend which medication to start with, and may also discuss an oral (under-the-tongue) option if injections aren't the right fit for you. See your medication-specific Patient Education Handout for full dosing, timing, and storage instructions.

Pricing

For current Weight Loss Program pricing, contact KORB Operations or ask your provider at your visit.

How Your Program Works

- Your first visit includes a full history and a review of whether GLP-1 therapy is a safe, appropriate fit.
- Your provider starts you on a low dose of your chosen medication and provides your first supply.

- You'll follow up with your provider every 4–8 weeks. At each visit, your provider reviews how you're responding and decides whether to increase your dose, hold steady, or make other adjustments.
- Dose increases happen gradually and only under your provider's direction — never increase your dose on your own, even if you feel ready.
- This program does not require routine lab work to participate. If your provider has a specific clinical reason to order labs for you individually, they'll explain why and what to expect.
- Most patients remain on the program on an ongoing basis, moving to a steady maintenance dose once they reach a level that's working well for them.

What to Expect

Response to GLP-1 medication varies between patients. Many people notice reduced appetite within the first few weeks, with weight loss building gradually over months rather than happening immediately. **GLP-1 medication is a tool that supports weight loss — it is not a “magic shot” that works independently of your daily habits.** It works best paired with attention to nutrition — including eating slowly, prioritizing protein, and avoiding greasy or fried foods that commonly worsen nausea — along with regular activity and consistent sleep. See your medication-specific Patient Education Handout for detailed nutrition and lifestyle guidance.

Safety Reminders

- Use your medication only as prescribed. Do not change your dose, schedule, or route without provider direction.
- Do not combine GLP-1 medications from more than one source at the same time.
- This program is not appropriate during pregnancy, breastfeeding, or if you are planning pregnancy.
- This program is not appropriate with a personal or family history of Medullary Thyroid Carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN2).
- Tell your KORB provider about any new medical condition or medication, or if you become pregnant.
- Your prescription label is the authoritative source for your dose and directions.
- Use a new insulin needle with syringe for every injection — do not reuse syringes.
- Follow your state or local regulations for sharps disposal.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations	Patient Portal — Message Your Provider	Seek Emergency Care Immediately
• Questions about timing, storage, or administration Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	• Mild but persistent side effects • Questions about whether to continue therapy • Questions about dose increases or your response so far	• Trouble breathing or throat / face swelling • Severe rash or widespread hives • Chest pain or fainting • Severe or rapidly worsening symptoms • Confusion or severe weakness • Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on clinic days. Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key Reminders

- Your provider selects your medication and starting dose — this guide is a general orientation, not personalized medical advice.
- Dose increases happen gradually, every 4 weeks or longer, and only under provider direction.
- Follow-up visits happen every 4–8 weeks so your provider can track your response and adjust your plan.
- Routine labs are not required for this program.
- For current program pricing, contact KORB Operations or ask your provider.