

When to Contact KORB or Seek Emergency Care

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This guide is for general educational reference only. It does not replace your prescription label, pharmacy-specific instructions, your KORB provider's guidance, or medication-specific patient education materials. If you are experiencing a medical emergency, do not wait for a response from KORB — call 911 or go to the nearest emergency room.

This guide applies to your injectable Functional Health & Longevity Program therapy — Sermorelin, BPC-157, CJC-1295/Ipamorelin, GHK-Cu, or Tesamorelin — and any other injectable therapy your KORB provider prescribes as part of the program.

Quick Reference

Contact KORB Operations

- Non-emergency questions
- Medication questions
- Follow-up needs

Email: info@korbhealth.com

Phone: +1 (888) 959-7299

Hours: Mon–Fri, 9 AM–6 PM CT

Patient Portal — Message Your Provider

- Mild, persistent side effects
- Injection-site concerns (not severe)
- Unsure whether to continue therapy

Seek Emergency Care Immediately

- Trouble breathing
- Swelling of face, lips, tongue, or throat
- Chest pain or fainting
- Severe allergic reaction
- Severe abdominal pain
- Severe weakness or confusion
- High fever / signs of serious infection
- Any severe or unsafe symptom

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

If in doubt about whether a symptom is serious, seek urgent or emergency care. Do not wait for a response from KORB for symptoms that feel severe, rapidly worsening, or unsafe. Call 911 or go to the nearest emergency room.

What This Guide Covers

This guide explains when to contact KORB and when to seek urgent or emergency care while you're on a Functional Health & Longevity Program therapy.

It does not replace your prescription label, pharmacy instructions, your KORB provider's guidance, medication-specific patient education materials, or emergency medical care.

Important Safety Reminder

Use your therapy exactly as prescribed. Unless your KORB provider specifically directs otherwise, do not change your dose, frequency, timing, route, formulation, treatment schedule, or medication combination.

Do not stop, restart, double up, combine therapies, or change your plan without KORB guidance — unless you're experiencing a medical emergency or another treating clinician has instructed you to stop.

Contacting KORB Operations — Non-Medical Questions

Reach out to KORB Operations by phone or email for logistics like:

- Questions about medication instructions, timing, storage, or administration
- Pharmacy or shipping questions
- Questions about lab timing or follow-up expectations

Messaging Your Provider — Medical Questions via Patient Portal

Use the Patient Portal — your only HIPAA-compliant way to reach your provider directly — for:

- Mild side effects that are not urgent
- Questions about whether to continue therapy
- Concerns that therapy isn't helping
- Follow-up questions after a recent visit

Please include clear details so your care team can respond appropriately.

What to Include in Your Message

Whichever channel you use — Operations or the Patient Portal — include:

- Your full name and medication name
- Dose and schedule you're using
- When you started therapy
- What symptom or concern you're having, and when it started
- Whether it's improving, worsening, or staying the same
- Any recent illness, dehydration, new medication, or procedure
- Photos of injection-site reactions, rashes, or skin concerns, if relevant

Do not send urgent or emergency symptoms through routine messaging.

Medication & Injection Questions

Contact KORB if you have questions about:

- Dose instructions, timing, or storage
- Injection instructions
- Medication that looks abnormal or may have been stored incorrectly
- Pharmacy instructions that don't match what you expected
- Whether to hold medication around illness, surgery, or a procedure
- Whether another provider recommended stopping or changing therapy

Don't guess if you're unsure — contact KORB before using the medication.

Non-Emergency Side Effects

Contact KORB if you develop side effects that are persistent, worsening, unexpected, interfering with daily activities, or making you unsure whether to continue therapy. Examples include:

- Ongoing nausea, upset stomach, or GI discomfort
- Headache, fatigue, or dizziness that is mild but persistent
- Sleep changes or flushing
- Mild rash or itching
- Injection-site irritation that is not severe

Injection-Site Concerns

Contact KORB if you notice:

- Increasing redness, swelling, warmth, drainage, or increasing pain
- Persistent bruising, lumps, or skin changes that aren't improving
- A rash near the injection site
- A reaction that worsens over time
- Concern that you injected incorrectly, or that the medication, syringe, or needle was contaminated

Send a clear photo if appropriate. Do not inject into red, swollen, hot, painful, bruised, infected, broken, or abnormal skin.

Wellness or Systemic Changes

Contact KORB if you develop:

- Worsening fatigue, persistent brain fog, or significant sleep disruption
 - New or unexplained systemic symptoms
 - New swelling or fluid retention
 - New numbness or tingling
 - New weakness that is not severe or sudden
 - New symptoms after starting a therapy
 - Symptoms that make you unsure whether therapy remains appropriate
- Some symptoms may need broader evaluation and shouldn't automatically be attributed to your Longevity Program therapy.

Tell KORB Before Continuing Therapy If Something Changes

Contact KORB before continuing therapy if you develop:

- New pregnancy or breastfeeding status
- New cancer diagnosis or cancer treatment
- New serious infection or severe medical condition
- New kidney or liver problems, heart failure, or a major cardiovascular event
- New autoimmune, inflammatory, or significant GI disease
- A new medication that may interact with your current plan
- Upcoming surgery or imaging with contrast
- Restricted oral intake, significant dehydration, or acute illness

Your treatment plan may need to be held, adjusted, or reassessed.

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep your medication with you — in your carry-on or personal item, not checked baggage — in its original labeled vial or packaging. It doesn't need to be packed in ice or kept refrigerated during travel; just keep it away from excessive heat (above 86°F / 30°C) and from freezing. Once you reach your destination, refrigerate it again. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

When to Seek Emergency Care

Seek urgent or emergency care right away for symptoms that feel severe, unsafe, or rapidly worsening. Don't wait to hear back from KORB.

Seek Emergency Care Right Away For: trouble breathing or wheezing; swelling of the face, lips, tongue, or throat; severe rash or widespread hives; chest pain or fainting; severe dizziness or confusion; severe weakness; severe or rapidly worsening pain; new neurologic symptoms — weakness, numbness, confusion, trouble speaking, or loss of bladder/bowel control; severe headache with neurologic symptoms; severe abdominal pain or persistent vomiting; signs of severe dehydration; high fever or signs of serious infection; rapidly spreading redness, red streaking, severe warmth, or drainage from an injection site; severe glucose-related symptoms; or any symptom that feels urgent, severe, or unsafe.

Possible allergic reaction: trouble breathing, wheezing, or chest tightness; swelling of the face, lips, tongue, or throat; severe rash or widespread hives; fainting, severe dizziness, or a feeling like your throat is closing. Don't wait to hear back from KORB.

Possible serious infection: high fever; rapidly spreading redness or red streaking from an injection site; severe warmth, swelling, or pain at an injection site; drainage or pus with worsening symptoms; confusion or fainting. Contact KORB for mild or uncertain injection-site concerns, but seek urgent care if symptoms are severe or rapidly worsening.

Phone and email are not appropriate for emergencies. Call 911 or go to the nearest emergency room.

Key Reminders

Contact KORB Operations for non-emergency logistics; use the Patient Portal to message your provider about medical concerns.

Seek urgent or emergency care for severe, rapidly worsening, or unsafe symptoms — don't wait to hear back from KORB first.

Phone and email are not appropriate for emergencies. Call 911 or go to the ER.

Do not change your dose, timing, route, schedule, or medication combination without KORB provider guidance.

Do not continue therapy if you're unsure it's safe — message your provider via the Patient Portal for guidance.

Follow your prescription label and pharmacy-specific instructions at all times.

How to Reach KORB Operations

For anything in this guide that isn't a medical emergency and doesn't need your provider directly — medication questions, follow-up needs, or logistics like pharmacy and shipping — contact KORB Operations:

Email: info@korbhealth.com

Phone: [+1 \(888\) 959-7299](tel:+1(888)959-7299)

Hours: Monday – Friday, 9 AM – 6 PM (Central Time)

If you reach voicemail, please leave one message. Our staff will return your call within 24 hours on weekdays, excluding observed holidays.