

Foundayo — brand orforglipron

Weight Loss & Metabolic Health • Provider Reference

KORB Health Medical Texas PA • korb-glp1-data.js v2.21 • built 2026-09-15

Everything needed to prescribe Foundayo brand orforglipron, complete on its own. Values are copied literally into Tebra — do not paraphrase, and do not adjust quantity, refill or days supply.

At a glance

Type	Brand product — dispensed by the manufacturer, not compounded by KORB
Fulfilment	Every brand-name pathway — LillyDirect, NovoCare and the patient's local pharmacy. One fee, one code, no variation by product or channel.
Order via	Tebra standard prescription
Visit fee	\$79 • FITGLP1001
Medication pricing	Not held by KORB — see below

Brand pathway — where it is available and how to order

Available in 4 states only

TX • CA • AL • WI

Brand-name GLP-1 through the manufacturer programs is only available in Texas, California, Alabama and Wisconsin. Source: the brand prescribing document. This is a program limit, not a shipping limit — do not offer a brand pathway to a patient outside these four states.

Order via	Tebra Standard prescription (NOT Tebra Compound)
Dispensed by	LillyDirect Self Pay Pharmacy Solutions
Billing	Patient pays the manufacturer program or the local pharmacy directly.
Prior authorisation	KORB does not complete insurance prior authorisations

Coupons and savings cards	Not handled by KORB
Local pharmacy	A brand prescription can be sent to the patient's local pharmacy on request

Dispensing pharmacy

LillyDirect Self Pay Pharmacy Solutions
4343 Equity Dr
Columbus, OH, 432283842
(833) 432-4322

KORB does not complete insurance prior authorizations and does not handle coupons or manufacturer savings cards. The patient is responsible for checking pricing, coverage and every other aspect of the medication.

A brand prescription can be sent to the patient's local pharmacy on request, under the same rules. KORB will not re-send a prescription between pharmacies to chase a lower price. The patient does that comparison before ordering.

Pricing and charge codes

Foundayo (orforglipron) — brand name oral

Supply	Price	Charge code
Prescription visit fee	\$79	FITGLP1001

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Includes: Telehealth visit, medication, shipping and supplies. Never quote pricing to a patient without confirming with Operations.

Brand product

Brand product dispensed through the manufacturer programme. Not compounded. Fulfilment, payment and shipping are handled by the manufacturer, not KORB.

Program options

Foundayo (orforglipron) — brand name oral

Program	Quantity	Refill	Days supply	When to use
30-day supply	30	0	30	New prescription or any dose change. Use until the dose is stable.

60-day supply	60	0	60	Clinically stable patients only. Quantity 60, no refill.
90-day supply	90	0	90	Clinically stable patients on a settled dose. Quantity 90, no refill.

Orals dispense as one fill

Orals are not subject to that ceiling and dispense as a single fill, so the 30, 60 and 90 day supplies are quantity changes with NO refill.

KORB does not control brand fulfilment

KORB does not control brand fulfilment. The prescription is sent like any other, and the patient pays the pharmacy directly. Price and coverage are between the patient and the manufacturer programme.

Tebra — copy these values literally

Do not paraphrase, and do not adjust quantity, refill or days supply.

Foundayo (orforglipron) — brand name oral

Tebra Standard prescription

Foundayo (orforglipron) — brand name oral • Tebra Standard prescription • 0.8 mg		30-DAY
Drug	Foundayo 0.8 mg tablet Select this from the Tebra drop-down. Do not copy and paste.	
Name	Foundayo 0.8 mg tablet 30-day supply	
Quantity	30	
Refill	0	
Days Supply	30	
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.	

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 0.8 mg
60-DAY

Drug	Foundayo 0.8 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 0.8 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 0.8 mg
90-DAY

Drug	Foundayo 0.8 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 0.8 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 2.5 mg
30-DAY

Drug	Foundayo 2.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 2.5 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 2.5 mg
60-DAY

Drug	Foundayo 2.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 2.5 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 2.5 mg
90-DAY

Drug	Foundayo 2.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 2.5 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 5.5 mg
30-DAY

Drug	Foundayo 5.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 5.5 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 5.5 mg
60-DAY

Drug	Foundayo 5.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 5.5 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 5.5 mg
90-DAY

Drug	Foundayo 5.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 5.5 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 9 mg
30-DAY

Drug	Foundayo 9 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 9 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 9 mg
60-DAY

Drug	Foundayo 9 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 9 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 9 mg
90-DAY

Drug	Foundayo 9 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 9 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral · Tebra Standard prescription · 14.5 mg
30-DAY

Drug	Foundayo 14.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 14.5 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral • Tebra Standard prescription • 14.5 mg
60-DAY

Drug	Foundayo 14.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 14.5 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral • Tebra Standard prescription • 14.5 mg
90-DAY

Drug	Foundayo 14.5 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 14.5 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral • Tebra Standard prescription • 17.2 mg
30-DAY

Drug	Foundayo 17.2 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 17.2 mg tablet 30-day supply
Quantity	30
Refill	0
Days Supply	30
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral • Tebra Standard prescription • 17.2 mg
60-DAY

Drug	Foundayo 17.2 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 17.2 mg tablet 60-day supply
Quantity	60
Refill	0
Days Supply	60
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Foundayo (orforglipron) — brand name oral • Tebra Standard prescription • 17.2 mg
90-DAY

Drug	Foundayo 17.2 mg tablet Select this from the Tebra drop-down. Do not copy and paste.
Name	Foundayo 17.2 mg tablet 90-day supply
Quantity	90
Refill	0
Days Supply	90
Patient Instructions	Take 1 tablet by mouth once daily with or without food. Swallow whole. Do not break, crush, or chew.

Clinical reference — Orforglipron — oral non-peptide GLP-1 receptor agonist
Indications

- Reduction of excess body weight in adults with obesity
- Reduction of excess body weight in adults with overweight who have at least one weight-related comorbid condition
- Long-term maintenance of weight reduction
- Used in combination with a reduced-calorie diet and increased physical activity

What KORB treats

KORB treats weight loss. The indications above are the approved uses of the molecule and are not a list of conditions KORB manages - any other diagnosis the patient carries stays with their PCP or specialist, and KORB does not order or monitor labs for it. Adults 18 and older only.

Identifying the appropriate candidate

Age. Adults 18 years and older. KORB does not treat patients under 18.

BMI

- BMI greater than 25 is overweight
- BMI greater than 30 is obese

Diabetes Note. KORB treats weight loss. Patients who are diabetic or pre-diabetic are acceptable. The patient must continue diabetes management with their PCP — KORB does not manage diabetes and does not order or monitor labs for it.

Definition

A small-molecule, non-peptide GLP-1 receptor agonist taken orally once daily. Because it is not a peptide, it does not require the absorption enhancer that oral semaglutide depends on, and it is taken without the empty-stomach and water-volume restrictions that apply to oral semaglutide.

Mechanism

- Binds and activates the GLP-1 receptor as a small molecule rather than a peptide analogue.
- Glucose-dependent insulin secretion and glucagon suppression.
- Slowed gastric emptying and central appetite reduction, as with the class.
- Not a peptide, so it needs no absorption enhancer and carries none of the empty-stomach or water-volume restrictions that apply to oral semaglutide.

Clinical evidence

- Supported by the phase 3 programme in obesity and type 2 diabetes.
- Evidence base is smaller and shorter than for semaglutide or tirzepatide, simply because the agent is newer.
- Long-term outcome data are correspondingly limited.

Absolute contraindications

- Personal or family history of medullary thyroid carcinoma (MTC)
- Multiple endocrine neoplasia syndrome type 2 (MEN2)
- Known hypersensitivity to orforglipron or any component of the formulation
- Pregnancy, breastfeeding, or planning pregnancy

Cautions

- History of pancreatitis
- Gastroparesis or other significant gastrointestinal motility disorder
- Active gallbladder disease
- Severe renal impairment
- History of suicidal ideation or active depression — monitor and ask

Medication interactions

- Delayed gastric emptying can alter absorption of concomitant oral medication.
- Insulin and sulfonylureas: additive hypoglycaemia risk.
- ANAESTHESIA AND PROCEDURES: the same rule as the injectable GLP-1 agents. HOLD for 7 days (one week) before an elective procedure requiring sedation, or for whatever longer period the anaesthesia team or the surgeon's office requires - their instruction takes precedence. The patient must disclose use before any procedure requiring sedation.
- Other GLP-1 receptor agonists: do not combine.
- Verify the current prescribing information — the interaction profile for this agent is less established than for the older molecules.

Side effects

Common

- Pain, redness or swelling at the injection site
- Nausea and vomiting
- Diarrhea
- Stomach pain
- Constipation
- Low appetite
- Headaches
- Dizziness

Less Common

- Pancreatitis
- Gastroparesis
- Bowel obstruction
- Gallstone attacks

Monitoring

- Weight and BMI at each visit
- Ask about blood pressure — monitored by the PCP, not measured by KORB
- Ask about recent HbA1c or glucose if diabetic — ordered by the PCP, not by KORB
- Gastrointestinal tolerance through titration
- Abdominal pain that is severe or radiates to the back — assess for pancreatitis
- Mood and mental health
- Lean mass preservation — protein intake and resistance training
- Adherence — daily dosing is easier to miss than weekly

Patient counseling

This is a once-daily tablet that works on the same gut hormone pathway as the injectable medications, but it is a different kind of molecule, so you can take it with or without food and without the water restrictions that apply to some other oral options. Swallow it whole. Nausea is the most common side effect and is usually worst after a dose increase. It is newer than the injectables, so there is less long-term data. Tell any surgeon or anaesthetist that you take it.

Escalation and discontinuation

Hard Stop

- Suspected pancreatitis — severe persistent abdominal pain, often radiating to the back, with or without vomiting. Stop the medication and evaluate.
- Hypersensitivity or anaphylaxis-type reaction.
- Pregnancy identified or confirmed.
- Medullary thyroid carcinoma or MEN2 identified at any point.
- Persistent severe vomiting with dehydration or acute kidney injury.

Flag And Reassess

- Gastrointestinal side effects that do not settle within two weeks of a dose increase — hold at the current dose or step back rather than pushing forward.
- Right-upper-quadrant pain, or gallstone symptoms.
- Rapid or excessive weight loss, or loss of lean mass.
- New or worsening depression, or any mention of suicidal thoughts.
- Worsening retinopathy in a diabetic patient during rapid glycaemic improvement.
- Plateau with no further response at the maximum tolerated dose — reassess the plan rather than continuing indefinitely.
- Patient scheduled for surgery or a procedure requiring sedation — coordinate holding.

Chart attestation

Patient counseled on orforglipron as an oral GLP-1 receptor agonist for chronic weight management. Mechanism, once-daily administration, expected time course, gastrointestinal side effects during titration, pancreatitis warning signs, the more limited long-term evidence base relative to the injectable agents, the need to disclose use before any procedure requiring sedation, and the monitoring and follow-up plan were reviewed. Patient verbalized understanding and consented to proceed.

ICD-10

Note. DOCUMENTATION ONLY — NOT BILLING. Select the code matching the documented presentation.

Primary

- E66.9 — Obesity, unspecified
- E66.01 — Morbid (severe) obesity due to excess calories

- E66.3 — Overweight

Secondary

- Z68.— — Body mass index (add the matching BMI code)
- Z71.3 — Dietary counseling and surveillance
- Z72.3 — Lack of physical exercise

Clinically reviewed

Reviewed and signed off by Donald Stevenson, PA-C (Director of Clinical Operations and Lead Provider) on 2026-09-06, against korb-glp1-data.js v2.13.

Monograph fingerprint fp-2c45be91-4791. This document re-checks the fingerprint every time it is opened: if the monograph is edited after sign-off, this line is replaced by a notice that the sign-off no longer covers the text.

Questions and escalation

- Pharmacy or shipping — Operations
- Charge codes and billing — Operations
- Clinical protocol — Clinical Director
- Corrections to this document — Clinical Operations

Do not improvise

- Do not substitute an unavailable pharmacy
- Do not alter quantity, refill or days supply
- Do not paraphrase patient instructions
- Do not quote pricing without Operations

KORB Health Group LLC is a management services organisation. Clinical care is delivered by the affiliated medical practice and its licensed providers.