

Add-On Clinical Reference

Provider Source of Truth · Sexual Health · Hair Loss · Skin Care · Anti-Aging

KORB Health Medical Texas PA · Add-On Clinical Reference v2.0 · effective 2026-09-12 · built 2026-09-15

This is the single source of truth for all KORB add-on products. Add-ons attach to any of the four main programs. The program references link here rather than repeating this content, so a product is never described in two places.

Supersedes. The Add-On Clinical Reference generated 2026-09-10, and before it the individual Get Intimate Now, Hair Loss and Skin Care documents dated 11/26/2025.

How add-ons work

Add-ons are not standalone programs. They attach to whichever program the patient is already enrolled in: Weight Loss and Metabolic Health, Men's Health, Women's Health, or Functional Health and Longevity. A patient in any of those four can be offered any appropriate add-on.

Two products are gender-specific and must not be crossed. Finasteride is for men only. Spironolactone is the women's oral agent. The topical formulations also differ by gender and are not interchangeable. Use the Add-On Selector to confirm before prescribing.

Add-on	Men	Women	Follow-up
Sexual Health	PERFORM, KORB Rise, PT-141 nasal spray, PT-141 injection	KORB Electric, PT-141 nasal spray, PT-141 injection	No follow-up scheduled. Patient requests refills through the portal; provider completes an async visit.
Hair Loss	Finasteride 1 mg, Topical spray, Topical foam	Spironolactone 50 mg, Topical spray, Topical foam	90-day check. No routine labs.
Skin Care	Combo cream, Estriol cream 0.3%, Tretinoin cream	Combo cream, Estriol cream 0.3%, Tretinoin cream	As needed. No labs.
Anti-Aging	Metformin ER 500 mg, NAD+, NAD+ nasal spray, NAD+ FastSL sublingual	Metformin ER 500 mg, NAD+, NAD+ nasal spray, NAD+ FastSL sublingual	NAD+ every 28 days. Metformin every 90 days.

Not every formulation is available to every patient. Availability depends on the state the patient is in, the pharmacy, and what that pharmacy currently compounds. Confirm in the Add-On provider tool before quoting a product to a patient.

Hair loss products are for patients 18 and older. KORB does not treat patients under 18.

Pharmacy and state availability

Every add-on is sourced from Premier or FarmaKeio, except in California. Neither ships to California.

California is not a blanket exclusion. Belmar ships to all 50 states and DC, and supplies PT-141 and the NAD+ forms there. Those two have vetted pricing. Everything else Belmar could source is priced above what KORB charges patients once shipping is added, so it would dispense at a loss. That is a cost decision, not a licensure one, and it is under review.

Confirm the patient's state before offering an add-on, so you are not withdrawing something you have already raised.

Pharmacy	Footprint	Sources
Premier Pharmacy	Licensed shipping list. Premier cannot ship anywhere outside it, including California. Confirmed 2026-08-09.	KORB Rise, KORB Electric, PT-141 injection, all hair loss, all skin care, NAD+ injection, metformin
FarmaKeio Pharmacy	Ships to 49 states plus DC. California is the single state FarmaKeio will not ship to, and that cannot be overridden. Preferred everywhere except CA, TX, FL, AZ and NV.	PERFORM, KORB Electric, hair loss, NAD+ injection, metformin. No skin care.
Belmar Pharmacy	Ships to all 50 states and DC. Preferred for California only.	California only in practice: PT-141 nasal spray, NAD+ nasal spray, NAD+ FastSL sublingual.

Routing rule

If the patient's GLP-1 already comes from FarmaKeio, take the add-on from FarmaKeio when it stocks the product. Otherwise Premier.

Skin care is Premier only. It cannot be routed to FarmaKeio.

Sexual health

Product	Pharmacy	Who	Dosing	Supply
KORB Electric	FarmaKeio Pharmacy	Women	Apply 1-2 clicks to the clitoris 30 min before sexual activity	One bottle, approx. 20 encounters
KORB Electric	Premier Pharmacy	Women	Apply 1-2 clicks to the clitoris 30 min before sexual activity	One bottle, approx. 20 encounters
KORB Rise	Premier Pharmacy	Men	½ to 1 troche 30 min before intercourse, increase by ½ as needed. Up to 36 hours.	20 troches, approx. 20-40 encounters
PERFORM	FarmaKeio Pharmacy	Men	½ to 1 ODT 30 min before intercourse, increase by ½ as needed. Up to 36 hours.	20 ODTs, approx. 20-40 encounters

Product	Pharmacy	Who	Dosing	Supply
PT-141 injection	Premier Pharmacy	Men and women	Inject 1 to 2 mg subcutaneously about 2 hours before anticipated activity	28-day supply
PT-141 nasal spray	Belmar Pharmacy	Men and women	One spray in each nostril 1 to 2 hours before activity	28-day supply

Pricing and charge codes are deliberately not in this document. They are maintained in the Add-On provider tool, which is the single place they are kept current.

Absolute contraindication: nitrates

Patients taking nitrates for chest pain must not receive KORB Rise, PERFORM or KORB Electric. The combination can cause a life-threatening drop in blood pressure. Ask directly at every visit rather than relying on the medication list.

PT-141 is off the 90-day cadence

The labelled beyond-use date is 45 days, but the 28-day-after-first-puncture rule governs, so it is dispensed as a 28-day supply. Do not schedule refills alongside quarterly visits. The patient contacts KORB when they need more.

Raise skin darkening before starting, not after

It is a known effect of the PT-141 mechanism rather than a reaction, and patients who are not warned tend to stop.

Prescribing detail

KORB Electric

Arginine HCl 6% / Sildenafil 2% / DHEA 1% topical — FarmaKeio Pharmacy

Sildenafil 1.5% / Theophylline 3% / L-Arginine 6% topical — Premier Pharmacy

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - KORB Electric Topical - 90-Day Supply

Drug Formulation	Arginine HCl/Sildenafil/DHEA 6/2/1% topical
Name	FARMAKEIO - KORB Electric Topical - 90-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	90
Patient Instructions	KORB ELECTRIC Apply 1-2 clicks topically to the clitoris 30 minutes before a sexual encounter
Reason for Compounding	Female arousal topical
Pharmacy Instructions	Bill to office / ship to patient. No FDA-approved or commercially available equivalent.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - KORB Electric Topical - 90-Day Supply

Drug Formulation	Sildenafil/Theophylline/L-Arginine 1.5/3/6% topical
Name	PREMIER - KORB Electric Topical - 90-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	90
Patient Instructions	KORB ELECTRIC Apply 1-2 clicks topically to the clitoris 30 minutes before a sexual encounter
Reason for Compounding	Female arousal topical
Pharmacy Instructions	Bill to office / ship to patient. No FDA-approved or commercially available equivalent.

KORB Rise

Sildenafil 50 mg / Tadalafil 10 mg / L-Arginine 90 mg / Oxytocin 0.04 mg troche

Sildenafil and tadalafil increase blood flow. L-Arginine reduces headache, the most common side effect. Oxytocin was added to the formulation in September 2026. Side effects: headache, nasal congestion, blurred vision, nausea, upset stomach, prolonged erection. Advise emergency care for an erection lasting more than 4 hours.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - KORB Rise Troche - 90-Day Supply

Drug Formulation	Sildenafil/Tadalafil/L-Arginine/Oxytocin 50/10/90/0.04mg troche
Name	PREMIER - KORB Rise Troche - 90-Day Supply
Quantity	20
Unit	troche
Refill	0
Days Supply	90
Patient Instructions	KORB RISE Take 1/2 - 1 troche 30 minutes before intercourse as needed
Reason for Compounding	ED combination therapy
Pharmacy Instructions	Bill to office / ship to patient. No FDA-approved equivalent.

PERFORM

Sildenafil 88 mg / Tadalafil 22 mg rapid ODT

No L-Arginine in this formulation, so headache may be more noticeable than with KORB Rise. Advise emergency care for an erection lasting more than 4 hours.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - PERFORM Rapid ODT - 90-Day Supply

Drug Formulation	Sildenafil/Tadalafil 88/22mg rapid ODT
Name	FARMAKEIO - PERFORM Rapid ODT - 90-Day Supply
Quantity	20
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	PERFORM Rapid ODT: Take 1/2 - 1 ODT 30 minutes before intercourse as needed
Reason for Compounding	ED combination therapy
Pharmacy Instructions	Bill to office / ship to patient. No FDA-approved or commercially available equivalent.

PT-141 injection

Bremelanotide 10 mg/mL, 5 mL vial

Tebra favorite fields are not yet defined for this product. Build the prescription from the Add-On Clinical Reference.

Acts on the nervous system to increase desire and arousal, independent of hormone levels. Side effects: nausea, dizziness, fatigue, cough, skin darkening. Raise skin darkening before starting. Does not follow the 90-day cadence.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - PT-141 Injection - 28-Day Supply

Drug Formulation	Bremelanotide 10mg/mL 5mL vial
Name	PREMIER - PT-141 Injection - 28-Day Supply
Quantity	1
Unit	vial
Refill	0
Days Supply	28
Patient Instructions	Inject 1 to 2 mg subcutaneously about 2 hours before intercourse as needed
Pharmacy Instructions	Bill to office / ship to patient.

PT-141 nasal spray

Bremelanotide 5 mg/mL, 15 mL bottle

Tebra favorite fields are not yet defined for this product. Build the prescription from the Add-On Clinical Reference.

Acts on the nervous system to increase desire and arousal, independent of hormone levels. Side effects: nausea, dizziness, fatigue, cough, skin darkening. Raise skin darkening before starting, not after - it is a known effect of the mechanism and patients who are not warned tend to stop. Does not follow the 90-day cadence; do not schedule refills alongside quarterly visits.

Belmar Pharmacy · Compounded Drug Favorite Entry · BELMAR - PT-141 Nasal Spray - 28-Day Supply

Drug Formulation	Bremelanotide 5mg/mL nasal spray 15mL
Name	BELMAR - PT-141 Nasal Spray - 28-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	28
Patient Instructions	One spray in each nostril 1 to 2 hours before intercourse as needed
Pharmacy Instructions	Bill to office / ship to patient.

6 of 6 formulation strings in this section were carried forward from the prior documentation and were approved when those programs launched. They will be reviewed as each program is re-evaluated.

Hair loss

Men and women, 18 and older. Oral and topical are billed separately. Days supply varies by product and by how often the patient uses it. Pricing and charge codes are in the Add-On provider tool.

Product	Pharmacy	Who	Dosing	Supply
Finasteride 1 mg	FarmaKeio Pharmacy	Men	One tablet daily	90 tablets, 90 days
Finasteride 1 mg	Premier Pharmacy	Men	One tablet daily	90 tablets, 90 days
Spironolactone 50 mg	FarmaKeio Pharmacy	Women	50 mg daily. Increase to 100 mg at the 90-day follow-up if needed.	90 tablets, 90 days
Spironolactone 50 mg	Premier Pharmacy	Women	50 mg daily. Increase to 100 mg at the 90-day follow-up if needed.	90 tablets, 90 days
Topical foam	Premier Pharmacy	Women	Apply 1 to 2 pumps to dry hair once daily	42 days at 2/day, up to 85 at 1/day
Topical foam	Premier Pharmacy	Men	Apply 1 to 2 pumps to dry hair once daily	42 days at 2/day, up to 85 at 1/day
Topical spray	FarmaKeio Pharmacy	Women	Apply 1 to 2 sprays to dry hair once daily, leave on at least 4 hours or overnight	60 days at 2/day, up to 120 at 1/day
Topical spray	FarmaKeio Pharmacy	Men	Apply 1 to 2 sprays to dry hair once daily, leave on at least 4 hours or overnight	60 days at 2/day, up to 120 at 1/day

Pricing and charge codes are deliberately not in this document. They are maintained in the Add-On provider tool, which is the single place they are kept current.

Never prescribe finasteride to a woman

Finasteride is not prescribed to women at KORB under any circumstance, regardless of childbearing potential. It is teratogenic. Women use spironolactone as the oral agent.

Do not label topicals as a 90-day supply

Neither product lasts 90 days at twice-daily use, and the FarmaKeio spray exceeds 90 days at once-daily use. Enter the conservative figure in the Days field so the patient is never left short, and tell them the realistic range. Premier foam, 60 mL, one pump is 0.7 mL: 85 days at once daily, 42 at twice daily, Days field 42. FarmaKeio spray, 30 mL and 120 sprays: 120 days at once daily, 60 at twice daily, Days field 60.

Counsel on timeline up front

Both oral agents take 3 to 6 months to show results. Patients who are not told this discontinue early and conclude the product failed. Spironolactone is also used for acne, hirsutism and PCOS, which may be relevant for Women's Health patients.

Prescribing detail**Finasteride 1 mg**

Finasteride 1 mg tablet

Never prescribe finasteride to women. Teratogenic.

3 to 6 months before visible results. Side effects: reduced libido, mood changes, gynecomastia, orthostatic hypotension.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - Finasteride 1 mg - 90-Day Supply

Drug Formulation	Finasteride 1mg tablet
Name	FARMAKEIO - Finasteride 1 mg - 90-Day Supply
Quantity	90
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	Take one tablet daily for hair loss
Pharmacy Instructions	Bill to office/Ship to patient

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Finasteride 1 mg - 90-Day Supply

Drug Formulation	Finasteride 1mg tablet
Name	PREMIER - Finasteride 1 mg - 90-Day Supply
Quantity	90
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	Take one tablet daily for hair loss
Pharmacy Instructions	Bill to office/Ship to patient

Spironolactone 50 mg

Spironolactone 50 mg tablet

3 to 6 months before visible results. Side effects: dizziness, low energy, raised potassium, breast enlargement and tenderness. Also used for acne, hirsutism, PCOS.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - Spironolactone 50 mg - 90-Day Supply

Drug Formulation	Spironolactone 50mg tablet
Name	FARMAKEIO - Spironolactone 50 mg - 90-Day Supply
Quantity	90
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	Take one tablet daily for hair loss
Pharmacy Instructions	Bill to office/Ship to patient

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Spironolactone 50 mg - 90-Day Supply

Drug Formulation	Spironolactone 50mg tablet
Name	PREMIER - Spironolactone 50 mg - 90-Day Supply
Quantity	90
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	Take one tablet daily for hair loss
Pharmacy Instructions	Bill to office/Ship to patient

Topical foam

Minoxidil 2% / Spironolactone 0.05% / Latanoprost 0.01% — 60 mL foam — Premier Pharmacy

Minoxidil 6% / Finasteride 0.3% / Tretinoin 0.025% — 60 mL foam — Premier Pharmacy

Days field is 42, not 90. This product does not last 90 days at two applications daily.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Minoxidil 2% Topical Foam - 42-Day Supply

Drug Formulation	Minoxidil 2% / Spironolactone 0.05% / Latanoprost 0.01% topical foam, 60 mL bottle
Name	PREMIER - Minoxidil 2% Topical Foam - 42-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	42
Patient Instructions	Apply 1-2 pumps topically to dry hair once daily
Reason for Compounding	Multi-active combination
Pharmacy Instructions	Bill to office / ship to patient. No equivalent commercial option.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Minoxidil 6% Topical Foam - 42-Day Supply

Drug Formulation	Minoxidil 6% / Finasteride 0.3% / Tretinoin 0.025% topical foam, 60 mL bottle
Name	PREMIER - Minoxidil 6% Topical Foam - 42-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	42
Patient Instructions	Apply 1-2 pumps topically to dry hair once daily
Reason for Compounding	Multi-active combination
Pharmacy Instructions	Bill to office / ship to patient. No equivalent commercial option.

Topical spray

Minoxidil 2% / Spironolactone 0.05% / Latanoprost 0.01% — 30 mL spray, 120 sprays — FarmaKeio Pharmacy

Minoxidil 6% / Finasteride 0.3% / Tretinoin 0.025% — 30 mL spray, 120 sprays — FarmaKeio Pharmacy

Days field is 60, not 90. This product does not last 90 days at two applications daily.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - Minoxidil 2% Topical Spray - 60-Day Supply

Drug Formulation	Minoxidil 2% / Spironolactone 0.05% / Latanoprost 0.01% topical spray, 30 mL (120 sprays)
Name	FARMAKEIO - Minoxidil 2% Topical Spray - 60-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	60
Patient Instructions	Apply 1-2 sprays topically once daily to dry hair, leave on for at least 4 hours or overnight
Reason for Compounding	Multi-active combination
Pharmacy Instructions	Bill to office / ship to patient. No equivalent commercial option.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - Minoxidil 6% Topical Spray - 60-Day Supply

Drug Formulation	Minoxidil 6% / Finasteride 0.3% / Tretinoin 0.025% topical spray, 30 mL (120 sprays)
Name	FARMAKEIO - Minoxidil 6% Topical Spray - 60-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	60
Patient Instructions	Apply 1-2 sprays topically once daily to dry hair, leave on for at least 4 hours or overnight
Reason for Compounding	Multi-active combination
Pharmacy Instructions	Bill to office / ship to patient. No equivalent commercial option.

8 of 8 formulation strings in this section were carried forward from the prior documentation and were approved when those programs launched. They will be reviewed as each program is re-evaluated.

Skin care

Topical treatment for acne, oily skin, dry skin, wrinkles, loss of elasticity and discolouration. Available to men and women. Most patients add this to an existing program. Tretinoin is for acne, oily skin, scarring and spots; estriol is for dry skin, wrinkles, spots and elasticity. All creams are applied once daily.

Skin care is Premier only and cannot be routed to FarmaKeio. Patients in FarmaKeio-only states cannot receive it.

Product	Pharmacy	Who	Dosing	Supply
Combo cream	Premier Pharmacy	Men and women	Apply 1 click (0.25 mL) once daily	120 days
Estriol cream 0.3%	Premier Pharmacy	Men and women	Apply 1 click (0.25 mL) once daily	120 days
Tretinoin cream	Premier Pharmacy	Men and women	Apply a pea-sized amount once daily	approx. 80 days

Pricing and charge codes are deliberately not in this document. They are maintained in the Add-On provider tool, which is the single place they are kept current.

The days figures are the conservative ones, and they are not 90

Tretinoin runs about 80 days on a 20 g tube at a pea-sized amount. Estriol 0.3% and the combo cream both run 120 days on 30 mL at one click, 0.25 mL, once daily. Enter those figures, not 90. Earlier versions of this document carried a 90-day sig line that contradicted its own supply table; the supply table was right.

Estriol contains a hormone

It can cause breast tenderness or bleeding. Consider carefully before adding it for a patient already on hormone therapy in Women's Health, and counsel accordingly. General side effects across skin care: dry skin and flaking, sun sensitivity, redness, itching. Counsel on sun protection.

Prescribing detail

Combo cream

Estriol 0.3% / GHK-Cu 0.3% / Hyaluronic Acid 0.2% / Niacinamide 0.5% — 30 mL

Contains estriol, a hormone. Same consideration as above.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Estriol Combo Cream - 120-Day Supply

Drug Formulation	Estriol 0.3% / GHK-Cu 0.3% / Hyaluronic Acid 0.2% / Niacinamide 0.5% cream, 30 mL
Name	PREMIER - Estriol Combo Cream - 120-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	120
Patient Instructions	Apply (1 click/0.25 ml/0.25 g) to the affected area once daily as directed
Reason for Compounding	Multi-active combination
Pharmacy Instructions	Bill to office / ship to patient. No FDA-approved or commercially available equivalent.

Estriol cream 0.3%

Estriol 0.3% — 30 mL

Contains a hormone. Can cause breast tenderness or bleeding. Consider carefully for a patient already on hormone therapy.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Estriol Cream 0.3% - 120-Day Supply

Drug Formulation	Estriol 0.3% topical cream, 30 mL
Name	PREMIER - Estriol Cream 0.3% - 120-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	120
Patient Instructions	Apply (1 click/0.25 ml/0.25 g, containing estriol 0.075%) to the affected external area once daily as directed
Reason for Compounding	Compounded strength
Pharmacy Instructions	Bill to office / ship to patient. No FDA-approved or commercially available equivalent.

Tretinoin cream

0.025%, 0.05%, or 0.1% — 20 g tube

For acne, oily skin, scarring, spots. Causes sun sensitivity — counsel on sun protection.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Tretinoin Cream 0.025% - 80-Day Supply

Drug Formulation	Tretinoin 0.025% topical cream, 20 g tube
Name	PREMIER - Tretinoin Cream 0.025% - 80-Day Supply
Quantity	1
Unit	tube
Refill	0
Days Supply	80
Patient Instructions	Apply a pea-sized amount daily as needed
Reason for Compounding	Strength not commercial
Pharmacy Instructions	20 gm Tube, Bill to office/Ship to patient

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Tretinoin Cream 0.05% - 80-Day Supply

Drug Formulation	Tretinoin 0.05% topical cream, 20 g tube
Name	PREMIER - Tretinoin Cream 0.05% - 80-Day Supply
Quantity	1
Unit	tube
Refill	0
Days Supply	80
Patient Instructions	Apply a pea-sized amount daily as needed
Reason for Compounding	Strength not commercial
Pharmacy Instructions	20 gm Tube, Bill to office/Ship to patient

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Tretinoin Cream 0.1% - 80-Day Supply

Drug Formulation	Tretinoin 0.1% topical cream, 20 g tube
Name	PREMIER - Tretinoin Cream 0.1% - 80-Day Supply
Quantity	1
Unit	tube
Refill	0
Days Supply	80
Patient Instructions	Apply a pea-sized amount daily as needed
Reason for Compounding	Strength not commercial
Pharmacy Instructions	20 gm Tube, Bill to office/Ship to patient

3 of 3 formulation strings in this section were carried forward from the prior documentation and were approved when those programs launched. They will be reviewed as each program is re-evaluated.

Anti-aging: NAD+ and metformin

Longevity-focused add-on, available to patients in any program. NAD+ and metformin are prescribed independently of each other; a patient may receive either or both. No laboratory testing is required to start either product, though metformin has monitoring requirements once started.

NAD+ is a naturally occurring coenzyme involved in metabolism, DNA repair and immune function, and levels decline with age. Injection is the fastest route, usually into the lower abdomen, and may also be given in the arm, buttock or thigh.

Product	Pharmacy	Who	Dosing	Supply
Metformin ER 500 mg	FarmaKeio Pharmacy	Men and women	One tablet daily	90 tablets, 90 days
Metformin ER 500 mg	Premier Pharmacy	Men and women	One tablet daily	90 tablets, 90 days
NAD+	FarmaKeio Pharmacy	Men and women	Start 50 mg SQ twice weekly. Increase to 100 mg twice weekly if needed.	28 days
NAD+	Premier Pharmacy	Men and women	Start 50 mg SQ twice weekly. Increase to 100 mg twice weekly if needed.	28 days
NAD+ FastSL sublingual	Belmar Pharmacy	Men and women	Start half a tablet under the tongue each morning, Monday to Friday, off at weekends. Increase to one tablet.	28 days
NAD+ nasal spray	Belmar Pharmacy	Men and women	Start 1 spray per nostril once daily. Increase to 2 sprays per nostril once daily.	28 days

Pricing and charge codes are deliberately not in this document. They are maintained in the Add-On provider tool, which is the single place they are kept current.

Every NAD+ form is a 28-day supply

That is driven by beyond-use dating on a compounded product. The injection vial holds 1000 mg, which outlasts 28 days at either dose, so the patient discards the remainder. Say so on the prescription. NAD+ follow-up is every 28 days, not on the 90-day cadence.

Metformin is not contraindication-free

It carries a boxed warning for lactic acidosis. Do not prescribe with eGFR below 30 mL/min/1.73m², and do not initiate between 30 and 45. Also screen for acute or unstable heart failure or any state causing tissue hypoxia, severe hepatic impairment, heavy or binge alcohol use, and metabolic acidosis of any cause. Hold before and after iodinated contrast imaging or surgery. Renal function must be known before starting. Metformin is cleared and in use; the screening criteria here come from standard prescribing guidance because the source protocol listed none.

Metformin rationale, stated with its limits

On the WHO List of Essential Medicines. The evidence cited for lower rates of dementia and some cancers, reduced cardiovascular risk, reduced inflammation and reduced diabetes risk is largely observational for an off-label longevity indication. Patients will not feel noticeably different; set that expectation up front.

Prescribing detail

Metformin ER 500 mg

Metformin ER 500 mg tablet

Do not prescribe with eGFR under 30. Do not initiate between 30 and 45. Also screen for acute or unstable heart failure or any state causing tissue hypoxia, severe hepatic impairment, heavy or binge alcohol use, and metabolic acidosis of any cause. Boxed warning for lactic acidosis. Hold before and after iodinated contrast imaging or surgery.

Blunts training adaptations. In the MASTERS trial (randomized, double-blind, placebo-controlled, 94 adults 65+, 14 weeks of progressive resistance training) metformin 1,700 mg/day produced smaller gains than placebo in lean body mass ($p=.003$), thigh muscle mass ($p<.001$) and thigh muscle area ($p=.005$). A companion trial found blunted mitochondrial adaptations to aerobic training. Those doses are well above KORB's 500 mg ER, so the size of the effect here is unknown, but raise it with any patient whose goal is strength or performance, including gym-partner referrals.

Patient will not feel noticeably different. Set that expectation. Side effects: nausea, upset stomach, headache, fatigue, mild hypoglycaemia symptoms.

Monitoring

- Baseline: eGFR, hepatic function, and a vitamin B12 level. A longevity indication has no endpoint, so treat this as long-term therapy from day one.
- eGFR: annually at 60 or above, every 3 to 6 months at 45 to 60, every 3 months at 30 to 45 with a 50% dose reduction (FDA).
- Vitamin B12: ADA Standards of Care 2026 recommendation 3.10 (grade B) is periodic assessment on long-term metformin, especially with anemia or peripheral neuropathy. Annual is reasonable past 4 years of therapy or with independent risk factors. Deficiency is below 148 pmol/L, borderline below 200; methylmalonic acid or holotranscobalamin resolve an equivocal level.
- KORB doses 500 mg ER daily, below the 1,500 mg/day threshold where B12 risk rises most sharply, but duration is the other driver and a longevity course is indefinite.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - Metformin ER 500 mg - 90-Day Supply

Drug Formulation	Metformin ER 500 mg tablet
Name	FARMAKEIO - Metformin ER 500 mg - 90-Day Supply
Quantity	90
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	TAKE ONE TAB PO QD
Pharmacy Instructions	Bill to KORB / ship to patient

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - Metformin ER 500 mg - 90-Day Supply

Drug Formulation	Metformin ER 500 mg tablet
Name	PREMIER - Metformin ER 500 mg - 90-Day Supply
Quantity	90
Unit	tablet
Refill	0
Days Supply	90
Patient Instructions	TAKE ONE TAB PO QD
Pharmacy Instructions	Bill to KORB / ship to patient

NAD+

NAD+ 200 mg/mL, 5 mL vial (1000 mg) SQ injection

28-day supply, not 90. The vial outlasts 28 days at either dose — patient discards the remainder. Follow-up every 28 days.

Do not combine with isotretinoin (Accutane). Counsel on nausea, fatigue, headache, indigestion.

FarmaKeio Pharmacy · Compounded Drug Favorite Entry · FARMAKEIO - NAD+ Injection - 28-Day Supply

Drug Formulation	NAD+ 200mg/mL 5mL vial
Name	FARMAKEIO - NAD+ Injection - 28-Day Supply
Quantity	1
Unit	vial
Refill	0
Days Supply	28
Patient Instructions	Injectable 50 mg subcutaneously twice a week, discard unused medication 28 days after first puncture
Reason for Compounding	No FDA-approved NAD+
Pharmacy Instructions	Bill to office / ship to patient. Compounded, no FDA-approved NAD+, individualized dosing.

Premier Pharmacy · Compounded Drug Favorite Entry · PREMIER - NAD+ Injection - 28-Day Supply

Drug Formulation	NAD+ 200mg/mL 5mL vial
Name	PREMIER - NAD+ Injection - 28-Day Supply
Quantity	1
Unit	vial
Refill	0
Days Supply	28
Patient Instructions	Injectable 50 mg subcutaneously twice a week, discard unused medication 28 days after first puncture
Reason for Compounding	No FDA-approved NAD+
Pharmacy Instructions	Bill to office / ship to patient. Compounded, no FDA-approved NAD+, individualized dosing.

NAD+ FastSL sublingual

NAD+ 100 mg sublingual tablet

Tebra favorite fields are not yet defined for this product. Build the prescription from the Add-On Clinical Reference.

28-day supply. Follow-up every 28 days, not on the 90-day cadence.

Belmar Pharmacy · Compounded Drug Favorite Entry · BELMAR - NAD+ FastSL Sublingual - 28-Day Supply

Drug Formulation	NAD+ 100mg sublingual tablet
Name	BELMAR - NAD+ FastSL Sublingual - 28-Day Supply
Quantity	20
Unit	tablet
Refill	0
Days Supply	28
Patient Instructions	Dissolve 1/2 to 1 tablet under the tongue each morning Monday to Friday. No dose at weekends.
Reason for Compounding	No FDA-approved NAD+
Pharmacy Instructions	Bill to office / ship to patient. Compounded, no FDA-approved NAD+, individualized dosing.

NAD+ nasal spray

NAD+ 300 mg/mL, 15 mL bottle

Tebra favorite fields are not yet defined for this product. Build the prescription from the Add-On Clinical Reference.

28-day supply, driven by beyond-use dating on a compounded product. Follow-up every 28 days, not on the 90-day cadence.

Belmar Pharmacy · Compounded Drug Favorite Entry · BELMAR - NAD+ Nasal Spray - 28-Day Supply

Drug Formulation	NAD+ 300mg/mL nasal spray 15mL
Name	BELMAR - NAD+ Nasal Spray - 28-Day Supply
Quantity	1
Unit	bottle
Refill	0
Days Supply	28
Patient Instructions	Start 1 spray in each nostril once daily. May increase to 2 sprays in each nostril once daily.
Reason for Compounding	No FDA-approved NAD+
Pharmacy Instructions	Bill to office / ship to patient. Compounded, no FDA-approved NAD+, individualized dosing.

6 of 6 formulation strings in this section were carried forward from the prior documentation and were approved when those programs launched. They will be reviewed as each program is re-evaluated.

Documentation

- Record the add-on in the patient's chart and add it to the CC section so it carries forward.
- Document that nitrate use was asked about and excluded before prescribing any sexual health product.
- For hair loss and skin care, document the counselling on the 3 to 6 month timeline and on sun sensitivity.
- Add-ons bill separately from the base program. Confirm the charge code against the product record rather than from memory.

Questions and escalation

- Pharmacy or shipping issues — Operations
- Charge codes and billing — Clinical Operations
- Clinical protocol questions — Clinical Operations
- Corrections to this document — Clinical Operations

Route everything through Clinical Operations. Do not contact the clinical sign-off physician directly; Clinical Operations escalates when it needs to.

Do not improvise

- Do not alter quantity, refill or days supply
- Do not paraphrase patient instructions
- Do not cross a gender-specific product
- Do not quote pricing to a patient without confirming with Operations

For KORB internal and provider use only. Do not distribute to patients. Clinical care is delivered by KORB Health Medical Texas PA and its licensed providers. KORB Health Group LLC is a management services organisation and does not practise medicine.