

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB using the details at the end of this handout.

What BPC-157 is

BPC-157 is a synthetic peptide derived from a protein found naturally in the stomach lining. Its full name is Body Protection Compound 157. It is being studied for possible support of soft tissue recovery, musculoskeletal health and gut health.

What it may support

This medication is one possible adjunctive support option for patients with goals related to:

- Soft tissue recovery, including tendon, ligament and connective tissue support
- Musculoskeletal wellness during rehabilitation or recovery
- Exercise recovery quality and return-to-activity support
- Gut support in select clinical contexts, discussed with your provider

Response to BPC-157 varies from patient to patient and results are not guaranteed. Some patients notice changes during their active course and others do not. Your provider will assess whether to continue at your follow-up visit.

How to use it

Your prescription label, or the instructions given to you by your KORB clinical provider, is the authoritative source for your dose and directions. Some pharmacies label vials "as directed by provider" rather than printing exact instructions. What follows is a general reference. Do not adjust anything without speaking to your provider first.

How to inject	Subcutaneous (SQ) injection — fatty tissue under the skin
When to inject	Any consistent time · No food restriction
Schedule	Every day (7 days a week)
Foundation — active weeks	Weeks 1–8
Gateway / Peak — active weeks	Weeks 3–8

On Gateway and Peak the course starts two weeks after your start date, which is why it begins later and runs shorter.

If you miss a dose

If you miss a dose on a scheduled day, skip it. Do not double up the next day.

Storage and handling

What to do	Detail
Refrigerate	36°F – 46°F (2°C – 8°C)
Do not freeze	Freezing damages medication
Protect from light	Keep away from direct sunlight
Discard at 28 days	Write the open date on the vial

Before first use: refrigerate, protect from light, do not freeze or shake.

After opening: write the open date on your vial and discard 28 days after first use, even if medication remains. If your label shows an earlier beyond-use date, follow the earlier one.

Do not use if the solution appears cloudy or discolored, or contains particles.

Traveling with your medication

Traveling does not have to interrupt your program. Keep your medication with you, in your carry-on or personal item rather than checked baggage, in its original labeled vial or packaging. It does not need to be packed in ice during travel; keep it away from excessive heat (above 86°F / 30°C) and from freezing, and refrigerate it again when you arrive. Medication and syringes are permitted in carry-on luggage, and TSA recommends keeping the original pharmacy label visible. Crossing time zones can shift your injection time by a few hours without a problem; if you will be off your usual schedule for several days, contact KORB.

What to expect

Timeline	Phase	What to expect
Weeks 1–2	Getting started	No immediate noticeable effect is common. The therapy works gradually. Continue your schedule.
Active course	Foundation 8 weeks; Gateway and Peak 6 weeks	Recovery quality may begin to shift. Individual response varies from patient to patient.
Off weeks	Washout — labs and lifestyle	No injections during this window; this is intentional. Your labs, drawn between weeks 12 and 14, are reviewed during this time, and it is a good stretch to focus on nutrition, exercise and sleep before your next course.

Your provider will assess whether therapy is appropriate to continue based on your response, tolerability and goals.

Side effects and what to watch for

What you may notice

What you may notice	What to do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell your KORB provider if it persists.
Nausea or GI discomfort	Occasionally reported. Usually mild and self-limiting. Tell your KORB provider if it persists.
Fatigue	Uncommon. Tell your KORB provider if it persists.

Tell your KORB provider at your next visit

What you may notice	What to do
Unusual pain or swelling at the injection site	Could indicate infection. Contact KORB promptly if redness spreads or a fever develops.
Systemic rash or hives	Tell your KORB provider at your next visit, or sooner if it is spreading or severe.
Any unexplained systemic symptom	Tell your KORB provider. Do not continue without guidance if you have a significant new symptom.

Stop injecting and seek emergency care immediately if you have trouble breathing, swelling of the face, lips, tongue or throat, a severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab monitoring

Lab monitoring is a required part of your program, not optional. Your provider uses labs to assess your response and your safety.

- Standard longevity panel: CBC, CMP, lipid panel, HbA1c, fasting glucose, fasting insulin, IGF-1, TSH, free T4, free T3, copper, zinc and ceruloplasmin, at baseline and every 16-week follow-up.
- PSA, added for men aged 45 and older at every draw.

Your lab draw should be completed between weeks 12 and 14 of your active cycle, so results are back in time for your 16-week follow-up visit. Some results, such as IGF-1, can take about a week. Unless your lab order says otherwise, plan to fast overnight and have blood drawn first thing in the morning.

Safety reminders

- Use BPC-157 only as prescribed. Do not change your dose, course length or schedule without provider direction.
- BPC-157 is not appropriate during active malignancy or pregnancy.
- Do not restart the course early after your off weeks without provider direction.
- Tell your KORB provider if you develop a new medical condition or become pregnant.
- Use a new insulin needle and syringe for every injection. Do not reuse.
- Never mix peptides. Use a separate needle and syringe for each peptide and each injection; do not combine them into one syringe.
- Follow your state or local rules for sharps disposal. Use an FDA-cleared sharps container, and do not put used needles or syringes in household trash unless your local guidance specifically allows it.

When to contact KORB

Contact KORB Operations	Patient Portal — message your provider	Seek emergency care immediately
• Questions about timing, storage or administration • Scheduling, billing and shipping Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	• Injection-site irritation or redness that does not resolve • Mild but persistent side effects • Questions about whether to continue therapy	• Trouble breathing, or swelling of the face, lips, tongue or throat • Severe rash or widespread hives • Chest pain or fainting • Severe or rapidly worsening symptoms • Confusion or severe weakness • Any symptom that feels urgent or unsafe

The Patient Portal is the only HIPAA-compliant way to message your provider directly. Providers check portal messages once per day on their clinic days. For anything that does not need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key reminders

- BPC-157 is taken every day, including weekends, for the length of your course.
- Write the open date on your vial and discard 28 days after first use.
- Labs between weeks 12 and 14 are required before your 16-week follow-up visit.
- Do not restart your course early after the off weeks.

- Contact KORB Operations for non-urgent questions, your provider through the Patient Portal for medical questions, and emergency care for severe symptoms.