

BPC-157

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB by phone or email (see contact information below).

What Is BPC-157

BPC-157 is a synthetic peptide derived from a protein found naturally in the stomach lining. Its full name is Body Protection Compound 157. It is being studied for possible support of soft tissue recovery, musculoskeletal health, and gut health.

Possible Support From BPC-157

This medication is one possible adjunctive support option for patients with goals related to:

- Soft tissue recovery — tendon, ligament, and connective tissue support
- Musculoskeletal wellness during rehabilitation or recovery
- Exercise recovery quality and return-to-activity support
- Gut support in select clinical contexts (discussed with your provider)

Response to BPC-157 varies from patient to patient, and results aren't guaranteed. Some patients notice changes during their active course; others don't. Your provider will assess whether to continue at your follow-up visit.

How to Use BPC-157

Your prescription label — or the instructions given to you by your KORB clinical provider — is the authoritative source for your dose and directions. (Some pharmacies label vials “as directed by provider” rather than printing exact instructions on the label.) What follows is a general reference. **Do not adjust anything without speaking to your provider first.**

How to inject	Schedule	Timing	Active weeks
Subcutaneous (SQ) injection — fatty tissue under the skin	Every day (7 days a week)	Any consistent time daily — no food restriction	Foundation: the first 8 weeks of your 16-week cycle, followed by an 8-week washout before your next cycle begins. Gateway / Peak: a shorter 6-week course, beginning two weeks after your start date.

Missed a Dose?

- If you miss a dose on a scheduled day: skip it — do not double up the next day

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep your medication with you — in your carry-on or personal item, not checked baggage — in its original labeled vial or packaging. It doesn't need to be packed in ice or kept refrigerated during travel; just keep it away from excessive heat (above 86°F / 30°C) and from freezing. Once you reach your destination, refrigerate it again. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Storage and Handling

Refrigerate 36°F – 46°F (2°C – 8°C)	Do not freeze Freezing damages medication	Protect from light Keep away from direct sunlight	Discard at 28 days Write open date on vial when opened
---	---	---	--

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

What to Expect — Realistic Timeline

Timeline	Phase	What to Expect
Course Weeks 1–2	Getting started	No immediate noticeable effect is common. The therapy works gradually. Continue your schedule.
Foundation: first 8 weeks Gateway/Peak: 6 weeks, beginning two weeks after your start date	Active course	Recovery quality may begin to shift. Foundation runs a full 8-week course; Gateway and Peak run a shorter 6-week course beginning two weeks after your start date. Individual response varies from patient to patient.
Off weeks	Washout — Labs & Lifestyle Focus	No injections during this window — this is intentional. Your labs (drawn between weeks 12 and 14 of your cycle) are reviewed during this time, and it's a good time to focus on nutrition, exercise, sleep, and other lifestyle habits before your next course.

Your provider will assess whether therapy is appropriate to continue based on your response, tolerability, and goals.

Side Effects and What to Watch For

What You May Notice — Common	What to Do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell KORB provider if persistent.
Nausea or GI discomfort	Occasionally reported. Usually mild and self-limiting. Tell KORB provider if persistent.
Fatigue	Uncommon; tell KORB provider if persistent.

Worth Monitoring — Tell KORB Provider at Your Next Visit

Unusual pain or swelling at injection site	Could indicate infection. Contact KORB promptly if redness spreads or fever develops.
Systemic rash or hives	Tell KORB provider at next visit or sooner if spreading or severe.
Any unexplained systemic symptom	Tell KORB provider. Do not continue without guidance if you have a significant new symptom.

Stop injecting and seek emergency care immediately if you experience: trouble breathing, swelling of the face, lips, tongue, or throat, severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab Monitoring

Lab monitoring is a required part of your program — not optional. Your provider uses labs to assess your response and safety.

- Standard longevity panel (CBC, CMP, lipid panel, HbA1c, fasting glucose, fasting insulin, IGF-1, TSH, free T4, free T3, copper, zinc, and ceruloplasmin) — baseline and every 16-week follow-up.
- PSA — added for men aged 45 and older at every draw.

Your lab draw should be completed between weeks 12 and 14 of your active cycle, so results are back in time for your 16-week follow-up visit — some results, like IGF-1, can take about a week to come back. Unless your lab order says otherwise, plan to fast overnight and have your blood drawn first thing in the morning.

Safety Reminders

- Use BPC-157 only as prescribed. Do not change your dose, course length, or schedule without provider direction.
- BPC-157 is not appropriate during active malignancy or pregnancy.
- Do not restart the course early after your off weeks without provider direction.
- Tell KORB provider if you develop any new medical condition or become pregnant.
- Your prescription label is the authoritative source for your dose and instructions.
- Use a new insulin needle with syringe for every injection — do not reuse.
- Never mix peptides — use one insulin needle with syringe for every peptide, every injection. Do not combine into one injection or syringe.
- Follow your state or local regulations for sharps disposal — use an FDA-cleared sharps container; do not place used needles or syringes in household trash unless your local guidelines specifically allow it.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations

- Questions about timing, storage, or administration

Email: info@korbhealth.com

Phone: +1 (888) 959-7299

Hours: Mon–Fri, 9 AM–6 PM CT

Patient Portal — Message Your Provider

- Injection-site irritation or redness that does not resolve
- Mild but persistent side effects
- Questions about whether to continue therapy

Seek Emergency Care Immediately

- Trouble breathing or throat / face swelling
- Severe rash or widespread hives
- Chest pain or fainting
- Severe or rapidly worsening symptoms
- Confusion or severe weakness
- Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key Reminders

Foundation: BPC-157 is an 8-week course, followed by an 8-week washout before your next cycle.

Gateway / Peak: BPC-157 is a shorter 6-week course, beginning two weeks after your start date.

Inject daily at any consistent time — no food restriction.

Write the open date on your vial — discard 28 days after first use.

Your provider will assess benefit at your follow-up visit.

Contact KORB Operations by phone or email for non-urgent questions, or your provider via the Patient Portal for medical questions. Seek emergency care for severe symptoms.