

Sexual Health

Patient Education Handout

KORB Rise · KORB Electric · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label and your KORB provider's guidance. If your label differs from anything here, follow your label and contact KORB.

Do not use these medications if you take nitrates for chest pain. Nitrates include nitroglycerin, isosorbide, and similar heart medications, whether you take them daily or only occasionally. Combining them with this treatment can cause a sudden, dangerous drop in blood pressure that can be life-threatening. Tell your provider about every medication you take, including ones you use rarely.

For Men: KORB Rise

KORB Rise is a dissolvable troche you place in your mouth. It contains the same active ingredients found in Viagra and Cialis, combined with L-Arginine and oxytocin. L-Arginine is included specifically to reduce headache, which is the most common side effect.

How to take it	When	How long it lasts
Let half to one troche dissolve in your mouth. Do not swallow it whole.	About 30 minutes before sex	Up to 36 hours, so it does not need tight planning

Start with half a troche. If that is not enough, your provider may tell you to increase to a full one. Do not take more than one troche in a 24-hour period unless your provider specifically tells you to.

If you live in a state we serve through a different pharmacy, you may receive **PERFORM** instead. It is the same idea in a fast-dissolving tablet, taken 30 minutes before sex. It does not contain L-Arginine, so headache may be a little more noticeable.

For Women: KORB Electric

KORB Electric is a topical gel applied directly to the clitoris about 30 minutes before sexual activity. It works by increasing local blood flow rather than acting on the whole body.

Apply one to two clicks from the applicator. Wash your hands before and after. One bottle covers roughly 20 uses.

What to Expect

- These medications support your body's normal response. They do not create arousal on their own, so interest and stimulation still matter.

- The first attempt is not always the best measure. It is common to need a few tries to find your timing and the right amount.
- Alcohol and heavy meals both blunt the effect. A large, fatty meal beforehand is the most common reason people feel it did not work.

Side Effects

What you may notice	What to do
Headache	The most common effect. Usually mild. Stay hydrated. Tell your provider if it persists.
Stuffy nose or flushing	Common and temporary.
Upset stomach or nausea	Common. Taking it on a lighter stomach may help.
Changes in vision, such as a blue tinge or blurring	Uncommon. Stop and contact KORB if it happens.
Local irritation (KORB Electric)	Wash the area. Tell your provider if it does not settle.

Seek emergency care right away for: an erection lasting more than 4 hours, chest pain during or after sex, sudden loss of vision or hearing, fainting, or severe dizziness. An erection that will not go down is a medical emergency and can cause permanent damage if untreated.

Getting Refills

There is no lab work and no scheduled follow-up for this add-on. When you need more, send a message through the patient portal and your provider will review it. Do not wait until you are completely out.

Key Reminders

- Never combine with nitrates for chest pain.
- Take about 30 minutes before sex. Effect lasts up to 36 hours.
- Do not exceed one dose in 24 hours unless your provider directs it.
- Heavy meals and alcohol reduce how well it works.
- An erection lasting more than 4 hours needs emergency care.
- Request refills through the patient portal.