

When to Contact KORB or Seek Emergency Care

Patient Education Handout

KORB Health Group · For Patient Reference

This guide is for general educational reference only. It does not replace your prescription label, pharmacy-specific instructions, your KORB provider's guidance, or medication-specific patient education materials. If you are experiencing a medical emergency, do not wait for a response from KORB — call 911 or go to the nearest emergency room.

This guide applies across all KORB programs — Weight Loss/GLP-1, Men's Health, Women's Health, and Functional Health & Longevity — regardless of whether your medication is an injection, patch, cream, or oral tablet.

Quick Reference

Contact KORB Operations

- Questions about timing, storage, or administration

Email: info@korbhealth.com
Phone: +1 (888) 959-7299
Hours: Mon–Fri, 9 AM–6 PM CT

Patient Portal — Message Your Provider

- Mild but persistent side effects
- Questions about whether to continue therapy
- Unsure whether to continue therapy

Seek Emergency Care Immediately

- Trouble breathing or throat / face swelling
- Severe rash or widespread hives
- Chest pain or fainting
- Severe or rapidly worsening symptoms
- Confusion or severe weakness
- Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once per day on clinic days. Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

If in doubt about whether a symptom is serious, seek urgent or emergency care. Do not wait for a response from KORB for symptoms that feel severe, rapidly worsening, or unsafe. Call 911 or go to the nearest emergency room.

What This Guide Covers

This guide explains when to contact KORB and when to seek urgent or emergency care while you're on a KORB program. It does not replace your prescription label, pharmacy instructions, your KORB provider's guidance, medication-specific patient education materials, or emergency medical care.

Important Safety Reminder

Use your therapy exactly as prescribed. Unless your KORB provider specifically directs otherwise, do not change your dose, frequency, timing, route, formulation, treatment schedule, or medication combination.

Do not stop, restart, double up, combine therapies, or change your plan without KORB guidance — unless you're experiencing a medical emergency or another treating clinician has instructed you to stop.

Contacting KORB Operations — Non-Medical Questions

Reach out to KORB Operations by phone or email for logistics like:

- Questions about medication instructions, timing, storage, or administration
- Pharmacy or shipping questions
- Questions about lab timing or follow-up expectations, if labs are part of your program

Messaging Your Provider — Medical Questions via Patient Portal

Use the Patient Portal — your only HIPAA-compliant way to reach your provider directly — for:

- Mild side effects that are not urgent
- Questions about whether to continue therapy
- Concerns that therapy isn't helping
- Follow-up questions after a recent visit

Please include clear details so your care team can respond appropriately.

What to Include in Your Message

Whichever channel you use — Operations or the Patient Portal — include:

- Your full name and medication name
- Dose and schedule you're using
- When you started therapy
- What symptom or concern you're having, and when it started
- Whether it's improving, worsening, or staying the same
- Any recent illness, dehydration, new medication, or procedure
- Photos of injection-site reactions, rashes, or skin concerns, if relevant

Do not send urgent or emergency symptoms through routine messaging.

Medication Questions

Contact KORB if you have questions about:

- Dose instructions, timing, or storage
- How to use your medication (injection, patch, cream, or oral)
- Medication that looks abnormal or may have been stored incorrectly
- Pharmacy instructions that don't match what you expected
- Whether to hold medication around illness, surgery, or a procedure
- Whether another provider recommended stopping or changing therapy

Don't guess if you're unsure — contact KORB before using the medication.

Non-Emergency Side Effects

Contact KORB if you develop side effects that are persistent, worsening, unexpected, interfering with daily activities, or making you unsure whether to continue therapy. Examples include:

- Ongoing nausea, upset stomach, or GI discomfort
- Headache, fatigue, or dizziness that is mild but persistent
- Sleep changes or flushing
- Mild rash or itching
- Injection-site, patch-site, or application-site irritation that is not severe

Injection or Application Site Concerns

If your therapy is an injection, patch, or topical medication, contact KORB if you notice:

- Increasing redness, swelling, warmth, drainage, or increasing pain
- Persistent bruising, lumps, or skin changes that aren't improving
- A rash near the site
- A reaction that worsens over time
- Concern that you applied or injected incorrectly, or that the medication or supplies were contaminated

Send a clear photo if appropriate. Do not inject into, or apply a patch/cream over, red, swollen, hot, painful, bruised, infected, broken, or abnormal skin.

Wellness or Systemic Changes

Contact KORB if you develop:

- Worsening fatigue, persistent brain fog, or significant sleep disruption
- New or unexplained systemic symptoms
- New swelling or fluid retention
- New numbness or tingling
- New weakness that is not severe or sudden
- New symptoms after starting a therapy

- Symptoms that make you unsure whether therapy remains appropriate

Some symptoms may need broader evaluation and shouldn't automatically be attributed to your KORB therapy.

Tell KORB Before Continuing Therapy If Something Changes

Contact KORB before continuing therapy if you develop:

- New pregnancy or breastfeeding status
- New cancer diagnosis or cancer treatment
- New serious infection or severe medical condition
- New kidney or liver problems, heart failure, or a major cardiovascular event
- New autoimmune, inflammatory, or significant GI disease
- A new medication that may interact with your current plan
- Upcoming surgery or imaging with contrast
- Restricted oral intake, significant dehydration, or acute illness

Your treatment plan may need to be held, adjusted, or reassessed.

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep your medication with you — in your carry-on or personal item, not checked baggage — in its original labeled packaging. It doesn't need to be packed in ice or kept refrigerated during travel unless your medication requires it; just keep it away from excessive heat (above 86°F / 30°C) and from freezing. Once you reach your destination, store it as directed again. If flying, medication and supplies are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. If you'll be off your usual schedule for several days, contact KORB for guidance.

When to Seek Emergency Care

Seek urgent or emergency care right away for symptoms that feel severe, unsafe, or rapidly worsening. Don't wait to hear back from KORB.

Seek Emergency Care Right Away For: trouble breathing or wheezing; swelling of the face, lips, tongue, or throat; severe rash or widespread hives; chest pain or fainting; severe dizziness or confusion; severe weakness; severe or rapidly worsening pain; new neurologic symptoms — weakness, numbness, confusion, trouble speaking, or loss of bladder/bowel control; severe headache with neurologic symptoms; severe abdominal pain or persistent vomiting; signs of severe dehydration; high fever or signs of serious infection; rapidly spreading redness, red streaking, severe warmth, or drainage from an injection or application site; severe glucose-related symptoms; or any symptom that feels urgent, severe, or unsafe.

Possible allergic reaction: trouble breathing, wheezing, or chest tightness; swelling of the face, lips, tongue, or throat; severe rash or widespread hives; fainting, severe dizziness, or a feeling like your throat is closing. Don't wait to hear back from KORB.

Possible serious infection: high fever; rapidly spreading redness or red streaking from an injection or application site; severe warmth, swelling, or pain at the site; drainage or pus with worsening symptoms; confusion or fainting. Contact KORB for mild or uncertain concerns, but seek urgent care if symptoms are severe or rapidly worsening.

Phone and email are not appropriate for emergencies. Call 911 or go to the nearest emergency room.

Key Reminders

- Contact KORB Operations for non-emergency logistics; use the Patient Portal to message your provider about medical concerns.
- Seek urgent or emergency care for severe, rapidly worsening, or unsafe symptoms — don't wait to hear back from KORB first.
- Phone and email are not appropriate for emergencies. Call 911 or go to the ER.
- Do not change your dose, timing, route, schedule, or medication combination without KORB provider guidance.
- Do not continue therapy if you're unsure it's safe — message your provider via the Patient Portal for guidance.
- Follow your prescription label and pharmacy-specific instructions at all times.

For anything in this guide that isn't a medical emergency and doesn't need your provider directly — medication questions, follow-up needs, or logistics like pharmacy and shipping — contact KORB Operations. If you reach voicemail, please leave one message; our staff will return your call within 24 hours on weekdays, excluding observed holidays.