

Wegovy

Brand-Name GLP-1 · Provider Reference

Weight Loss Program · KORB Health Group · korb-glp1-data.js v1.2 · generated 2026-08-09

Wegovy is written as a Tebra STANDARD prescription, never Tebra Compound. KORB does not manage fulfillment, does not complete prior authorizations, and does not handle coupons.

Available states

TX, CA, AL, WI

Brand-name GLP-1 through the manufacturer programs is only available in Texas, California, Alabama and Wisconsin. Source: the brand prescribing document. This is a program limit, not a shipping limit — do not offer a brand pathway to a patient outside these four states.

Rules and visit fee

Order via	Tebra Standard prescription (NOT Tebra Compound)
Billing	Patient pays the manufacturer program or the local pharmacy directly.
Prior authorizations	KORB does NOT complete insurance prior authorizations.
Coupons and savings cards	KORB does NOT handle coupons or savings cards.
Patient responsibility	KORB does not complete insurance prior authorizations and does not handle coupons or manufacturer savings cards. The patient is responsible for checking pricing, coverage and every other aspect of the medication.
Local pharmacy	A brand prescription can be sent to the patient's local pharmacy on request, under the same rules. KORB will not re-send a prescription between pharmacies to chase a lower price. The patient does that comparison before ordering.
Prescription visit fee	\$79 — charge code FITGLP1001
Fee covers	Cash-pay only. The fee covers the clinical visit, prescription issuance and program administration. Medication cost, pharmacy processing, shipping and supplies are set by the external pharmacy and paid directly by the patient.
Non-refundable	The visit fee is non-refundable once the visit has occurred or a prescription has been issued. Medication cost is not included.

Medication pricing is deliberately not listed

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Wegovy — Pen

Route: subcutaneous · **Frequency:** once weekly

Order via: Tebra Standard prescription

Pharmacy: NovoCare Pharmacy, 2400 Sand Lake Road, Suite 200B, Orlando, FL 32809 · (833) 949-5527

Wegovy pens are SINGLE-DOSE, so 4 pens covers 4 weeks. This is the opposite of the Zepbound KwikPen, where ONE pen holds all four weekly doses. The Tebra entries confirm it: each Wegovy line is one strength at one volume, with no total-content

figure in brackets.

Volume varies by dose. 0.25, 0.5 and 1 mg are 0.5 mL; 1.7, 2.4 and 7.2 mg are 0.75 mL. The 7.2 mg strength is listed as "Wegovy hd".

Dispensing option	Quantity	Refill	Days	When to use
4-week (28-day)	4	0	28	New prescription or any dose change. Use until the dose is stable.
8-week (56-day) — 4 pens + 1 refill	4	1	28	Clinically stable patients only.

Tebra STANDARD fields — 4-week (28-day)

Dose	Drug (dropdown)	Patient Instructions	Qty	RF	Days
0.25 mg	Wegovy 0.25 mg/0.5 ml subcutaneous pen injector	Inject 0.25 mg subcutaneously once weekly for 4 weeks.	4	0	28
0.5 mg	Wegovy 0.5 mg/0.5 ml subcutaneous pen injector	Inject 0.5 mg subcutaneously once weekly for 4 weeks.	4	0	28
1 mg	Wegovy 1 mg/0.5 ml subcutaneous pen injector	Inject 1 mg subcutaneously once weekly for 4 weeks.	4	0	28
1.7 mg	Wegovy 1.7 mg/0.75 ml subcutaneous pen injector	Inject 1.7 mg subcutaneously once weekly for 4 weeks.	4	0	28
2.4 mg	Wegovy 2.4 mg/0.75 ml subcutaneous pen injector	Inject 2.4 mg subcutaneously once weekly for 4 weeks.	4	0	28
7.2 mg [Wegovy HD]	Wegovy hd 7.2 mg/0.75 ml subcutaneous pen injector	Inject 7.2 mg subcutaneously once weekly for 4 weeks.	4	0	28

Tebra STANDARD fields — 8-week (56-day) — 4 pens + 1 refill

Dose	Drug (dropdown)	Patient Instructions	Qty	RF	Days
0.25 mg	Wegovy 0.25 mg/0.5 ml subcutaneous pen injector	Inject 0.25 mg subcutaneously once weekly for 4 weeks.	4	1	28
0.5 mg	Wegovy 0.5 mg/0.5 ml subcutaneous pen injector	Inject 0.5 mg subcutaneously once weekly for 4 weeks.	4	1	28
1 mg	Wegovy 1 mg/0.5 ml subcutaneous pen injector	Inject 1 mg subcutaneously once weekly for 4 weeks.	4	1	28
1.7 mg	Wegovy 1.7 mg/0.75 ml subcutaneous pen injector	Inject 1.7 mg subcutaneously once weekly for 4 weeks.	4	1	28
2.4 mg	Wegovy 2.4 mg/0.75 ml subcutaneous pen injector	Inject 2.4 mg subcutaneously once weekly for 4 weeks.	4	1	28
7.2 mg [Wegovy HD]	Wegovy hd 7.2 mg/0.75 ml subcutaneous pen injector	Inject 7.2 mg subcutaneously once weekly for 4 weeks.	4	1	28

- The 7.2 mg strength is marketed as Wegovy HD. Same molecule, higher strength.

- Dose escalation follows FDA-approved labeling and is individualized.

Wegovy — Oral tablet

Route: oral · **Frequency:** once daily

Order via: Tebra Standard prescription

Pharmacy: NovoCare Pharmacy, 2400 Sand Lake Road, Suite 200B, Orlando, FL 32809 · (833) 949-5527

Dispensing option	Quantity	Refill	Days	When to use
30-day supply	30	0	30	New prescription or any dose change. Use until the dose is stable.
60-day supply	60	0	60	Clinically stable patients only. Quantity 60, no refill — not 30 with a refill.

Tebra STANDARD fields — 30-day supply

Dose	Drug (dropdown)	Patient Instructions	Qty	RF	Days
1.5 mg	Wegovy 1.5 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	30	0	30
4 mg	Wegovy 4 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	30	0	30
9 mg	Wegovy 9 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	30	0	30
25 mg	Wegovy 25 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	30	0	30

Tebra STANDARD fields — 60-day supply

Dose	Drug (dropdown)	Patient Instructions	Qty	RF	Days
1.5 mg	Wegovy 1.5 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	60	0	60
4 mg	Wegovy 4 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	60	0	60
9 mg	Wegovy 9 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	60	0	60
25 mg	Wegovy 25 mg tablet	Take 1 tablet by mouth daily on empty stomach with up to 4 oz water. Swallow whole. No food, drink or other oral meds for 30 min.	60	0	60

- Sent electronically through Tebra as a STANDARD prescription. No insurance billing and no prior authorization.
- Patient completes enrolment, payment and shipping directly with NovoCare.
- Oral semaglutide has lower and more variable bioavailability than injectable. Adherence to the empty-stomach and 30-minute rules is critical to efficacy.
- Weight loss response may be less pronounced than with injectable GLP-1 therapy.

Is this the right patient

Program: **Get Fit Now** · Visit cadence: **Every 4 to 8 weeks**

Indications

- Weight loss
- Medically driven weight loss
- Increased energy
- Better sleep
- Improved mental health
- Improved cardiovascular health

Identifying the appropriate candidate

Body mass index	BMI greater than 25 is overweight · BMI greater than 30 is obese
FDA-approved uses	• Weight loss • Type 2 diabetes • Prevention of cardiovascular death, myocardial infarction and stroke in adults with established cardiovascular disease who are overweight or obese
Diabetic and pre-diabetic patients	KORB treats weight loss. Patients who are diabetic or pre-diabetic are acceptable. Diabetic patients can usually obtain coverage through insurance; where cost or availability prevents that, KORB can supply the medication. The patient must continue diabetes management with their PCP.

Clinical reference

Contraindications

• Personal or family history of medullary thyroid carcinoma (MTC)
• Multiple endocrine neoplasia syndrome type 2 (MEN2)
• Hypersensitivity to the active drug or any component of the formulation
• Current pregnancy, breastfeeding, or planning pregnancy
• Active gallbladder disease or history of gallbladder-related surgical complications
• History of pancreatitis (use with caution)
• Severe gastrointestinal disorders such as gastroparesis
• Severe renal impairment (eGFR below 30 mL/min/1.73 m ²)
• Uncontrolled diabetic retinopathy (primarily relevant in type 2 diabetes)

Side effects

Common	Less common
• Pain, redness or swelling at the injection site	• Pancreatitis
• Nausea and vomiting	• Gastroparesis
• Diarrhea	• Bowel obstruction
• Stomach pain	• Gallstone attacks
• Constipation	
• Low appetite	
• Headaches	
• Dizziness	

Counseling

• Gastrointestinal side effects are most common during dose escalation.
• Slower titration or holding at the current dose may improve tolerability.
• Instruct the patient to report persistent, severe or worsening symptoms.
• New or severe symptoms may require dose adjustment, temporary hold, or discontinuation.
• Vials are good for 28 days from first use regardless of the pharmacy BUD. Counsel the patient to discard at 28 days even if medication remains.

Questions and escalation

- Pharmacy or shipping issues — Operations
- Charge codes and billing — Operations
- Clinical protocol questions — Clinical Director
- Corrections to this document — Clinical Operations

Do not improvise

- Do not substitute a pharmacy the tool marks as unavailable
- Do not alter quantity, refill or days supply
- Do not paraphrase patient instructions
- Do not quote pricing without confirming with Operations