

Zepbound

Brand-Name GLP-1 · Provider Reference

Weight Loss Program · KORB Health Group · korb-glp1-data.js v1.2 · generated 2026-08-09

Zepbound is written as a Tebra STANDARD prescription, never Tebra Compound. KORB does not manage fulfillment, does not complete prior authorizations, and does not handle coupons.

Available states

TX, CA, AL, WI

Brand-name GLP-1 through the manufacturer programs is only available in Texas, California, Alabama and Wisconsin. Source: the brand prescribing document. This is a program limit, not a shipping limit — do not offer a brand pathway to a patient outside these four states.

Rules and visit fee

Order via	Tebra Standard prescription (NOT Tebra Compound)
Billing	Patient pays the manufacturer program or the local pharmacy directly.
Prior authorizations	KORB does NOT complete insurance prior authorizations.
Coupons and savings cards	KORB does NOT handle coupons or savings cards.
Patient responsibility	KORB does not complete insurance prior authorizations and does not handle coupons or manufacturer savings cards. The patient is responsible for checking pricing, coverage and every other aspect of the medication.
Local pharmacy	A brand prescription can be sent to the patient's local pharmacy on request, under the same rules. KORB will not re-send a prescription between pharmacies to chase a lower price. The patient does that comparison before ordering.
Prescription visit fee	\$79 — charge code FITGLP1001
Fee covers	Cash-pay only. The fee covers the clinical visit, prescription issuance and program administration. Medication cost, pharmacy processing, shipping and supplies are set by the external pharmacy and paid directly by the patient.
Non-refundable	The visit fee is non-refundable once the visit has occurred or a prescription has been issued. Medication cost is not included.

Medication pricing is deliberately not listed

Manufacturer pricing changes without notice and KORB does not control it. Storing it creates a maintenance burden and risks providers quoting incorrect pricing to patients. Price and coverage are the patient's responsibility.

Zepbound — KwikPen

Route: subcutaneous · **Frequency:** once weekly

Order via: Tebra Standard prescription

Pharmacy: LillyDirect Self Pay Pharmacy Solutions, 4343 Equity Dr, Columbus, OH 43228-3842 · (833) 432-4322

KwikPen only. Do NOT prescribe the single-dose vial presentation.

One KwikPen holds 4 fixed weekly doses. The pen locks out after the fourth dose and is then discarded, including any leftover

medicine. Do not transfer from the pen into a syringe.

Two presentations in the dropdown — pick carefully

Tebra lists BOTH presentations. The vial reads "Zepbound X mg/0.5 ml subcutaneous solution"; the pen reads "Zepbound kwikpen X mg/0.6 ml ...". Selecting the 0.5 mL vial with quantity 1 dispenses ONE WEEK instead of four. The bracketed figure in the pen entry is total pen content — 4 doses x 0.6 mL = 2.4 mL — which is Tebra confirming the 4-dose device.

Dispensing option	Quantity	Refill	Days	When to use
4-week (28-day)	1	0	28	New prescription or any dose change. Use until the dose is stable.
8-week (56-day) — 1 pen + 1 refill	1	1	28	Clinically stable patients only. Each fill is one pen covering 4 weeks.

Tebra STANDARD fields — 4-week (28-day)

Dose	Drug (dropdown)	Patient Instructions	Qty	RF	Days
2.5 mg	Zepbound kwikpen 2.5 mg/0.6 ml (10 mg/2.4 ml) subcutaneous pen injector	Inject 2.5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	0	28
5 mg	Zepbound kwikpen 5 mg/0.6 ml (20 mg/2.4 ml) subcutaneous pen injector	Inject 5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	0	28
7.5 mg	Zepbound kwikpen 7.5 mg/0.6 ml (30 mg/2.4 ml) subcutaneous pen injector	Inject 7.5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	0	28
10 mg	Zepbound kwikpen 10 mg/0.6 ml (40 mg/2.4 ml) subcutaneous pen injector	Inject 10 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	0	28
12.5 mg	Zepbound kwikpen 12.5 mg/0.6 ml (50 mg/2.4 ml) subcutaneous pen injector	Inject 12.5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	0	28
15 mg	Zepbound kwikpen 15 mg/0.6 ml (60 mg/2.4 ml) subcutaneous pen injector	Inject 15 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	0	28

Tebra STANDARD fields — 8-week (56-day) — 1 pen + 1 refill

Dose	Drug (dropdown)	Patient Instructions	Qty	RF	Days
2.5 mg	Zepbound kwikpen 2.5 mg/0.6 ml (10 mg/2.4 ml) subcutaneous pen injector	Inject 2.5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	1	28
5 mg	Zepbound kwikpen 5 mg/0.6 ml (20 mg/2.4 ml) subcutaneous pen injector	Inject 5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	1	28
7.5 mg	Zepbound kwikpen 7.5 mg/0.6 ml (30 mg/2.4 ml) subcutaneous pen injector	Inject 7.5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	1	28
10 mg	Zepbound kwikpen 10 mg/0.6 ml (40 mg/2.4 ml) subcutaneous pen injector	Inject 10 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	1	28
12.5 mg	Zepbound kwikpen 12.5 mg/0.6 ml (50 mg/2.4 ml) subcutaneous pen injector	Inject 12.5 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	1	28
15 mg	Zepbound kwikpen 15 mg/0.6 ml (60 mg/2.4 ml) subcutaneous pen injector	Inject 15 mg subcutaneously once weekly for 4 weeks. Multi-dose pen — discard after 4 doses.	1	1	28

- Sent electronically through Tebra as a STANDARD prescription. No insurance billing and no prior authorization.
- Patient completes enrolment, payment and shipping directly with LillyDirect.

- PEN NEEDLES ARE REQUIRED and are not included. The patient purchases them, and they can be added at LillyDirect checkout. Each injection uses a new needle.
- Zepbound is sold in three formats. KORB uses the KwikPen only.
- Rotate injection sites. Follow standard tirzepatide titration principles.
- Early refills or replacement shipments are decided by the pharmacy, not KORB.

Is this the right patient

Program: **Get Fit Now** · Visit cadence: **Every 4 to 8 weeks**

Indications

- Weight loss
- Medically driven weight loss
- Increased energy
- Better sleep
- Improved mental health
- Improved cardiovascular health

Identifying the appropriate candidate

Body mass index	BMI greater than 25 is overweight · BMI greater than 30 is obese
FDA-approved uses	• Weight loss • Type 2 diabetes • Prevention of cardiovascular death, myocardial infarction and stroke in adults with established cardiovascular disease who are overweight or obese
Diabetic and pre-diabetic patients	KORB treats weight loss. Patients who are diabetic or pre-diabetic are acceptable. Diabetic patients can usually obtain coverage through insurance; where cost or availability prevents that, KORB can supply the medication. The patient must continue diabetes management with their PCP.

Clinical reference

Contraindications

- Personal or family history of medullary thyroid carcinoma (MTC)
- Multiple endocrine neoplasia syndrome type 2 (MEN2)
- Hypersensitivity to the active drug or any component of the formulation
- Current pregnancy, breastfeeding, or planning pregnancy
- Active gallbladder disease or history of gallbladder-related surgical complications
- History of pancreatitis (use with caution)
- Severe gastrointestinal disorders such as gastroparesis
- Severe renal impairment (eGFR below 30 mL/min/1.73 m²)
- Uncontrolled diabetic retinopathy (primarily relevant in type 2 diabetes)

Side effects

Common	Less common
• Pain, redness or swelling at the injection site	• Pancreatitis
• Nausea and vomiting	• Gastroparesis

Common	Less common
• Diarrhea	• Bowel obstruction
• Stomach pain	• Gallstone attacks
• Constipation	
• Low appetite	
• Headaches	
• Dizziness	

Counseling

• Gastrointestinal side effects are most common during dose escalation.
• Slower titration or holding at the current dose may improve tolerability.
• Instruct the patient to report persistent, severe or worsening symptoms.
• New or severe symptoms may require dose adjustment, temporary hold, or discontinuation.
• Vials are good for 28 days from first use regardless of the pharmacy BUD. Counsel the patient to discard at 28 days even if medication remains.

Questions and escalation • Pharmacy or shipping issues — Operations • Charge codes and billing — Operations • Clinical protocol questions — Clinical Director • Corrections to this document — Clinical Operations	Do not improvise • Do not substitute a pharmacy the tool marks as unavailable • Do not alter quantity, refill or days supply • Do not paraphrase patient instructions • Do not quote pricing without confirming with Operations
--	---