

Farmakeio — Tirzepatide

GLP-1 Provider Reference

Weight Loss Program · KORB Health Group · korb-glp1-data.js v1.2 · generated 2026-08-09

Everything needed to prescribe Farmakeio tirzepatide, complete on its own. Values are copied literally into the Tebra Compound section — do not paraphrase, and do not adjust quantity, refill or days supply.

Farmakeio Pharmacy at a glance

Status	Active
Preferred states	AK, AL, AR, CO, CT, DC, DE, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NY, OH, OK, OR, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY
Ships to	AK, AL, AR, AZ, CO, CT, DC, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, WY
CANNOT ship to	CA
Order via	Tebra Compound
Billing	Bill to KORB Health Group, ship to patient
Dispensing address	Farmakeio Pharmacy, 1736 N Greenville Ave, Richardson, TX 75081

• Ships to 49 states plus DC. California is the single state Farmakeio will not ship to, and that cannot be overridden. Preferred everywhere except CA, TX, FL, AZ and NV.

Before you prescribe

- Products ship as a HOME KIT that includes syringes and supplies.
- Compounded with pyridoxine (B-6) for nausea prevention, not B-12.
- Semaglutide is a single concentration (2.5 mg/25 mg per mL). Dose is set by volume; the only variable on the Rx is how many mL.
- Tirzepatide is a single concentration (18 mg/25 mg per mL).

Farmakeio — Tirzepatide / B-6 Home Kit

Formulation: Tirzepatide 18 mg / Pyridoxine (B-6) 25 mg per mL — HOME KIT

Single concentration for every dose. Dose is set by volume.

Route: subcutaneous · **Frequency:** once weekly · **Order via:** Tebra Compound

Same for every dose

Drug Formulation	Tirzepatide/B-6 18mg/25mg per mL Home Kit
Select	Allow Substitution
Unit	ml
Reason for Compounding	N/V mitigation & flexibility
Pharmacy Instructions	Bill to KORB Health Group and ship to the patient. Home Kit. Custom Rx for N/V mitigation, dosing flexibility, and added B-6.

4-week program

Dose	Name	Patient Instructions	Qty	RF	Days
2.5 mg	FARMAKEIO – Tirzepatide 2.5 mg – 4-Week Supply	INJECT 0.13ML (13 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	1	0	28
4.5 mg	FARMAKEIO – Tirzepatide 4.5 mg – 4-Week Supply	INJECT 0.25ML (25 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	1	0	28
7.5 mg	FARMAKEIO – Tirzepatide 7.5 mg – 4-Week Supply	INJECT 0.42ML (42 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	2	0	28
9 mg	FARMAKEIO – Tirzepatide 9 mg – 4-Week Supply	INJECT 0.50ML (50 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	2	0	28
13.5 mg	FARMAKEIO – Tirzepatide 13.5 mg – 4-Week Supply	INJECT 0.75ML (75 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	3	0	28
15 mg	FARMAKEIO – Tirzepatide 15 mg – 4-Week Supply	INJECT 0.83ML (83 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	4	0	28

8-week program

Dose	Name	Patient Instructions	Qty	RF	Days
2.5 mg	FARMAKEIO – Tirzepatide 2.5 mg – 8-Week Supply	INJECT 0.13ML (13 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	2	0	56
4.5 mg	FARMAKEIO – Tirzepatide 4.5 mg – 8-Week Supply	INJECT 0.25ML (25 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	2	0	56
7.5 mg	FARMAKEIO – Tirzepatide 7.5 mg – 8-Week Supply	INJECT 0.42ML (42 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	4	0	56
9 mg	FARMAKEIO – Tirzepatide 9 mg – 8-Week Supply	INJECT 0.50ML (50 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	4	0	56
13.5 mg	FARMAKEIO – Tirzepatide 13.5 mg – 8-Week Supply	INJECT 0.75ML (75 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	6	0	56
15 mg	FARMAKEIO – Tirzepatide 15 mg – 8-Week Supply	INJECT 0.83ML (83 UNITS) SUBCUTANEOUSLY ONCE WEEKLY	8	0	56

Vials dispensed

Dose	Units	Concentration	4-week	8-week
2.5 mg	13 u	—	1 ml	1 ml x 2
4.5 mg	25 u	—	1 ml	1 ml x 2
7.5 mg	42 u	—	2 ml	2 ml x 2
9 mg	50 u	—	2 ml	2 ml x 2
13.5 mg	75 u	—	3 ml	3 ml x 2
15 mg	83 u	—	4 ml	4 ml x 2

Pricing and charge codes

Provider and internal only. Never quote a price to a patient from this document without confirming with Operations. Includes: Telehealth visit, medication, shipping and supplies.

Tirzepatide — by dose tier

Tier	Doses	4-week (corp partner)	4-week (standard)	8-week
T1A	2 / 2.5 / 4 / 4.5 / 5 mg depending on pharmacy	\$349 FITTirzCP2	\$349 FITTirz002	\$599 FITTirzMT1
T2A	6.5 / 7.5 / 8.5 / 9 / 10 mg depending on pharmacy	\$399 FITTirzCP3	\$399 FITTirz003	\$649 FITTirzMT2

Tier	Doses	4-week (corp partner)	4-week (standard)	8-week
T3A	12.5 / 13.5 / 15 / 16 mg depending on pharmacy	\$449 FITTirzCP4	\$449 FITTirz004	\$799 FITTirzMT3

- Tirzepatide has no corporate discount. The corp partner code exists for attribution and tracking, not a lower price — it is the same figure as the standard 4-week at every tier. Do not present it to a corporate partner as a saving.

If the patient moves state

Patient state	Pharmacy
TX, NV, AZ, FL	Premier Pharmacy
CA	Belmar Pharmacy
Everywhere else	Farmakeio Pharmacy ← this document

- Farmakeio Pharmacy cannot ship to: CA. Those states must move to another pharmacy.
- Established patients normally stay with their current pharmacy unless there is a clinical or supply reason to move, or the new state is one this pharmacy cannot serve.

Is this the right patient

Program: **Get Fit Now** · Visit cadence: **Every 4 to 8 weeks**

Indications

• Weight loss
• Medically driven weight loss
• Increased energy
• Better sleep
• Improved mental health
• Improved cardiovascular health

Identifying the appropriate candidate

Body mass index	BMI greater than 25 is overweight · BMI greater than 30 is obese
FDA-approved uses	• Weight loss • Type 2 diabetes • Prevention of cardiovascular death, myocardial infarction and stroke in adults with established cardiovascular disease who are overweight or obese
Diabetic and pre-diabetic patients	KORB treats weight loss. Patients who are diabetic or pre-diabetic are acceptable. Diabetic patients can usually obtain coverage through insurance; where cost or availability prevents that, KORB can supply the medication. The patient must continue diabetes management with their PCP.

Program length and follow-up

INTERNAL ONLY. Internal and provider-facing only. Do not put this rule in a patient handout, the website, a pricing flyer, or any Circle post. A patient should not arrive expecting to graduate to 8-week.

Start rule	Every patient starts on the 4-week program. No exceptions.
Move to 8-week	A patient may be offered the 8-week program once they are stable at the dose they have landed on after escalation.
Who decides	Provider. Provider-driven throughout. The 8-week program is an offer the provider extends, not a milestone the patient reaches or requests. Escalation, holding, reducing and weaning off are all provider discretion.

Visit cadence	4-week: Monthly provider visit. 8-week: Provider visit every 8 weeks. Either a video visit or an asynchronous visit. Both are acceptable for 4-week and 8-week follow-ups.
Weaning	Dose reduction and weaning off therapy are entirely at provider discretion. No fixed taper schedule is defined at the program level.
Reasons to stay at 4-week	• Cost — the 4-week program is the lower per-visit outlay and carries the discounted rate. • Accountability — some patients want to see the provider monthly. • Clinical — the provider judges that side effects are not adequately controlled and the patient needs to be seen more often.

Clinical reference

Contraindications

• Personal or family history of medullary thyroid carcinoma (MTC)
• Multiple endocrine neoplasia syndrome type 2 (MEN2)
• Hypersensitivity to the active drug or any component of the formulation
• Current pregnancy, breastfeeding, or planning pregnancy
• Active gallbladder disease or history of gallbladder-related surgical complications
• History of pancreatitis (use with caution)
• Severe gastrointestinal disorders such as gastroparesis
• Severe renal impairment (eGFR below 30 mL/min/1.73 m ²)
• Uncontrolled diabetic retinopathy (primarily relevant in type 2 diabetes)

Side effects

Common	Less common
• Pain, redness or swelling at the injection site	• Pancreatitis
• Nausea and vomiting	• Gastroparesis
• Diarrhea	• Bowel obstruction
• Stomach pain	• Gallstone attacks
• Constipation	
• Low appetite	
• Headaches	
• Dizziness	

Counseling

• Gastrointestinal side effects are most common during dose escalation.
• Slower titration or holding at the current dose may improve tolerability.
• Instruct the patient to report persistent, severe or worsening symptoms.
• New or severe symptoms may require dose adjustment, temporary hold, or discontinuation.
• Vials are good for 28 days from first use regardless of the pharmacy BUD. Counsel the patient to discard at 28 days even if medication remains.

Questions and escalation

- Pharmacy or shipping issues — Operations
- Charge codes and billing — Operations
- Clinical protocol questions — Clinical Director
- Corrections to this document — Clinical Operations

Do not improvise

- Do not substitute a pharmacy the tool marks as unavailable
- Do not alter quantity, refill or days supply
- Do not paraphrase patient instructions
- Do not quote pricing without confirming with Operations