

Untitled

PHS 7010 Fall 2023 - Final Report

Kline Dubose, Haojia Li

April 18, 2024

1 Likelihood ratio tests in linear mixed models with one variance component

Consider an LMM with one variance component

$$\mathbf{Y} = X\beta + Z\mathbf{b} + \epsilon, \quad E \begin{pmatrix} \mathbf{b} \\ \epsilon \end{pmatrix} = \begin{pmatrix} \mathbf{0}_K \\ \mathbf{0}_n \end{pmatrix}, \quad \text{cov} \begin{pmatrix} \mathbf{b} \\ \epsilon \end{pmatrix} = \begin{pmatrix} \sigma_b^2 \Sigma & \mathbf{0} \\ \mathbf{0} & \sigma_\epsilon^2 I_n \end{pmatrix}$$

where,

- \mathbf{Y} is the $n \times 1$ vector of observations,
- X is the $n \times p$ design matrix for the fixed effects,
- Z is the $n \times K$ design matrix for the random effects,
- β is a p -dimensional vector of fixed effects parameters,
- \mathbf{b} is a K -dimensional vector of random effects,
- (\mathbf{b}, ϵ) has a normal distribution.

Under these conditions it follows that

$$E(\mathbf{Y}) = X\beta, \\ \text{var}(\mathbf{Y}) = \sigma_\epsilon^2 V_\lambda$$

where

- $\lambda = \sigma_b^2 / \sigma_\epsilon^2$, which can be considered a signal-to-noise ratio,
- $V_\lambda = I_n + \lambda Z \Sigma Z'$.