Introduction to Matrices

Revise Matrices unit from G.C.E. (A/L) Combined Mathematics and G.C.E. (O/L) Mathematics.

Types of matrices

Square matrix

Number of columns equal to number of rows.

(i) Main diagonals of a square matrix

Formed by elements having equal subscripts.

Diagonal matrix

A square matrix whose only non-zero elements are main-diagonal elements. Denoted by $m{D}$. Subset of triangular matrices.

Identity matrix or Unit matrix

A diagonal matrix whose diagonal elements are all equal to ${f 1}$. Denoted by ${f I}$. Subset of diagonal matrices.

Zero matrix / Null matrix

All elements are 0.

Column matrix (column vector)

Only ${f 1}$ column.

Row matrix (row vector)

Only ${f 1}$ row.

Triangular matrix

Upper triangular matrix or lower triangular matrix.

Upper triangular matrix

All elements below the main diagonal are $\mathbf{0}$. Subset of square matrices.

Lower triangular matrix

All elements above the main diagonal are $\mathbf{0}$. Subset of square matrices.

Matrix operations

Addition and subtraction

Order of the 2 matrices must be same. Matrix obtained by adding or subtracting corresponding elements.

Scalar multiplication

Matrix obtained by multiplying all elements by the scalar.



Matrix multiplication is also defined.

Transpose

Matrix obtained from a given matrix by interchanging its rows and columns. Denoted by a superscript T, like A^T .

Properties

- $_{1.}(A^T)^T=A$
- 2. Distributive over addition:

$$(A+B)^T = A^T + B^T$$

- $(A+B)^T = A^T + B^T$ 3. $(kA)^T = kA^T$ 4. $(A imes B)^T = B^T imes A^T$

More types of matrices

Symmetric matrix

If $A=A^T$. Subset of square matrices.

Skew-symmetric matrix

If $A=-A^T$. Subset of square matrices. All elements in main diagonal are 0.

(i) Note

Any square matrix can be expressed as a sum of a symmetric matrix and a skewsymmetric matrix.

Matrix multiplication

Defined only if the number of columns of the first matrix is equal to the number of rows of the second matrix.

If
$$A=(a_{ij})_{m imes p}$$
 and $B=(b_{ij})_{p imes n}$, then $A imes B=C=(c_{ij})_{m imes n}$ where $c_{ij}=a_{i1}b_{1j}+a_{i2}b_{2j}+\cdots+a_{ip}b_{pj}$.

(i) Note

$$A \times B \neq B \times A$$

• Generally $A\times B\neq B\times A$. . $A\times B=0 \Longrightarrow A=0 \vee B=0$. $A\neq 0 \wedge B\neq 0 \Longrightarrow A\times B\neq 0$

$$A \neq 0 \land B \neq 0 \Longrightarrow A \times B \neq 0$$

Properties of matrix multiplication

A,B,C,I matrices must be chosen so that below-mentioned product matrices are defined.

1. Associative:

$$A(BC) = (AB)C$$

2. Right distributive over addition:

$$(A+B)C = AC + BC$$

3. Left distributive over addition:

$$C(A+B) = CA + CB$$

4. $\overrightarrow{AI} = \overrightarrow{IA} = \overrightarrow{A}$

is an identity matrix.

Determinant

Defined only for square matrices. Denoted by |A|.

For 2x2

$$|A| = egin{array}{c|c} a_{11} & a_{12} \ a_{21} & a_{22} \ \end{array} = a_{11}a_{22} - a_{12}a_{21}$$

For higher order

Minor of an element

Suppose $A=(a_{ij})$.

Minor of an element a_{ij} , is the matrix obtained by deleting $i^{
m th}$ row and $j^{
m th}$ column of A. Denoted by M_{ij} .

Co-factor of an element

Suppose $A=(a_{ij})$.

Co-factor of an element a_{ij} , is defined as (commonly denoted as A_{ij}):

$$A_{ij} = (-1)^{i+j} \, |M_{ij}|$$

Definition

If $A=(a_{ij})_{n imes n}$ then the ${f determinant}$ of A is denoted by |A| and is defined by:

$$|A| = \sum_{j=1}^n a_{ij} A_{ij}$$

where $1 \leq j \leq n$.

Properties of determinants

- Every element of a row or column of a matrix is ${\bf 0}$ then the value of its determinant is ${\bf 0}$
- If 2 columns or 2 rows of a matrix are identical then its determinant is $\boldsymbol{0}$

• If A and B are two square matrices then

$$|AB| = |A||B|$$

- The value of the determinant of a matrix remains unchanged if a scalar multiple of a row or column is added to any other row or column.
- If a matrix

 \boldsymbol{B}

is obtained from a square matrix

 \boldsymbol{A}

by an interchange of two columns or rows:

$$|B| = -|A|$$

• If every entry in any row or column is multiplied by

k

, then the whole determinant is multiplied by

 \boldsymbol{k}

.

Adjoint

Suppose $A=(a_{ij})_{n imes n}$.

$$\mathrm{adj}A = (A_{ij})_{n \times n}^T$$

Where A_{ij} is the <u>co-factor of</u> a_{ij} .

Inverse

Suppose A and B are square matrices of the same order. If AB=BA=I then B is called the inverse of \boldsymbol{A} and is denoted by \boldsymbol{A}^{-1} .

$$A^{-1} = \frac{\operatorname{adj} A}{|A|}$$

 \odot Singular vs Non-singular A square matrix is singular if |A|=0. Otherwise non-singular or invertible matrices.

Properties of Inverse

- . $(AB)^{-1} = B^{-1}A^{-1}$. $(A^T)^{-1} = (A^{-1})^T$. $A \operatorname{adj} A = \operatorname{adj} A A = |A|I$

Elementary Transformations

- · Interchange of any columns or rows
- · Addition of multiple of any row or column to any other row or column
- · Multiplication of each element of a column or a row by a non-zero constant

When a matrix B is obtained by applying elementary transformations to a matrix A, then A is equivalent to B. Denoted by A pprox B.

Theorem

The elementary row operations that reduce a given matrix A to the identity matrix, also transform the identity matrix to the inverse of A.

Augmented Matrix

Two matrices are written as a single matrix with a vertical line in-between. Denoted by (Aert B) . Example:

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

Inverse using elementary row transformations

Let A be a square matrix with order $n \times n$.

- Start with $(A_{n imes n} | I_n)$
- Repeatedly add ${f row}$ transformations (not column) to both of the matrices until the LHS becomes an identity matrix.

 \circ Convert all elements outside the main diagonal to $oldsymbol{\Omega}$

 \circ Convert elements on the main diagonal to $\ensuremath{\mathbf{1}}$ by multiplying by a constant.

• When LHS is an identity matrix, RHS is A^{-1}



What about singular matrices?

System of Linear Equations

Any system of linear equations can be represented in matrix notation as shown below.

- $a_{11}x + a_{12}y + a_{13}z = b_1$
- $a_{21}x + a_{22}y + a_{23}z = b_2$
- $a_{31}x + a_{32}y + a_{33}z = b_3$

$$egin{pmatrix} a_{11} & a_{12} & a_{13} \ a_{21} & a_{22} & a_{23} \ a_{31} & a_{32} & a_{33} \ \end{pmatrix} egin{pmatrix} x \ y \ z \ \end{pmatrix} = egin{pmatrix} b_1 \ b_2 \ b_3 \ \end{pmatrix} \implies AX = B$$

2 types based on $oldsymbol{B}$:

- $\cdot = 0$
 - : Homogeneous
- $. \neq 0$
 - : Non-homogeneous

Solution of non-homogenous systems

Method 1: Direct approach

Used when coefficient matrix $oldsymbol{A}$ is invertible. It means the system has a unique set of solutions.

$$AX = B \implies X = A^{-1}B$$

Method 2: Cramer's Rule

Let AX=B, where A is the coefficient matrix and $X=(x_i)_{n imes 1}$.

$$x_i = rac{|A_i|}{|A|}$$

Where A_i is the matrix obtained by replacing ith column in matrix A by B.

Method 1: Reducing to Echelon Form

Start with (A|B). Convert the LHS to echelon form using elementary row transformations. The solution can be found now. If a contradiction is encountered while solving the equation, that means the system has no solutions.

Echelon Form

A matrix is in echelon form iff:

- All rows having only zero entries are at the bottom.
- For all row that does not contain entirely zeros, the first non-zero entry is 1.
- For 2 successive non-zero rows, the leading 1 in the higher row is further left than the leading 1 in the lower row.

The process of reducing the augmented matrix to row Echelon form is known as **Gaussian elimination**.

Solution of homogenous systems

TODO