

Package ‘truncnorm’

July 2, 2014

Version 1.0-7

Title Truncated normal distribution

Description r/d/p/q functions for the truncated normal distribution

Author Heike Trautmann <trautmann@statistik.uni-dortmund.de>
and Detlef Steuer <detlef.steuer@hsu-hamburg.de>
and Olaf Mersmann <olafm@statistik.uni-dortmund.de>
and Björn Bornkamp <bornkamp@statistik.tu-dortmund.de>

Maintainer Olaf Mersmann <olafm@statistik.uni-dortmund.de>

Depends R (>= 2.15.0)

License GPL-2

Encoding UTF-8

Date

NeedsCompilation yes

Repository CRAN

Date/Publication 2014-01-21 10:17:26

R topics documented:

truncnorm	2
Index	4

truncnorm

*The Truncated Normal Distribution***Description**

Density, distribution function, quantile function, random generation and expected value function for the truncated normal distribution with mean equal to 'mean' and standard deviation equal to 'sd'.

Usage

```
dtruncnorm(x, a=-Inf, b=Inf, mean = 0, sd = 1)
ptruncnorm(q, a=-Inf, b=Inf, mean = 0, sd = 1)
qtruncnorm(p, a=-Inf, b=Inf, mean = 0, sd = 1)
rtruncnorm(n, a=-Inf, b=Inf, mean = 0, sd = 1)
etruncnorm(a=-Inf, b=Inf, mean=0, sd=1)
vtruncnorm(a=-Inf, b=Inf, mean=0, sd=1)
```

Arguments

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If 'length(n) > 1', the length is taken to be the number required.
a	vector of lower bounds. These may be -Inf
b	vector of upper bounds. These may be Inf
mean	vector of means.
sd	vector of standard deviations.

Details

If mean or sd are not specified they assume the default values of 0 and 1, respectively. The values of a, b, mean and sd are recycled as needed.

Value

'dtruncnorm' gives the density, 'ptruncnorm' gives the distribution function, 'qtruncnorm' gives the quantile function, 'rtruncnorm' generates random deviates, 'etruncnorm' gives the expected value and 'vtruncnorm' the variance of the distribution.

Author(s)

Heike Trautmann <trautmann@statistik.tu-dortmund.de>, Detlef Steuer <steuer@hsu-hamburg.de>, Olaf Mersmann <olafm@statistik.tu-dortmund.de> and Björn Bornkamp <bornkamp@statistik.tu-dortmund.de> who donated a much improved rtruncnorm implementation using an accept-reject sampler.

References

The accept-reject sampler follows the description given in

Geweke, J. (1991). *Efficient simulation from the multivariate normal and student-t distributions subject to linear constraints*. In Computing Science and Statistics: Proceedings of the 23rd Symposium on the Interface, Ed. E. Keramidas and S. Kaufman, pp. 571-8. Fairfax Station, VA: Interface Foundation of North America.

Index

*Topic **distribution**

truncnorm, [2](#)

dtruncnorm (truncnorm), [2](#)

etruncnorm (truncnorm), [2](#)

ptruncnorm (truncnorm), [2](#)

qtruncnorm (truncnorm), [2](#)

rtruncnorm (truncnorm), [2](#)

truncnorm, [2](#)

vtruncnorm (truncnorm), [2](#)