RESEARCH BRIEF / OCTOBER 2026

When are knee pain predictions less reliable?

Sricharan Karthigeyan and Yuheng Tang | Ardrey Kell High School

THE QUESTION

Does the mismatch between X-ray severity and reported pain identify knees with less accurate predictions four years later?

8,232
knees / 4,146 participants

1.787
held-out MAE

48 months
prediction horizon

WHAT WE DID

We analyzed existing Osteoarthritis Initiative data. Elastic-net models predicted WOMAC pain (0-20). Nested five-fold validation grouped participants, keeping both knees together and learning preprocessing inside training data. Prediction error was evaluated only on held-out cases.

THE RESULT - AND THE CRITICAL QUALIFICATION

The frozen unadjusted analysis linked each extra discordance point to 0.301 points of absolute error (95% CI 0.275-0.326). But discordance correlated strongly with baseline pain ($r = 0.751$). In a post-freeze sensitivity analysis, adjusting for baseline pain reduced the coefficient to -0.011 (95% CI -0.043 to 0.022; $p = 0.52$). Discordance did not retain an independent association in these data.

WHAT SELECTIVE DEFERRAL ADDS

Deferring the 823 most uncertain knees retained 7,409 cases and lowered their MAE from 1.787 to 1.582 (11.47% relative reduction). This trades coverage for lower retained error. The uncertainty model included discordance, so it is not independent evidence of a discordance effect. Deferral also varied by subgroup.

TAKEAWAY

Baseline symptom burden better explains the observed association than discordance alone. The study shows why prediction reliability needs careful testing, adjustment, and external validation.

LIMITATIONS & STATUS

One cohort; internal validation; 7.7% missing 48-month outcomes; post hoc baseline-pain adjustment; nonmonotonic quartile means. Not a clinical tool. No journal acceptance or external replication is claimed in this brief.

Explore the study and source materials

ksricharan7.github.io/oai-model-reliability/