

## **STATISTICS WORKSHEET-5**

**Q1 to Q10 are MCQs with only one correct answer. Choose the correct option.**

1. Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.
- a) Mean
  - b) Actual
  - c) Predicted
  - d) Expected

Answer:-(d) **Expected**

2. Chisquare is used to analyse
- a) Score
  - b) Rank
  - c) Frequencies
  - d) All of these

Answer:-(d) **All of these**

3. What is the mean of a Chi Square distribution with 6 degrees of freedom?
- a) 4
  - b) 12
  - c) 6
  - d) 8

Answer:-(c) **6**

4. Which of these distributions is used for a goodness of fit testing?
- a) Normal distribution
  - b) Chisquared distribution
  - c) Gamma distribution
  - d) Poisson distribution

Answer:-(d) **Poisson distribution**

5. Which of the following distributions is Continuous
- a) Binomial Distribution
  - b) Hypergeometric Distribution
  - c) F Distribution
  - d) Poisson Distribution

Answer:-(a) **Binomial distribution**

6. A statement made about a population for testing purpose is called?
- a) Statistic
  - b) Hypothesis
  - c) Level of Significance
  - d) TestStatistic

Answer:-(b) **Hypothesis**

7. If the assumed hypothesis is tested for rejection considering it to be true is called?
- a) Null Hypothesis
  - b) Statistical Hypothesis
  - c) Simple Hypothesis
  - d) Composite Hypothesis

Answer:-(a) **Null Hypothesis**

8. If the Critical region is evenly distributed then the test is referred as?
- a) Two tailed
  - b) One tailed
  - c) Three tailed
  - d) Zero tailed

Answer:-(a) **Two tailed**

9. Alternative Hypothesis is also called as?
- a) Composite hypothesis
  - b) Research Hypothesis
  - c) Simple Hypothesis
  - d) Null Hypothesis

Answer:-(b) **Research Hypothesis**

10. In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by \_\_\_\_\_
- a) np
  - b) n

Answer:-(a) **np**

---