

Assignment 15

KUSUMA PRIYA
EE20MTECH11007

Download codes from

Rewriting (1.0.1)

<https://github.com/KUSUMAPRIYAPULAVARTY/assignment15>

$$T_2 T(z) = \begin{pmatrix} x+7y \\ 5y \\ -10y \\ x-7y \end{pmatrix} \quad (2.0.4)$$

$$= \begin{pmatrix} 1 & 7 \\ 0 & 5 \\ 0 & -10 \\ 1 & -7 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (2.0.5)$$

$$T_2 T : \mathbf{V} \rightarrow R^4 \quad (2.0.6)$$

$$rref \begin{pmatrix} 1 & 0 & 0 & 1 \\ 7 & 5 & -10 & -7 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & -\frac{14}{5} \end{pmatrix} \quad (2.0.7)$$

1 QUESTION

Let \mathbf{V} be the set of complex numbers regarded as a vector space over the field of real numbers. We define a function T from \mathbf{V} into the space of 2×2 real matrices, as follows. If $z = x + iy$ with x and y real numbers, then

$$T(z) = \begin{pmatrix} x+7y & 5y \\ -10y & x-7y \end{pmatrix} \quad (1.0.1)$$

How would you describe the range of T ?

So, the vectors $\begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 7 \\ 5 \\ -10 \\ -7 \end{pmatrix}$ are linearly independent.

From, (2.0.5)

$$\text{range}(T_2 T) = \text{columnspace of } \begin{pmatrix} 1 & 7 \\ 0 & 5 \\ 0 & -10 \\ 1 & -7 \end{pmatrix}$$

$$\text{range}(T) \text{ is span of } \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 7 & 5 \\ -10 & -7 \end{pmatrix} \right\}$$

2 SOLUTION

$$T : \mathbf{V} \rightarrow R^{2 \times 2} \quad (2.0.1)$$

where $R^{2 \times 2}$ is the space of all 2×2 real matrices
If R^4 is the space of all 4-tuple vectors, then $R^{2 \times 2}$ and R^4 are isomorphic with

$$T_2 \left[\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right] = \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} \quad (2.0.2)$$

$$T_2 : R^{2 \times 2} \rightarrow R^4 \quad (2.0.3)$$