

Assignment 19

KUSUMA PRIYA
EE20MTECH11007

Download codes from

<https://github.com/KUSUMAPRIYAPULAVARTY/assignment19>

1 QUESTION

Let \mathbf{V} be the vector space of all polynomial functions over the field of real numbers. Let a and b be fixed real numbers and let f be the linear functional on \mathbf{V} defined by

$$f(p) = \int_a^b p(x) dx \quad (1.0.1)$$

If D is the differentiator operator on \mathbf{V} , what is $D^t f$?

2 SOLUTION

PARAMETERS	DESCRIPTION
\mathbb{R}	Field of real numbers
\mathbf{V}	Vector space of all polynomials over \mathbb{R}
a, b	Fixed real numbers
f	Linear functional on \mathbf{V}
D	Differentiator operator on \mathbf{V}
D^t	Transpose of D

TABLE 0: Input Parameters

STATEMENTS	DERIVATIONS
D^t is transpose of D	$(D^t f)(p) = f[D(p)]$ (2.0.1)
A polynomial of degree $n + 1$	$p = \mathbf{c}^T \mathbf{x}$ (2.0.2)
$p(x) = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n$	$\mathbf{c} = \begin{pmatrix} c_0 \\ c_1 \\ c_2 \\ \vdots \\ c_n \end{pmatrix}, \mathbf{x} = \begin{pmatrix} 1 \\ x \\ x^2 \\ \vdots \\ x^n \end{pmatrix}$ (2.0.3)
$f(p) = \int_a^b p(x) dx$	$f(p) = \mathbf{c}^T \mathbf{F}$ (2.0.4)
	$\mathbf{F} = \begin{pmatrix} b-a \\ \frac{b^2-a^2}{2} \\ \frac{b^3-a^3}{3} \\ \vdots \\ \frac{b^{n+1}-a^{n+1}}{n+1} \end{pmatrix}$ (2.0.5)
$D(p) = c_1 + 2c_2 x + 3c_3 x^2 + \dots + nc_n x^{n-1}$	$D(p) = \mathbf{c}^T \mathbf{D} \mathbf{x}$ (2.0.6)
	$\mathbf{D} = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 2 & 0 & \dots & 0 & 0 \\ 0 & 0 & 3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & n & 0 \end{pmatrix}$ (2.0.7)
$f[D(p)] = c_1(b-a) + c_2(b^2-a^2) + \dots + c_n(b^n-a^n)$	$\text{Let } D(p) = p'$ (2.0.8)
	$\Rightarrow p' = \mathbf{c}'^T \mathbf{x}$ (2.0.9)
	$\text{where } \mathbf{c}'^T = \begin{pmatrix} c_1 \\ 2c_2 \\ 3c_3 \\ \vdots \\ nc_n \\ 0 \end{pmatrix}$ (2.0.10)
	$f[D(p)] = \mathbf{c}'^T \mathbf{F}$ (2.0.11)
$(D^t f)(p) = p(b) - p(a)$	$(D^t f)(p) = c_1(b-a) + c_2(b^2-a^2) + \dots + c_n(b^n-a^n) + c_0 - c_0$ (2.0.12)
	$= (c_0 + c_1 b + c_2 b^2 + \dots + c_n b^n) - (c_0 + c_1 a + c_2 a^2 + \dots + c_n a^n)$ (2.0.13)
	$= p(b) - p(a)$ (2.0.14)
	$= p(b) - p(a)$ (2.0.15)
	$= p(b) - p(a)$ (2.0.16)

TABLE1: Proof