

Assignment 9

KUSUMA PRIYA
EE20MTECH11007

Download codes from

<https://github.com/KUSUMAPRIYAPULAVARTY/assignment9>

where \mathbf{C} is a product of elementary matrices

$$\mathbf{C} = \mathbf{E}_5 \mathbf{E}_4 \mathbf{E}_3 \mathbf{E}_2 \mathbf{E}_1 \quad (2.1.2)$$

$$\mathbf{E}_1 = \begin{pmatrix} \frac{1}{A_{11}} & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.3)$$

$$\mathbf{E}_2 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -A_{21} & 1 & 0 & \dots & 0 \\ -A_{31} & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ -A_{n1} & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.4)$$

$$\mathbf{E}_3 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{A_{22} - \frac{A_{21}A_{12}}{A_{11}}} & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.5)$$

$$\mathbf{E}_4 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -A_{32} + \frac{A_{31}A_{12}}{A_{11}} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -A_{n2} + \frac{A_{n1}A_{12}}{A_{11}} & 0 & \dots & 1 \end{pmatrix} \quad (2.1.6)$$

$$\mathbf{E}_5 = \begin{pmatrix} 1 & \frac{-A_{21}}{A_{11}} & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.7)$$

$$\Rightarrow \mathbf{R}_1 = \mathbf{C}\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix} \quad (2.1.8)$$

1 QUESTION

Prove that if two homogenous systems of linear equations in two unknowns have the same solutions, then they are equivalent.

2 SOLUTION

Let the two systems of homogenous equations be

$$\mathbf{A}\mathbf{x} = \mathbf{0} \quad (2.0.1)$$

$$\mathbf{B}\mathbf{x} = \mathbf{0} \quad (2.0.2)$$

$$\Rightarrow \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \\ \vdots & \vdots \\ A_{n1} & A_{n2} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad (2.0.3)$$

$$\text{and } \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \\ \vdots & \vdots \\ B_{n1} & B_{n2} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad (2.0.4)$$

2.1 Case 1

Let us assume that the solution is unique. Since they have the same solution, both \mathbf{A}, \mathbf{B} must have their rank as 2, that is both of them have two independent equations.

Let the reduced row echelon form of \mathbf{A} be \mathbf{R}_1 and \mathbf{B} be \mathbf{R}_2

Finding the reduced row echelon form of \mathbf{A} ,

$$\mathbf{R}_1 = \mathbf{C}\mathbf{A} \quad (2.1.1)$$

Finding the reduced row echelon form of \mathbf{B} ,

$$\mathbf{R}_2 = \mathbf{D}\mathbf{B} \quad (2.1.9)$$

where \mathbf{D} is a product of elementary matrices

$$\mathbf{D} = \mathbf{E}'_5 \mathbf{E}'_4 \mathbf{E}'_3 \mathbf{E}'_2 \mathbf{E}'_1 \quad (2.1.10)$$

$$\mathbf{E}'_1 = \begin{pmatrix} \frac{1}{B_{11}} & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.11)$$

$$\mathbf{E}'_2 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -B_{21} & 1 & 0 & \dots & 0 \\ -B_{31} & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ -B_{n1} & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.12)$$

$$\mathbf{E}'_3 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{B_{22} - \frac{B_{21}B_{12}}{B_{11}}} & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.13)$$

$$\mathbf{E}'_4 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -B_{32} + \frac{B_{31}B_{12}}{B_{11}} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -B_{n2} + \frac{B_{n1}B_{12}}{B_{11}} & 0 & \dots & 1 \end{pmatrix} \quad (2.1.14)$$

$$\mathbf{E}'_5 = \begin{pmatrix} 1 & \frac{-B_{21}}{B_{11}} & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix} \quad (2.1.15)$$

$$\Rightarrow \mathbf{R}_2 = \mathbf{DB} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix} \quad (2.1.16)$$

$$\Rightarrow \mathbf{R}_1 = \mathbf{R}_2 \quad (2.1.17)$$

Hence the given systems are equivalent.

2.2 Case 2

Let us assume that (2.0.3),(2.0.4) have infinitely many solutions

So,

$$\text{either rank}(\mathbf{A}) = \text{rank}(\mathbf{B}) = 1 \quad (2.2.1)$$

$$\text{or rank}(\mathbf{A}) = \text{rank}(\mathbf{B}) = 0 \quad (2.2.2)$$

If, rank of $\mathbf{A} = \text{rank of } \mathbf{B} = 1$

Row reduced echelon forms of \mathbf{A}, \mathbf{B} become $\mathbf{R}_1, \mathbf{R}_2$

$$\mathbf{R}_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix} \quad (2.2.3)$$

$$\mathbf{R}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix} \quad (2.2.4)$$

Hence the same approach as in case 1 yields equivalence of two systems.

Rank zero indicates both \mathbf{A} and \mathbf{B} are null matrices and hence the systems (2.0.3) and (2.0.4) are equivalent.