

Math Notes

K

March 6, 2020


Contents

1	Trigonometric Concepts	3
1.1	Unit Circle	3
1.2	Basic Trigonometry	4
1.3	Pythagorean Theorem Identities	4
1.4	Half Angle Identities	4
1.5	Double Angle Identities	4
1.6	Power Reduction Identities	5
2	Trigonometric Substitution	5
3	Table of Basic Derivatives	5
4	Table of Basic Integrals	5
5	Substitution Techniques	5
5.1	U Substitution	6
5.2	Integration by Parts	6
6	Engineering Formulas	6
6.1	Spring Formulas	6
6.2	Fluid Formulas	7
7	Method of Partial Fractions	7
7.1	Decomposition Types	7
7.2	Numerical Integration	7
7.2.1	Midpoint Rule	7
7.2.2	Trapezoidal Rule	8
7.2.3	Error Approximation (Actual Error)	8
7.2.4	Error Bounds	8
7.3	Improper Integrals	8
7.3.1	Type 1	8
7.3.2	Type 2	8
7.3.3	Type 3	9

7.4 The Length of the Curve	9
8 Area of Revolution	9
9 Separable Equations	9
10 Parametric Equations of Curves	10
11 Polar Coordinates	10
12 Cartesian Coordinates	10
13 Sequences	10
14 Curves in Polar Coordinates	10
15 Sequences (Continued)	10
16 Squeeze Theorem	11
17 Series	11
18 Additional Resources	12

1 Trigonometric Concepts

1.1 Unit Circle



Source: https://etc.usf.edu/clipart/43200/43215/unit-circle7_43215.htm

1.2 Basic Trigonometry

$$\begin{aligned}\sin(\theta) &= \frac{\textit{Opposite}}{\textit{Hypotenuse}} & \cos(\theta) &= \frac{\textit{Adjacent}}{\textit{Hypotenuse}} & \tan(\theta) &= \frac{\textit{Opposite}}{\textit{Adjacent}} \\ &= \frac{y}{1} = y & &= \frac{x}{1} = x & &= \frac{y}{x} \\ & & & & &= \frac{\sin(\theta)}{\cos(\theta)}\end{aligned}$$

$$\begin{aligned}\csc(\theta) &= \frac{1}{\sin(\theta)} & \sec(\theta) &= \frac{1}{\cos(\theta)} & \cot(\theta) &= \frac{\cos(\theta)}{\sin(\theta)}\end{aligned}$$

1.3 Pythagorean Theorem Identities

$$\begin{aligned}\sin^2\theta + \cos^2\theta &= 1 \\ \tan^2\theta + 1 &= \sec^2\theta \\ 1 + \cot^2\theta &= \csc^2\theta\end{aligned}$$

1.4 Half Angle Identities

$$\begin{aligned}\cos\left(\frac{\theta}{2}\right) &= \pm\sqrt{\frac{1 + \cos(\theta)}{2}} & \tan\left(\frac{\theta}{2}\right) &= \pm\sqrt{\frac{1 - \cos(\theta)}{2}} \\ \sin\left(\frac{\theta}{2}\right) &= \pm\sqrt{\frac{1 - \cos(\theta)}{2}} & &= \frac{\sin(\theta)}{1 + \cos(\theta)} \\ & & &= \frac{1 - \cos(\theta)}{\sin(\theta)}\end{aligned}$$

1.5 Double Angle Identities

$$\begin{aligned}\sin(2\theta) &= 2\sin\theta\cos\theta \\ \cos(2\theta) &= \cos^2\theta - \sin^2\theta \\ &= 2\cos^2\theta - 1 \\ &= 1 - 2\sin^2\theta \\ \tan(2\theta) &= \frac{2\tan\theta}{1 - \tan^2\theta}\end{aligned}$$

1.6 Power Reduction Identities

$$\sin^2\theta = \frac{1 - \cos 2\theta}{2}$$

$$\cos^2\theta = \frac{1 + \cos 2\theta}{2}$$

$$\tan^2\theta = \frac{1 - \cos 2\theta}{1 + \cos 2\theta}$$

$$\csc^2\theta = \frac{2}{1 - \cos 2\theta}$$

$$\sec^2\theta = \frac{2}{1 + \cos 2\theta}$$

$$\cot^2\theta = \frac{1 + \cos 2\theta}{1 - \cos 2\theta}$$

2 Trigonometric Substitution

Expression	Substitution	Identity
$\sqrt{a^2 - b^2x^2}$	$x = \frac{a}{b}\sin\theta$	$1 - \sin^2\theta = \cos^2\theta$
$\sqrt{a^2 + b^2x^2}$	$x = \frac{a}{b}\tan\theta$	$1 + \tan^2\theta = \sec^2\theta$
$\sqrt{b^2x^2 - a^2}$	$x = \frac{a}{b}\sec\theta$	$\sec^2\theta - 1 = \tan^2\theta$

3 Table of Basic Derivatives

y	$\frac{dy}{dx}$
C	0
x	1
$ax^2 + bx + c$	$2ax + b$
x^n	nx^{n-1}
$x^{-1}, \frac{1}{x}$	$-\frac{1}{x^2}$
\sqrt{x}	$\frac{1}{2\sqrt{x}}$
$\sqrt[n]{x}$	$\frac{1}{n\sqrt[n]{x^{n-1}}}$
$\ln(x)$	$\frac{1}{x}$
$\log_a(x)$	$\frac{1}{x\ln(a)}$
a^x	$a^x \ln(a)$
e^x	e^x
$\sin(x)$	$\cos(x)$
$\cos(x)$	$-\sin(x)$
$\tan(x)$	$\frac{1}{\cos^2 x}, \sec^2 x$
$\cot(x)$	$-\frac{1}{\sin^2 x}, -\csc^2 x$
$\sec(x)$	$\tan(x)\sec(x)$
$\csc(x)$	$-\cot(x)\csc(x)$

4 Table of Basic Integrals

$f(x)$	$\int f(x)dx = F(x) + C$
x^α $\alpha \neq 0$	$\frac{x^{\alpha+1}}{\alpha+1} + C$
$\sin(kx)$	$-\frac{\cos(kx)}{k} + C$
$\cos(kx)$	$\frac{\sin(kx)}{k} + C$
$\sec^2(kx)$	$\frac{\tan(kx)}{k} + C$
$\csc^2(kx)$	$-\frac{\cot(kx)}{k} + C$
e^{kx}	$\frac{e^{kx}}{k} + C$
$x^{-1}, \frac{1}{x}$	$\ln x + C$
$\frac{1}{\sqrt{1-x^2}}$	$\sin^{-1} x + C$
$\frac{1}{1+x^2}$	$\tan^{-1} x + C$
a^{kx}	$\frac{1}{k\ln a} a^{kx} + C$
$\frac{1}{\sqrt{a^2-x^2}}$	$\sin^{-1} \frac{x}{a} + C$
$\frac{1}{\sqrt{a^2+x^2}}$	$\frac{1}{a} \tan^{-1}(x/a) + C$

5 Substitution Techniques

5.1 U Substitution

$$\int f(g(x))g'(x)dx = \int f(u)du = F(u) + C$$

where: $u = g(x)$, $du = g'(x)$

5.2 Integration by Parts

$$\int f(x)g'(x)dx = f(x)g(x) - \int g(x)f'(x)dx$$

$$\int u dv = uv - \int v du$$

$$\int_a^b f(x)g'(x) = f(x)g(x)\Big|_a^b - \int_a^b g(x)f'(x)dx$$

6 Engineering Formulas

6.1 Spring Formulas

$$F = kx$$

$$W = \int_a^b kx dx$$

F = Force (Newtons [N])

k = spring constant (Newton meters $^N/m$)

x = change in distance (meters [m])

W = Work (Joules [J])

a = initial length (meters [m])

b = final length (meters [m])

6.2 Fluid Formulas

$$W = F \cdot d = \int F dx$$

$$V = \pi r^2 h (\text{apply to cylinders})$$

$$F = m \cdot a = V \cdot \rho$$

$$W = \text{Weight } ()$$

$$F = \text{Force (Newtons [N])}$$

$$d = \text{distance (meters [d])}$$

$$m = \text{mass (meters}^3[\text{m}^3])$$

$$a = \text{acceleration (meters per second}^2 [\text{m/s}^2])$$

$$\rho = \text{Something}$$

7 Method of Partial Fractions

$$\int \frac{P_n(x)}{Q_m(x)} dx \text{ when } m > n$$

n and m are defined as the degree of the numerator and the denominator.

7.1 Decomposition Types

Type	Factor Example	Decomposition
Linear Factor	$(x - 4)$	$\frac{A}{x-4}$
Repeated Linear Factor	$(x - 4)^2$	$\frac{A}{(x+4)} + \frac{B}{(x+4)^2}$
Quadratic Irreducible Factor	$(x^2 + 4)$	$\frac{Ax+B}{x^2+4}$
Repeated Quadratic Irreducible Factor	$(x^2 + 4)^2$	$\frac{Ax+B}{(x^2+4)} + \frac{Cx+D}{(x^2+4)^2}$

7.2 Numerical Integration

7.2.1 Midpoint Rule

$$\int_a^b f(x) dx \approx \sum_{i=1}^n f(\bar{x}_i) \Delta x$$

$$x_i^* = \frac{x_{i-1} + x_i}{2} = \bar{x}_i$$

$$x_i^* \in [x_{i-1}, x_i]$$

7.2.2 Trapezoidal Rule

$$\begin{aligned}\int_a^b f(x)dx &\approx \sum_{i=1}^n \frac{f(x_{i-1}) + f(x_i)}{2} \Delta x \\ &\approx \sum_{i=1}^n \frac{\Delta x}{2} [f(x_{i-1}) + f(x_i)]\end{aligned}$$

7.2.3 Error Approximation (Actual Error)

$$\begin{aligned}E_T &= \int_a^b f(x)dx - T_n \\ E_H &= \int_a^b f(x)dx - M_n\end{aligned}$$

7.2.4 Error Bounds

$$\begin{aligned}|f''(x)| &\leq K \text{ on } [b, a], f(x) = \frac{1}{x}, f''(x) = -\frac{2}{x^3} \\ \left| -\frac{2}{x^3} \right| &\leq 2 = K \\ |E_T| &\leq \frac{K(b-a)^3}{12n^2} = \frac{2(2-1)^3}{12 \cdot 5^2} = 0.0012 \\ |E_M| &\leq \frac{K(b-a)^3}{24n^2} = \frac{2(2-1)^3}{24 \cdot 5^2} = 0.0012 \\ E_T &\leq 0.6931 - 0.6956\end{aligned}$$

7.3 Improper Integrals

7.3.1 Type 1

$x \in [a, b]$, $f(x)$ is continuous on $[a, b)$ at $x = b$ $f(x)$ discontinues

$$\int_a^\infty f(x) = \lim_{x \rightarrow \infty} \left(\int_a^b f(x)dx \right)$$

7.3.2 Type 2

$f(x)$ dicontinues at $x = a$, continuous on $(a, b]$

$$\int_a^b f(x)dx = \lim_{t \rightarrow a} \int_t^b f(x)dx$$

7.3.3 Type 3

$f(x)$ discontinues at $x = c$, $a < c < b$ everywhere else on $[a, b]$
 $f(x)$ is continuous

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$

7.4 The Length of the Curve

$$\begin{array}{lll} x = x(t) & y = y(t) & a \leq t \leq b \\ S' = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} \end{array}$$

8 Area of Revolution

$$\begin{array}{l} ds = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\ dA = 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \\ 1) \int_a^b dA = A = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\ 2) x = xb \end{array}$$

9 Separable Equations

$$y'(x) = g(x)h(y), \frac{dy}{dx} = g(x)h(y) \Big| dx$$

$$dy = h(y)g(x)dx \Big| \frac{1}{h(y)}$$

$$\frac{1}{h(y)}dy = g(x)dx, H(y) = \frac{1}{h(y)} \longrightarrow H(y)dy = g(x)dx$$

$$\int H(y) = \int g(x)dx + C$$

Implicit Form of Solution

$$F(y) = G(x) + C$$

Explicit Form of Solution

$$y = F^{-1}(G(x) + C)$$

10 Parametric Equations of Curves

$$t_0 = \begin{pmatrix} x_0 = x(t_0) \\ y_0 = y(t_0) \end{pmatrix}$$

$$t_1 = \begin{pmatrix} x_1 = x(t_1) \\ y_1 = y(t_1) \end{pmatrix}$$

$$\begin{aligned} x &= x(t), y = y(t), \alpha \leq t \leq \beta \\ &= \varphi(x)y = (y(\varphi(x))) = f(x) \end{aligned}$$

$$t_2 = \begin{pmatrix} x_2 = x(t_2) \\ y_2 = y(t_2) \end{pmatrix}$$

11 Polar Coordinates

$$x = r \cos \theta, y = r \sin \theta$$

Note: Angles are CounterClockwise from the Positive X

12 Cartesian Coordinates

$$r^2 = x^2 + y^2, \tan \theta = \frac{y}{x}$$

13 Sequences

Discrete

$$a_n = f(n)$$

$$a_1, a_2, a_3, \dots, a_n, a_{n+1}, \dots$$

$$n \rightarrow a_n$$

$$1 \rightarrow a, 2 \rightarrow a_2, \dots$$

$$\{a_n\}_{n=1}^{\infty} = \{a_1, a_2, \dots, a_n, \dots\} = \{a_n\}$$

• can not take derivative

• can take the limit

14 Curves in Polar Coordinates

$$x = r \cos \theta, y = r \sin \theta, r = f(\theta)$$

$$x = x(t), u = y(t)$$

$$\alpha \leq \theta \leq \beta$$

15 Sequences (Continued)

$$a_1, a_2, a_3, \dots, a_n, a_{n+1}, \dots$$

if $\{a_n\}, \{b_n\}$ converges, then:

$$1. \lim_{n \rightarrow \infty} (a_n \pm b_n) = \lim a_n \pm \lim b_n$$

$$2. \lim_{n \rightarrow \infty} (c \cdot a_n) = c \lim a_n$$

3. $\lim_{n \rightarrow \infty} (a_n, b_n) = \lim a_n \cdot \lim b_n$
4. $\lim \frac{a_n}{b_n} = \frac{\lim a_n}{\lim b_n}, b \neq 0$
5. $\lim_{n \rightarrow \infty} (a_n^p) = (\lim a_n)^p, a_n, p > 0$

16 Squeeze Theorem

for $n > n_0, a_n \leq b_n \leq c_n$
 $\{a_n\}, \{b_n\}, \{c_n\} \rightarrow$ sequences
 $\lim a_n = \lim c_n = L$, then
 $\lim b_n = L$

1. $\lim_{n \rightarrow \infty} \frac{\lim}{n} = 0$
2. $\lim \sqrt[n]{n} = 1$
3. $\lim x^n = 0, |x| < 1$
4. $\lim x^{\frac{1}{n}}, x > 0 > 1$
5. $\lim (1 + \frac{x}{n})^n = e^x$
6. $\lim \frac{x^n}{n!} = 0, y$ any x

17 Series

$\{a_n\}_{n=1}^{\infty}, a_1 + a_2 + a_3 + \cdots + a_n + \cdots$
 Infinite Series
 $\sum_{n=1}^{\infty} a_n, \sum a_n$

18 Additional Resources

Print

Calculus Study Guide: <https://mt-jfk.com/ap-calculus-study-guide.pdf>

Video

The Organic Chemistry Tutor: <https://www.youtube.com/channel/UCeWpbFLzoYGPfuWUMFPSaoA>

Black Pen Red Pen: <https://www.youtube.com/user/blackpenredpen>