

Math Notes

K

March 6, 2020


Contents

1	Trigonometric Concepts	3
1.1	Unit Circle	3
1.2	Basic Trigonometry	4
1.3	Pythagorean Theorem Identities	4
1.4	Half Angle Identities	4
1.5	Double Angle Identities	4
1.6	Power Reduction Identities	5
2	Trigonometric Substitution	5
3	Table of Basic Derivatives	5
4	Table of Basic Integrals	5
5	Substitution Techniques	5
5.1	U Substitution	6
5.2	Integration by Parts	6
6	Engineering Formulas	6
6.1	Spring Formulas	6
6.2	Fluid Formulas	7
7	Method of Partial Fractions	7
7.1	Decomposition Types	7
7.2	Numerical Integration	7
7.2.1	Midpoint Rule	7
7.2.2	Trapezoidal Rule	8
7.2.3	Error Approximation (Actual Error)	8
7.2.4	Error Bounds	8
7.3	Improper Integrals	8
7.3.1	Type 1	8
7.3.2	Type 2	8
7.3.3	Type 3	9

7.4 The Length of the Curve	9
8 Additional Resources	10

1 Trigonometric Concepts

1.1 Unit Circle



Source: https://etc.usf.edu/clipart/43200/43215/unit-circle7_43215.htm

1.2 Basic Trigonometry

$$\begin{aligned}\sin(\theta) &= \frac{\textit{Opposite}}{\textit{Hypotenuse}} & \cos(\theta) &= \frac{\textit{Adjacent}}{\textit{Hypotenuse}} & \tan(\theta) &= \frac{\textit{Opposite}}{\textit{Adjacent}} \\ &= \frac{y}{1} = y & &= \frac{x}{1} = x & &= \frac{y}{x} \\ & & & & &= \frac{\sin(\theta)}{\cos(\theta)}\end{aligned}$$

$$\begin{aligned}\csc(\theta) &= \frac{1}{\sin(\theta)} & \sec(\theta) &= \frac{1}{\cos(\theta)} & \cot(\theta) &= \frac{\cos(\theta)}{\sin(\theta)}\end{aligned}$$

1.3 Pythagorean Theorem Identities

$$\begin{aligned}\sin^2\theta + \cos^2\theta &= 1 \\ \tan^2\theta + 1 &= \sec^2\theta \\ 1 + \cot^2\theta &= \csc^2\theta\end{aligned}$$

1.4 Half Angle Identities

$$\begin{aligned}\cos\left(\frac{\theta}{2}\right) &= \pm\sqrt{\frac{1+\cos(\theta)}{2}} & \tan\left(\frac{\theta}{2}\right) &= \pm\sqrt{\frac{1-\cos(\theta)}{2}} \\ \sin\left(\frac{\theta}{2}\right) &= \pm\sqrt{\frac{1-\cos(\theta)}{2}} & &= \frac{\sin(\theta)}{1+\cos(\theta)} \\ & & &= \frac{1-\cos(\theta)}{\sin(\theta)}\end{aligned}$$

1.5 Double Angle Identities

$$\begin{aligned}\sin(2\theta) &= 2\sin\theta\cos\theta \\ \cos(2\theta) &= \cos^2\theta - \sin^2\theta \\ &= 2\cos^2\theta - 1 \\ &= 1 - 2\sin^2\theta \\ \tan(2\theta) &= \frac{2\tan\theta}{1 - \tan^2\theta}\end{aligned}$$

1.6 Power Reduction Identities

$$\sin^2\theta = \frac{1 - \cos 2\theta}{2}$$

$$\cos^2\theta = \frac{1 + \cos 2\theta}{2}$$

$$\tan^2\theta = \frac{1 - \cos 2\theta}{1 + \cos 2\theta}$$

$$\csc^2\theta = \frac{2}{1 - \cos 2\theta}$$

$$\sec^2\theta = \frac{2}{1 + \cos 2\theta}$$

$$\cot^2\theta = \frac{1 + \cos 2\theta}{1 - \cos 2\theta}$$

2 Trigonometric Substitution

Expression	Substitution	Identity
$\sqrt{a^2 - b^2x^2}$	$x = \frac{a}{b}\sin\theta$	$1 - \sin^2\theta = \cos^2\theta$
$\sqrt{a^2 + b^2x^2}$	$x = \frac{a}{b}\tan\theta$	$1 + \tan^2\theta = \sec^2\theta$
$\sqrt{b^2x^2 - a^2}$	$x = \frac{a}{b}\sec\theta$	$\sec^2\theta - 1 = \tan^2\theta$

3 Table of Basic Derivatives

y	$\frac{dy}{dx}$
C	0
x	1
$ax^2 + bx + c$	$2ax + b$
x^n	nx^{n-1}
$x^{-1}, \frac{1}{x}$	$-\frac{1}{x^2}$
\sqrt{x}	$\frac{1}{2\sqrt{x}}$
$\sqrt[n]{x}$	$\frac{1}{n\sqrt[n]{x^{n-1}}}$
$\ln(x)$	$\frac{1}{x}$
$\log_a(x)$	$\frac{1}{x\ln(a)}$
a^x	$a^x \ln(a)$
e^x	e^x
$\sin(x)$	$\cos(x)$
$\cos(x)$	$-\sin(x)$
$\tan(x)$	$\frac{1}{\cos^2 x}, \sec^2 x$
$\cot(x)$	$-\frac{1}{\sin^2 x}, -\csc^2 x$
$\sec(x)$	$\tan(x)\sec(x)$
$\csc(x)$	$-\cot(x)\csc(x)$

4 Table of Basic Integrals

$f(x)$	$\int f(x)dx = F(x) + C$
x^α $\alpha \neq 0$	$\frac{x^{\alpha+1}}{\alpha+1} + C$
$\sin(kx)$	$-\frac{\cos(kx)}{k} + C$
$\cos(kx)$	$\frac{\sin(kx)}{k} + C$
$\sec^2(kx)$	$\frac{\tan(kx)}{k} + C$
$\csc^2(kx)$	$-\frac{\cot(kx)}{k} + C$
e^{kx}	$\frac{e^{kx}}{k} + C$
$x^{-1}, \frac{1}{x}$	$\ln x + C$
$\frac{1}{\sqrt{1-x^2}}$	$\sin^{-1} x + C$
$\frac{1}{1+x^2}$	$\tan^{-1} x + C$
a^{kx}	$\frac{1}{k\ln a} a^{kx} + C$
$\frac{1}{\sqrt{a^2-x^2}}$	$\sin^{-1} \frac{x}{a} + C$
$\frac{1}{\sqrt{a^2+x^2}}$	$\frac{1}{a} \tan^{-1}(x/a) + C$

5 Substitution Techniques

5.1 U Substitution

$$\int f(g(x))g'(x)dx = \int f(u)du = F(u) + C$$

where: $u = g(x)$, $du = g'(x)$

5.2 Integration by Parts

$$\int f(x)g'(x)dx = f(x)g(x) - \int g(x)f'(x)dx$$

$$\int u dv = uv - \int v du$$

$$\int_a^b f(x)g'(x) = f(x)g(x)\Big|_a^b - \int_a^b g(x)f'(x)dx$$

6 Engineering Formulas

6.1 Spring Formulas

$$F = kx$$

$$W = \int_a^b kx dx$$

F = Force (Newtons [N])

k = spring constant (Newton meters $^N/m$)

x = change in distance (meters [m])

W = Work (Joules [J])

a = initial length (meters [m])

b = final length (meters [m])

6.2 Fluid Formulas

$$W = F \cdot d = \int F dx$$

$$V = \pi r^2 h (\text{apply to cylinders})$$

$$F = m \cdot a = V \cdot \rho$$

$$W = \text{Weight } ()$$

$$F = \text{Force (Newtons [N])}$$

$$d = \text{distance (meters [d])}$$

$$m = \text{mass (meters}^3[\text{m}^3])$$

$$a = \text{acceleration (meters per second}^2 [\text{m/s}^2])$$

$$\rho = \text{Something}$$

7 Method of Partial Fractions

$$\int \frac{P_n(x)}{Q_m(x)} dx \text{ when } m > n$$

n and m are defined as the degree of the numerator and the denominator.

7.1 Decomposition Types

Type	Factor Example	Decomposition
Linear Factor	$(x - 4)$	$\frac{A}{x-4}$
Repeated Linear Factor	$(x - 4)^2$	$\frac{A}{(x+4)} + \frac{B}{(x+4)^2}$
Quadratic Irreducible Factor	$(x^2 + 4)$	$\frac{Ax+B}{x^2+4}$
Repeated Quadratic Irreducible Factor	$(x^2 + 4)^2$	$\frac{Ax+B}{(x^2+4)} + \frac{Cx+D}{(x^2+4)^2}$

7.2 Numerical Integration

7.2.1 Midpoint Rule

$$\int_a^b f(x) dx \approx \sum_{i=1}^n f(\bar{x}_i) \Delta x$$

$$x_i^* = \frac{x_{i-1} + x_i}{2} = \bar{x}_i$$

$$x_i^* \in [x_{i-1}, x_i]$$

7.2.2 Trapezoidal Rule

$$\begin{aligned}\int_a^b f(x)dx &\approx \sum_{i=1}^n \frac{f(x_{i-1}) + f(x_i)}{2} \Delta x \\ &\approx \sum_{i=1}^n \frac{\Delta x}{2} [f(x_{i-1}) + f(x_i)]\end{aligned}$$

7.2.3 Error Approximation (Actual Error)

$$\begin{aligned}E_T &= \int_a^b f(x)dx - T_n \\ E_H &= \int_a^b f(x)dx - M_n\end{aligned}$$

7.2.4 Error Bounds

$$\begin{aligned}|f''(x)| &\leq K \text{ on } [b, a], f(x) = \frac{1}{x}, f''(x) = -\frac{2}{x^3} \\ \left| -\frac{2}{x^3} \right| &\leq 2 = K \\ |E_T| &\leq \frac{K(b-a)^3}{12n^2} = \frac{2(2-1)^3}{12 \cdot 5^2} = 0.0012 \\ |E_M| &\leq \frac{K(b-a)^3}{24n^2} = \frac{2(2-1)^3}{24 \cdot 5^2} = 0.0012 \\ E_T &\leq 0.6931 - 0.6956\end{aligned}$$

7.3 Improper Integrals

7.3.1 Type 1

$x \in [a, b]$, $f(x)$ is continuous on $[a, b)$ at $x = b$ $f(x)$ discontinues

$$\int_a^\infty f(x) = \lim_{x \rightarrow \infty} \left(\int_a^b f(x)dx \right)$$

7.3.2 Type 2

$f(x)$ is continuous at $x = a$, continuous on $(a, b]$

$$\int_a^b f(x)dx = \lim_{t \rightarrow a} \int_t^b f(x)dx$$

7.3.3 Type 3

$f(x)$ discontinues at $x = c$, $a < c < b$ everywhere else on $[a, b]$
 $f(x)$ is continuous

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$

7.4 The Length of the Curve

$$x = x(t)$$

$$y = y(t)$$

$$a \leq t \leq b$$

$$S' = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2}$$

8 Additional Resources

Print

Calculus Study Guide: <https://mt-jfk.com/ap-calculus-study-guide.pdf>

Video

The Organic Chemistry Tutor: <https://www.youtube.com/channel/UCEWpbFLzoYGPfuWUMFPSaoA>

Black Pen Red Pen: <https://www.youtube.com/user/blackpenredpen>