
12/8/2020 no2b2

```
clear all;
close all;
f = @(x) \sin(pi*x/2) + 0.5*\sin(2*pi*x);
g0 = Q(t) 0;
g1 = @(t) exp((-pi^2*t)/4);
%Exact solution
uexact = @(x,t) \exp(-(pi^2)*t/4).*sin((pi*x)/2) + 0.5*exp(-(4*pi^2)*t).*sin(2*pi*x);
alp = 1;
tspan = 1;
n = 4:8;
m = 2.^n - 1;
N = 2.^n;
Re_Err = zeros(5,1);
for i =1:5
    %Numerical solution
   [u,t,x] = cnhteq(f,g0,g1,tspan,alp,N(i),m(i));
   u = u(end,:);
   %exact solution
    uex = uexact(x,1)';
    Re_Err(i) = RelNorm(u,uex);
end
loglog(m,Re_Err);
title('A graph of Relative Error against m');
xlabel('m');
ylabel('Relative error');
p = polyfit(log(m),log(Re_Err),1);
fprintf('The order of accuracy is %f \n', p(1));
function L2 = RelNorm(U,Uexact)
error = (U - Uexact).^2;
L2 = sqrt(sum(error)/sum(Uexact.^2));
end
```

The order of accuracy is -1.948416

12/8/2020 no2b2



Published with MATLAB® R2020a