

# Simulating Tropical Cyclone over the South-west Indian Ocean with MPAS: Sensitivity to cloud microphysics schemes

Brian KYANJO (brian.kyanjo@aims.ac.rw)  
African Institute for Mathematical Sciences (AIMS) Rwanda

Supervised by: Prof. Babatunde J. Abiodun  
University of Cape Town, South Africa

May 2020

*Submitted in partial fulfilment of the requirements of a Master of Science in Mathematical Sciences at AIMS Rwanda*



**AIMS**

**African Institute for  
Mathematical Sciences  
RWANDA**

# Abstract

Tropical cyclones (TCs) are one of the strongest atmospheric features known for massive devastating destructions. This study compares the ability of Model for Prediction Across Scales (MPAS) cloud microphysics schemes (Thompson and WSM6) in simulating TCs over the Southwest Indian Ocean (SWIO) basin. In the study, Thompson and WSM6 are tested with a 25 km variable resolution over the SWIO. The simulation is conducted for nine days (11<sup>th</sup>/02/2017 - 19<sup>th</sup>/02/2017) and evaluated against ERA5 reanalysis and the Regional Specialised meteorological Centre (RSMC) best track data. Thompson exhibits a reasonable simulation of TCs and depicts some competence in simulating the spatial and temporal distribution of TCs tracks and locations over SWIO. However, there are sometimes inconsistencies between the observed and the simulated TC by both WSM6 and Thompson over the entire domain. The pressure at the centre (eye) and the maximum sustained wind speeds (MSW) in the eyewall are well simulated by both schemes, even though Thompson is more realistic than WSM6. Thompson well exhibits the daily temporal variation of TCs. In this study, simulations of TCs show adaptations of improving and using Thompson for predicting TCs over the basin.

## Declaration

I, the undersigned, hereby declare that the work contained in this research project is my original work, and that any work done by others or by myself previously has been acknowledged and referenced accordingly.



---

Brian Kyanjo, May 2020

## List of Abbreviations

ARW	Advanced Research WRF.
CAM	Community Atmosphere Model.
CFS	Climate Forecast System.
CFSR	Climate Forecast System Reanalysis.
CHPC	Centre for High Performance Computing.
MPAS	Model for Prediction Across Scales.
MSW	Maximum Sustained winds.
MYNN	Mellor-Yamada-Nakanishi-Niino.
NCAR	National Center for Atmospheric Research.
NCEP	National Centres for Environmental Prediction.
NCL	NCAR Command language.
RDA	Research Data Archive.
RRTMG	Rapid Radiative Transfer Model for General Circulation Models.
SST	Sea Surface Temperature.
SWIO	South West Indian Ocean.
TCs	Tropical Cyclones.
WRF	Weather Research Forecast.
WSM6	WRF-Single-Moment-Microphysics Class 6.
YSU	Yonsei University.

# Contents

<b>Abstract</b>	<b>i</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Background . . . . .	1
1.2 Problem Statement . . . . .	2
1.3 Aim and objectives . . . . .	2
1.4 Overview . . . . .	2
<b>2 Literature Review</b>	<b>3</b>
2.1 Tropical cyclones (TCs) Over the SWIO . . . . .	3
2.2 MPAS overview . . . . .	3
<b>3 Methodology and Dataset</b>	<b>5</b>
3.1 Study Area . . . . .	5
3.2 Datasets . . . . .	6
3.3 MPAS . . . . .	6
3.4 Model simulation of TC Dineo . . . . .	10
3.5 Detecting and Tracking TC Dineo . . . . .	11
<b>4 Results and discussions</b>	<b>13</b>
4.1 Track of TC Dineo . . . . .	13
4.2 Horizontal structure of TC Dineo . . . . .	18
4.3 Vertical cross section of TC Dineo . . . . .	19
<b>5 Conclusion</b>	<b>22</b>
<b>References</b>	<b>25</b>

# 1. Introduction

## 1.1 Background

A TC is a rapidly rotating storm system distinguished by a warm core, a closed low-level atmospheric circulation, low-pressure center, robust cyclonic surface circulation, convection, and a spiral arrangement of thunderstorms that manufacture significant rain. TCs develop over tropical or subtropical waters. There are three fundamental stages in the evolution of a TC: its origin, mature stage, and dissipation stage where it finally stops. They occur in a continuous process, not as separate, distinct stages, each stage may occur more than once during the life cycle as the strength of the cyclone rises and falls it may also reach land weakened then returns to the sea where it strengthens again ([tropmet](#)). The formation of a cyclone occurs after satisfying certain conditions i.e., developing storm should maintain contact with warm water ( $\geq 27^\circ$ ), constant evaporation of seawater from the sea surface, wind pattern near the ocean surface that spirals air inwards, deep moisture at low levels, light winds throughout the troposphere and convergence of the easterly waves. Once satisfied, they allow a body of warm air to develop above the ocean surface which expands becomes lighter and rises, then other local winds blowing replaces it, then also warms and rises. The rising air (contains vast amounts of moisture evaporated from the ocean's surface) cools and condenses to form huge clouds about 10km up in the troposphere ([Joseph A. Zehnder](#)). More warm air rushes in and rises carrying moisture high into the atmosphere so that the formed clouds become very thick and heavy. Condensation then releases latent heat stored in the water vapor providing the cyclone with more power which creates a self-sustaining heat cycle drawn further upwards by the new release of energy; the clouds can grow into 12 to 25km high. Also, the force created by the Earth's rotation on a tilted axis (the Coriolis effect) causes rising currents of air to spiral around the center of the TC ([ThoughtCo](#)). It is at this stage that the TC matures and the eye of the storm is created as the air rises and cools, some of this dense air descends to form a still clear eye as the cyclone rages around it, the eyewall where the wind is strongest behaves like a rotating cylinder ([JetStream](#)). Once the TC is formed, its movement follows a pathway away from its source driven by global wind circulation as warm ocean waters feed it with heat and moisture ([Natural Disasters](#)). It continues to enlarge and will last as long as there atmospheric and oceanic conditions remain favourable, i.e., the storm should still be with warm ocean waters which can last as long as seven days. Once it makes a landfall, it loses its source of energy and momentum and then begins to disappear which is called dissipating ([Roy and Kovordányi, 2012](#)).

Tropical cyclones (TCs), one of the strongest atmospheric features, are known for massive destruction of properties and human lives, because they are associated with strong winds, heavy rains, and storm surges ([TROPICAL CYCLONES](#)). Since the beginning of the 20th century, TCs have affected more than 629 million people and killed more than 1.33 million people in the tropics ([Doocy et al., 2013](#)). TCs, which usually form over warm tropical oceans ( $\geq 26^\circ\text{C}$ ), are warm-core low-pressure systems with 100 - 1000 km diameter. Once established, TCs sustain their strength by absorbing warm moisture as they continue to enlarge and will last as long as the atmospheric and oceanic conditions remain favourable over the ocean. After making a landfall, TCs can last for seven days and cause severe damages as they move inland or along the coastline ([Doocy et al., 2013](#)). The TCs that form over the Southwest Indian Ocean (SWIO) are a threat to the neighbouring countries, but Madagascar, Mauritius and La Reunion experience the most devastating and destructive impacts of the TCs, followed by the extensive coastline and the borders of Zimbabwe and Mozambique ([Tropical cyclones in the South West Indian Ocean](#)). For instance, in 2017, Tropical Cyclone Dineo (name of a tropical cyclone), which formed on 13th February and made a landfall over Mozambique on 15th February ([Moses and Ramotonto, 2018](#)).

Dineo produced heavy rains ( $> 200\text{mm day}^{-1}$ ) and damaging winds in Inhambane (southern province in Mozambique) and destroyed around 20000 homes, directly affected about 130,000 people, and killed seven people. Its widespread torrential rains and the associated storm surge produced massive floods that killed about 271 people and caused damages that exceeded \$ 200 million in Zimbabwe (USD 2017) (Cyclone Dineo. , 2020, 3 15). Although Dineo weakened on February 17th, its remnant low still triggered heavy rainfall with flooding in Botswana. The flooding destroyed 650 households and public infrastructures (e.g. bridges, hospitals, schools, and fields), leaving hundreds of people homeless and poor (especially the low-income earners) (Moses and Ramotonto, 2018). A month after the landfall, the impacts of Dineo landfall induced an outbreak of cholera that infected 1,200 people and killed 2 people in Mozambique and Malawi (Cyclone Dineo. , 2020, 3 15). Despite the devastating impacts TCs over the region, from our best knowledge, there are still no reliable systems for providing early warning on TC occurrences over region. Hence, there is a need for more studies on how to improve simulation and prediction of TC characteristics over the SWIO. This will help reduce the devastating impacts of TCs in the region.

## 1.2 Problem Statement

Model for Prediction Across Scales (MPAS) cloud microphysics schemes: Weather Research Forecast (WRF)-Single-Moment-Microphysics Class 6 (WSM6) and Thompson, have a significant influence in the simulation of TC; however, according to past research, the capability of these two microphysics schemes in simulating TCs over the SWIO have not been tested yet. So comparing and analysing the influence of these microphysics schemes during simulations would help us to understand which scheme is best to use and under which available conditions. This will improve predictions of TCs over the basin and will reduce on the devastating damages caused by TCs ahead of time.

## 1.3 Aim and objectives

This project aims to evaluate the capability of MPAS in simulating TC Dineo and examine the sensitivity of TC Dineo to two prominent cloud microphysics schemes: WSM6 and Thompson over the SWIO basin.

The three specific objectives are:

- To describe the characteristics of TC Dineo.
- To evaluate how MPAS simulates TC Dineo over the SWIO basin.
- To compare the sensitivity of WSM6 and Thompson in the simulation of TC Dineo over the SWIO basin.

## 1.4 Overview

Chapter 2 reviews papers of previous work done relation to this project. It also describes the characteristics of TC Dineo. Chapter 3 describes the steps followed and methods used during the project. Datasets and setups used in designing the model are also explained in this chapter. The Discussion about the results produced during the simulations are well presented in chapter 4. Finally, conclusions and recommendations are discussed in the last chapter.

## 2. Literature Review

This chapter reviews the previous works and the characteristics of TC Dineo. The Model for Prediction Across Scales (MPAS) which is used to simulate TC Dineo is also discussed in this chapter.

### 2.1 Tropical cyclones (TCs) Over the SWIO

Of all the basins in the tropics in which TCs form, previous research shows that SWIO is one of the basin that is mostly affected by TCs. [Mavume et al. \(2009\)](#) research on climatology and landfall of TCs over SWIO showed that TCs develop in eastern Madagascar and Mozambique channel. Observations show that 5% of TCs that form a landfall over Madagascar finally reach Mozambique, 65.4% of the 5% form within the greater SWIO basin and the remaining 34.5% evolve within the Mozambique channel ([Fitchett and Grab, 2014](#)). Apart from cyclone Dineo whose remnants struck Botswana, research shows that in most cases, weakened TCs move further towards westward affecting South African countries but occasionally reach Botswana.

**2.1.1 Characteristics of Tropical Cyclone Dineo.** The TC Dineo which is ranked as the 5th TC of the SWIO season developed from the Mozambique channel on 13th February 2017. At 10 am EST on the same day, it had reached MSW of about 40 knots and was dragging towards the south-southwest at 2 knots with a storm height of 9.9 miles. It was crawling with developing low-level circulation center, that consisted of deep convective bands which were binding into the partially exposed low-level center, with its eastern side composed of bulk thunderstorms as it approached Mozambique ([Sciences, E. , n.d.](#)). On 14th February, Dineo's northeastern quadrant consisted of extremely heavy precipitation in the bands of thunderstorms dropping at over 132 mm per hour. At 10 am EST on 15th February, Dineo reached its peak strength with MSWs near 70 knots with a cloud-filled eye, at this point, it was near  $23.5^{\circ}S$  and  $35.8^{\circ}E$  which is just off Mozambique coast ([NASA, 2017](#)). Later at around 12:30 pm EST on the same day, it made a landfall over Inhambane (Southern Mozambique) with dangerous storm surge, torrential rain and high winds ([Mozambique. , n.d.](#)). On the same day, Dineo's clouds expanded over Inhambane, eastern Botswana, Swaziland, and northeastern South Africa. In Swaziland, the intense storm with height about 8.4 miles produced rainfall falling at over 86 mm per hour. At 7 pm EST, Ex-Tropical Cyclone Dineo was near  $23.4^{\circ}S$  and  $34.1^{\circ}E$  and moving to the west at 9 knots with MSWs at around 60 knots. At 4:16 am EST on February 16th, Dineo's storm elongated and progressed inland, its MSW had reduced to about 60 knots and was still dropping as it continued inland ([NASA, 2017](#)). On 17th February Dineo weakened to remnant low which struck Botswana and triggered heavy rains that initiated floods in the country ([Moses and Ramotonto, 2018](#)).

### 2.2 MPAS overview

Several studies have shown that MPAS is one of the atmospheric models that have great potential for providing reliable early warning information on TC occurrence and characteristics over the SWIO. Furthermore, MPAS has been frequently used to simulate different atmospheric features and processes, including real-time weather, seasonal forecasting of tropical cyclones, tornadoes, and convection ([ncar.ucar.edu. , n.d.](#)). For instance, [Hashimoto et al. \(2016\)](#) used MPAS to reproduce the frequency and locations of North Atlantic TCs using mesh refinement over the North Atlantic ocean. They show that MPAS simulations perform better than the Weather Research and Forecasting System (WRF) simulation because the smooth transitions between two grid resolutions in MPAS (i.e., variable mesh grid) helps the model to simulate smoother and more realistic mid-level jet structures and seasonal cloud patterns than the

WRF simulations, which is influenced nested boundary condition problems. [Harris and Lin \(2014\)](#) also show that MPAS performs better than the High Resolution Model (HiRAM) in simulating the frequency of TCs. Although HIRAM can run at high resolution, it ignores computation of vertical acceleration, and its representation of thunderstorms and convection are still small compared to those in observation and MPAS simulations. However, despite the great potential of MPAS in simulating TCs, no studies have investigated its capability in simulating tropical cyclones over the SWIO.

Successful simulations of TCs with any atmospheric model depend on how well microphysics processes are represented in the model. Microphysics contains resolved water vapour, cloud, and precipitation formation processes ([Skamarock, 2008](#)). For this study's case, the capability of the two microphysics schemes: WSM6 and Thompson in the simulation of TC Dineo over the SWIO basin are compared. The selection of these two microphysics schemes is based on the fact that WSM6 is a default scheme in microphysics and has to be compared with either Kessler or Thompson. Thompson is selected over Kessler because it is uniquely coupled with the rapid radiative transfer model for general circulation models (RRTMG) radiation scheme and also has a unique cooperation with other parameterizations in the same physics suit. Therefore, it provides consistent information about cloud droplet size distributions during modelling. This makes WSM6 and Thompson the two prominent microphysics schemes to be used in MPAS during the entire simulations.

Following the previous research about WSM6 and Thompson schemes, below are some of the comparisons made between the two schemes. The particles associated with graupel in WSM6 are assumed to follow an exponential size distribution ([Hong, 2006](#)). Contrarily, in Thompson, the distribution of the size of snow is assumed to be given by the sum of gamma and exponential distributions and depends on both temperature and water content in the snow. Observations show that Thompson assumes that the shape of snow is non-spherical with its diameter inversely proportional to its bulk density, making it different from WSM6, which assumes snow is spherical and has a constant radius. Furthermore, the vertical heating profiles of freezing and melting processes in WSM6 are computed with high accuracy towards the fall-time sub-steps, this reduces its sensitivity compared to Thompson ([Skamarock, 2008](#)). [Federico \(2016\)](#) used WSM6 in the Regional Atmospheric Modelling System (RAMS) at 5km horizontal grid resolution to simulate an idealized 2D thunderstorm and 3D real case heavy precipitation over Korea. WSM6 results showed a positive and significant impact on the temporal storm evolution and precipitation at the surface. [Su and Fung \(2018\)](#) implemented Thompson in the Weather Research and Forecast model coupled with chemistry (WRF-chem) to simulate the impact of dust on ice nucleation processes in the atmosphere during spring over East Asia. Although observations were consistent with satellite observations after evaluating the effect of dust on ice nucleation processes, simulated ice crystal number concentration and ice water mixing ratio were found to be 7 and 15% higher than the ones simulated without dust. [Hashimoto et al. \(2016\)](#) compared the capacity of WSM6 in WRF model and Thompson in MPAS in capturing the activity of Northwest Pacific TCs. Results showed that WSM6 determined only the mass mixing ratios of the rain species while Thompson determined both the number concentration and the mass mixing ratios of precipitation species. However, both schemes used only six water species during simulations and also sensitivity tests with Thompson in WRF gave approximately the same results as with WSM6.



### 3. Methodology and Dataset

This chapter discusses the basic theory of MPAS, the structure of MPAS and its different types of schemes. It also talks about the implementation of MPAS and testing sensitivity of WSM6 and Thompson schemes in simulating TC Dineo over the SWIO. Tracking and detecting TC Dineo over the SWIO basin is also discussed in this chapter.

#### 3.1 Study Area

Our study area is the SWIO basin ( $0 - 40^{\circ}S, 30 - 90^{\circ}E$ ), which is the main tropical cyclone basin in the southern hemisphere with cyclone season starting from 1st July to 30th June but more active between 15th November and 30th April. Cyclones in this basin are called TCs, and are named and classified after reaching 34 knots (10min MSW) near the center. For instance, on 13th February 2017 a TC that formed in the Mozambique channel was named Dineo by NASA, 2017 after reaching 40 knots MSW. Although the highest number of cyclones form between January and March, on average 9-10 tropical systems form in the basin in each season with 4-5 becoming TCs (2018 - South-West Indian Ocean Tropical Cyclones, n.d.). The SWIO basin is monitored by TC Regional specialized Meteorological Centre (RSMC) La Reunion Meteo France, which gives daily updates about TCs during the season. Figure. 3.1 illustrates the track intensity and history of TCs that occurred in the SWIO basin from 2011/2012 to 2017/2018. On the Figure. 3.1, traces in red represent track intensity  $> 110 \text{ km/h}$ , orange  $63 - 110 \text{ km/hr}$ , yellow  $< 63 \text{ km/hr}$  and white represents track intensity for all TCs in SWIO during those seasons.




Figure 3.1: TCs in the SWIO between 2011-2018 (2018 - South-West Indian Ocean Tropical Cyclones, n.d.)

## 3.2 Datasets

The meteorological data (Climate Forecast System Reanalysis (CFSR) (ds094)) used for initializing the model to simulate TC Dineo was downloaded from the Research Data Archive (RDA) website (ds0). CFSR (ds094) dataset was used because TC Dineo occurred between the period of 2011 and 2020. Before downloading the data from RDA website, the period in which TC Dineo occurred was selected starting three days before and ending three days after its occurrence, allowing the model to spin up. On downloading, two datasets, Sea Surface Temperature (SST) and pressure levels, were downloaded from the RDA. SST data sets was extracted for the entire period of occurrence of TC Dineo, i.e., 10th/02/2017 to 20th/02/2017. In contrast pressure levels were extracted for the first six hours of occurrence of TC Dineo, i.e., 13th/02/2017 at 00:00 to 13th/02/2017 at 06:00.

## 3.3 MPAS

MPAS aids atmospheric development and simulation of components of the Earth system which are used in regional climate and weather studies through collaborative projects. MPAS – Atmosphere (MPAS-A) is a non-hydrostatic atmosphere model that is part of MPAS and is led by the National Center for Atmospheric Research (NCAR) Earth System Laboratory (NESL)).

**3.3.1 Model description.** MPAS models use centroidal Voronic tessellations in their horizontal meshes, which provide a common software framework that gives an infrastructure for producing parallel execution, input, output, high-level driver program and other software infrastructure. The supported important features of MPAS - Atmosphere (MPAS-A) include fully-compressible, non-hydrostatic dynamics, split-explicit Runge-Kunta time integration, Exact conservation of dry-air mass and scalar, Positive-definite and monotonic transport options, Generalised terrain-following hight coordinate. Unstructured variable-resolution (horizontal) mesh integrations for the sphere and Cartesian planes is also a supported vital feature of MPAS-A (ucar, n.d.).

Version 6.0 of MPA-A consists of parameterizations of physical processes that are taken from the WRF Model. Explicitly Parameterizations of physical processes supported by MPAS-A include Radiation, Land-surface, Surface-layer, Boundary-layer, Convection, Cloud microphysics (ucar, n.d.). Table. 3.1 shows the set of parameterizations of physical processes and their respective schemes.

Parameterizations	Schemes
Radiation	Community Atmosphere Model (CAM), Rapid Radiative Transfer Model for General Circulation Models (RRTMG) both long and short wave radiation
Land-surface	Noah land-surface model
Boundary-layer	Yonesi University (YSU) and Mellor-Yamada-Nakanishi-Niino (MYNN) Planetary Boundary Layer (PBL)
Convection	Kain-Fritsch Tiedtke, New Tiedtke, and Grell-Freitas convection parameterizations
Cloud microphysics	WSM6, Kessler, and Thompson
Surface-layer	Monin-Obukhov and MYNN

Table 3.1: set of parameterizations and their schemes.

MPAS-A consists of two main components, i.e., the model and initialization, built as cores within the MPAS software framework. Both the model and initialization make use of software infrastructure and the same driver program but they are compiled as separate executables. The model component

consists of physics and atmospheric dynamics. The initialisation component consists of components for creating initial conditions for the atmospheric and land-surface state and updating files for the sea-surface temperature (SST) and sea ice (ucar, n.d.). Figure. 3.2 shows model components within the MPAS framework. As shown in the Figure. 3.2 the `init_atmosphere` (on the left) and `atmosphere` (on the right) are built as separate cores within the framework (MPAS superstructure). Both cores are symbolically linked to the MPAS common infrastructure (at the bottom).




Figure 3.2: The initialization and model components (ucar, n.d.)

**3.3.2 Dynamic core and physics of MPAS.** The effect of physical processes on state variable evolution is modelled or represented using a generic term called Model physics. The effect of these processes is observed either when these processes like radiation, phases of changes of water species are not contained in the Navier-stokes equations or when convective cells within coarse spatial resolution simulations are not resolved on the model mesh. Processes not to be resolved on the model mesh is caused by the failure of the structures to be represented on the mesh ( $L < 2\Delta x$ ), where  $L$  is the length of the grid and  $\Delta x$  grid spacing. Inaccurate representation of present structures on the mesh in the dynamics below the effective resolution also causes physical processes not to be resolved on the model mesh. The physics operates on cell-centered variables. The radial-basis-functions reconstruction and cell-edge normal velocities are used to compute the global mesh and the components of cell-centered velocity (Atmospheric physics).

The Physics in MPAS is divided into several parameterisations: microphysics, surface-layer, land-surface, short wave radiation, long wave radiation, boundary layer, convection, gravity wave drag by orography, and cloud fraction for radiation. Each parameterisation consists of different schemes for example, WSM6, Thompson and Kessler schemes are contained in microphysics, Noah is the only scheme in land-surface whereas Tiedtke, New Tiedtke, Kain-Fritsch, and Grell-Freitas are schemes in convection

parameterization etc.

WSM6 is developed by including graupel ( $q_g$ ) to the Weather Research and Forecasting System (WRF)-Single-Moment-Microphysics Class 5 (WSM5) prognostic variables: snow ( $q_s$ ), cloud ice ( $q_i$ ), cloud water ( $q_c$ ), and mixing ratios of water vapour ( $q_v$ ) (Federico, 2016). WSM5 is an updated version of Weather Research Forecast (WRF)-Single-Moment-Microphysics Class 3 (WSM3) by allowing super-cooled water and mixed phase processes. WSM3 is a simple and efficient scheme that consists of snow and ice processes that makes it suitable for mesoscale grid sizes (NOAA Research). According to the efficiency and theoretical backgrounds of WSM6, it is suitable for both cloud-resolving grids and high-resolution simulations. However, it only anticipates the mass mixing ratios of six water species (Skamarock, 2008). In WSM6, the number of graupel particles per cubic meter of air:  $n_g(D)dD_g(m^{-4})$  follows an exponential size distribution given by:

$$n_g(D)dD_g = n_{og} \exp(-\lambda_g D_g) dD_g \quad (3.3.1)$$

where  $D_g$  is the graupel diameter,  $n_{og}(m^{-4})$  represents the intercept values.  $\lambda_g(m^{-1})$  represents the slope of the exponential distribution. This slope is given by:

$$\lambda_g = \left( \frac{\pi n_{og} \rho_g}{\rho q_g} \right)^{\frac{1}{4}} \quad (3.3.2)$$

where  $\rho_g$  and  $\rho$  represents the graupel and air density, respectively. The slope parameter is obtained by multiplying equation (3.3.1) by the drop mass ( $\frac{\pi \rho_g D_g^3}{6}$ ) and then integrate by parts three times over all diameters, and finally, the resulting quantities are equated to appropriate water contents ( $\rho q_g$ ). The graupel particle terminal velocity of diameter  $D_g$  is given by:

$$V_{Dg} = \alpha_g D_g^{b_g} \left( \frac{\rho_o}{\rho} \right)^{\frac{1}{2}} \quad (3.3.3)$$

where  $\alpha_g$  represents the empirical formula of the mass weighted speed of graupel ( $V_g$ ),  $b_g$  is the empirical formula of the mass weighted speed of snow  $V_r$ , and  $\rho_o$  represents the density of air at reference state. According to Rutledge and Hobbs (1984),  $V_g$  is obtained by integrating equation (3.3.3) and is expressed as:

$$V_g = \frac{\alpha_g \Gamma(4 + b_g)}{6} \left( \frac{\rho_o}{\rho} \right)^{\frac{1}{2}} \frac{1}{\lambda_g^{b_g}} \quad (3.3.4)$$

where  $\Gamma(\gamma)$  is a gamma function. The continuity equation for graupel is given by:

$$\frac{\partial q_g}{\partial t} = -\vec{V} \cdot \nabla q_g - \frac{q_g}{\rho} \frac{\partial(\rho V_g)}{\partial z} + S_g \quad (3.3.5)$$

where  $S_g$  represents the sinks and sources of graupel.  $\vec{V} \cdot \nabla q_g$  and  $\frac{q_g}{\rho} \frac{\partial(\rho V_g)}{\partial z}$  represent the 3D advection and sedimentation of graupel respectively.

Thompson is a bulk microphysical parametrization scheme whose assumed snow size distribution depends on both ice water content and temperature. It consists of number concentration with six classes of moisture species as its prognostic variables: cloud water, graupel, cloud ice, rain, snow, and water vapour (NOAA Research). Thompson explicitly predicts cloud ice and droplet number concentration due to its inclusion of aerosols as its cloud condensation nuclei (CNN) and ice nuclei (IN). In this scheme, snow size distribution ( $N(D)$ ) is represented by the sum of gamma and exponential distributions, as

illustrated in equation (3.3.6), and snow assumes a non-spherical shape with a bulk density that varies inversely with its diameter (Skamarock, 2008).

$$N(D) = \frac{M_2^4}{M_3^3} \left[ K_o \exp \left( -\frac{M_2}{M_3} \wedge_o D \right) + K_1 \left( \frac{M_2}{M_3} D \right)^{\mu_s} \exp \left( -\frac{M_2}{M_3} \wedge_1 D \right) \right] \quad (3.3.6)$$

where  $K_o$ ,  $K_1$ ,  $\wedge_o$  and  $\wedge_1$  are snow size distribution constants.  $\mu_s$  represents the gamma size distribution shape parameter.  $M_n$  is the distribution  $n^{th}$  moment:  $M_n = \int D^n N(D) dD$ .  $M_n$  allows bulk snow density to be correctly modelled with an inverse diameter dependence in contrast to WSM6 which assumes constant snow density. The first and second term inside the brackets of equation (3.3.6) represents the exponential and gamma distributions, respectively. The sum of these two distributions accounts for the general slope of large particles and frequently observed super-exponential number of small particles. Through the use of equation (3.3.6) in the scheme enables the effective  $y$  intercept parameter to depend on both snow mixing ratios and temperature. The number concentrations (mass mixing ratios) of all forms of water species: vapour, liquid, and solid or aerosol number concentrations follow the same dominant conservation given by:

$$\frac{\partial \Phi}{\partial t} = -\frac{1}{\rho} \nabla \cdot (\rho U \Phi) - \frac{1}{\rho} \frac{\partial (\rho V_\Phi \Phi)}{\partial z} + \delta \Phi + S_\Phi \quad (3.3.7)$$

where  $\Phi$  represents the mass mixing ratio of any water species,  $U$  is the 3D wind vector,  $t$  is time,  $V_\Phi$  is the weighted fall speed of  $\Phi$ ,  $\rho$  is air density,  $z$  is height,  $\delta \Phi$  is the sub-grid scale mixing operator,  $S_\Phi$  is the microphysical process rate terms. The detailed description for newly predicted variables: number of Hygroscopic aerosol ( $N_{wfa}$ ), number of Non-hygroscopic aerosol ( $N_{ifa}$ ) i.e., ice-nucleating aerosol and number concentration of cloud droplet ( $N_c$ ) are provided respectively in equations (3.3.8) to (3.3.10).

$$\begin{aligned} \frac{dN_{wfa}}{dt} = & - \left( \text{rain, snow and graupel} \right) - \left( \text{homogeneous nucleated} \right) - \left( \text{Cloud Condensation} \right) \\ & + \left( \text{cloud and rain} \right) + \left( \text{surface emissions} \right) \\ & \text{collecting aerosols} \quad \text{deliquesced aerosols} \quad \text{Nuclei (CCN) activation} \end{aligned} \quad (3.3.8)$$

$$\frac{dN_{ifa}}{dt} = - \left( \text{rain, snow and graupel} \right) - \left( \text{Ice Nuclei (IN)} \right) + \left( \text{cloud ice} \right) + \left( \text{surface emissions} \right) \quad (3.3.9)$$

$$\text{collecting aerosols} \quad \text{activation} \quad \text{sublimation}$$

$$\begin{aligned} \frac{dN_c}{dt} = & - \left( \text{rain, snow and graupel} \right) - \left( \text{freezing into} \right) - \left( \text{collide into rain} \right) - \left( \text{evaporation} \right) + \left( \text{CCN activation} \right) \\ & + \left( \text{cloud ice melting} \right) \\ & \text{collecting droplets} \quad \text{cloud ice} \end{aligned} \quad (3.3.10)$$

Thompson assumes that water and ice species follows a generalised gamma distribution defined in equation (3.3.11) in instead of the exponential distribution.

$$N(D) = \frac{N_t}{\Gamma(\mu + 1)} \lambda^{\mu+1} D^\mu \exp(-\lambda D) \quad (3.3.11)$$

where  $\lambda$  represents the slope of the distribution,  $N_t$  gives the total number of particles in the distribution,  $D$  is the diameter of the particles, and  $\mu$  is the shape parameter of the distribution. All water and ice species are assumed

to use a standard power law form in relating their mass and terminal velocity as a function of diameter. The mass ( $m(D)$ ) and terminal velocity ( $V(D)$ ) relations are given by equations (3.3.12) and (3.3.13) respectively.

$$m(D) = aD^b \quad (3.3.12)$$

$$V(D) = \left(\frac{\rho_o}{\rho}\right)^{\frac{1}{2}} \alpha D^\beta \exp(-fD) \quad (3.3.13)$$

where  $a$  and  $b$  are mass power law constants and these vary for different species,  $f$ ,  $\beta$ , and  $\alpha$  are velocity power law constants,  $\rho_o$  and  $\rho$  are the reference and moist air densities respectively.

### 3.4 Model simulation of TC Dineo

The following steps were taken to effect model simulation: (1). We created a run directory which we named `case_studies` on the CHPC cluster. (2). Linked the `init_atmosphere_model` and `atmosphere_model` executables, physics lookup tables into the `case_studies` directory. (3). Copied the `namelist.*`, `streams.*`, files into the `case_studies` directory. (4). Downloaded 92km-25km variable resolution mesh (`x4.163842.grid.nc`) from <https://mpas-dev.github.io>, then push it to the CHPC account in the `grid_rotate` directory. (5). We used the `grid_rotate` utility by editing the `namelist.input` file to set the latitudes and longitudes to the center of interest, then use the `grid_rotate` utility to rotate the mesh to the new area of refinement (Figure 3.3), hence producing a new netCDF file used as an input in the `streamlist` file during static and terrestrial field processing. (6). Push both the downloaded datasets (SST and pressure levels) from the RDA website to the chpc server account into the `sst` and `press` directory respectively, run the executable file `ungrib.exe` to generate intermediate files for both sea surface temperature (SST) and pressure levels. (7). Configured the `namelist` and the `stream` files in the `init_atmosphere` directory, then run the `init_atmosphere_model` to create initial conditions. (8). Generate idealized initial conditions by interpolating the produced intermediate files with the land surface initial conditions, the created file is used as input in the `streams.atmosphere` file during model integration. Model integration simulation was simulated twice, for WSM6 and Thompson schemes. In the `namelist.atmosphere`, the `&physics` section gives as ability to choose our own schemes by changing the `namelist` variables during model integration. So this gave as chance to perform two simulations i.e., one with WSM6 with other schemes available and Thompson excluded, and finally Thompson with other schemes available and WSM6 excluded. For the case of this study, since we are using 25km variable resolution and we are interested in comparing the capability of WSM6 and Thompson microphysics schemes in simulating TC Dineo over SWIO, we used the `mesoscale_reference` suit, and override its default after the first simulation.

`Mesoscale_reference` suit contains WSM6 scheme as its default as shown in table. 3.2, so for the first simulations with WSM6, we don't adjust anything in the `&physics` record in the `namelist.atmosphere` file. For the second simulations, we over-ride the default and change WSM6 to Thompson scheme adding a `namelist` variable (`config_microp_scheme = 'mp_thompson'`) in the `&physics` record. This enabled the `atmosphere_model` to read Thompson scheme during integration instead of WSM6. Before running the `atmosphere_model`, we first run `build_tables` executable file to generate lookup tables for the Thompson scheme, since they were not generated initially. This is because the look up tables contain the essential Thompson physics variables necessary for model integration while running with Thompson scheme. Table. 3.2 illustrates default parameterization and schemes contained in the `mesoscale_reference` suit which have been tested for mesoscale resolution.



Parameterizations	Schemes
Convection New	Tiedtke
Microphysics	WSM6
Land surface	Noah
Boundary layer	YSU
Surface layer	Monin-Obukhov
Radiation, LW	RRTMG
Radiation, SW	RRTMG
Cloud fraction for radiation	Xu-Randall
Gravity wave drag by orography	YSU

Table 3.2: Set of parameterization schemes used by mesoscale\_reference suit

Having finished configuring the namelist.atmosphere file, the atmosphere\_model program produced history and diagnostic files for the entire period of simulation which were remapped using the convert\_mpas utility to produce a netCDF file name latlon.nc which was visualized and used to make analysis.

The Figure. 3.3 shows the new transformed mesh, Dineo.grid.nc, to our new area of interest rotated about 90° using the grid\_rotate utility.




Figure 3.3: After the grid\_rotate

### 3.5 Detecting and Tracking TC Dineo

After diagnostic and history files were generated during model integration as explained in section 3.4 for the two schemes: WSM6 and Thompson, convert\_utility were used to generate a latlon file which contained mean sea level pressure (mslp), zonal wind component (u10), meridional wind component (v10), temperature\_250hpa, temperature\_500hpa and vorticity\_850hpa variables. All these variables were created only for the domain in which TC Dineo occurred, this simplified the tracking because we dealt with a small data set.

The objective measures that we used for detecting and tracking TC Dineo in the tracking algorithm that uses a 6 hourly output data in the latlon file are similar to that described by Kleppek et al. (2008): (1). The minimum

value of mslp over the entire domain termed as the centre of a low pressure system (TC) was determined. (2).  $2^\circ$  from the centre of the TC, the warm core which is the average integrated temperature between 250 hpa and 500 hpa was calculated, this core is at least  $1^\circ C$  warmer than the surrounding environment which is defined as a  $10^\circ \times 10^\circ$  block centred at the mslp. (3). The wind speed which is used to classify the intensity of TC Dineo according to Saffir Simpson scale was calculated using u10 and v10 components over the domain. Then we searched for the maximum value of the wind speed within  $5^\circ$  of the cyclone centre. (4). The minimum vorticity within  $4^\circ$  of the cyclone centre is determined. Since we are in the SWIO basin, negative vorticity is associated with a cyclone spin.

TC Dineo was detected by comparing the calculated variables above with the cyclone threshold values as follows: (1). The maximum surface wind speed within  $5^\circ$  of the cyclone centre must be greater than  $17 ms^{-1}$ . (2). The minimum vorticity must be less than  $3.5 \times 10^{-5} s^{-1}$ . (3). The warm core temperature must be at least  $1^\circ C$  greater than the surrounding environment. If the cyclone we identified satisfied the above TC criteria, it was assigned a tracking ID of 1 otherwise 0 for the respective 6 hourly intervals during each day TC Dineo occurred.

The output of the respective computed variables: Date, mslp, latitude, longitude, wind speed, vorticity and warm core temperature with their respective track IDs were visualized for the respective schemes to view the track path of TC Dineo as shown in Figure. 4.1.



## 4. Results and discussions

This chapter discusses all the results obtained from the methodology described in chapter 3, and it is divided into three sections. The first section gives the description and analysis for the track of TC Dineo over the SWIO basin. The time evolution of different variables are displayed and discussed in the same chapter. The second and third chapter discusses the horizontal and vertical structures, respectively of TC Dineo for reanalysis, MPAS (Thompson) and MPAS (WSM6).

### 4.1 Track of TC Dineo

As far as we know, the capability of MPAS microphysics schemes (WSM6 and Thompson) in tracking TCs over SWIO has not been investigated yet. Therefore we started the project by comparing various tracks from four different data sets: reanalysis (red), observation (green), WSM6 (blue), and Thompson (orange) as shown in Figure 4.1 for the entire lifetime of cyclone Dineo. The small triangles along the trajectories of the various data sets with their respective colours indicate the timing at which TC Dineo matured from being a tropical storm to a category one tropical cyclone.




Figure 4.1: Track of TC Dineo over SWIO depicted by four different datasets (observations, reanalysis, MPAS (Thompson), and MPAS(WSM6)).

On the Figure 4.1, over water (  $19^{\circ}$ - $24^{\circ}$  S;  $35^{\circ}$  -  $42^{\circ}$  E), WSM6 and Thompson almost followed the same path as the observation and the reanalysis data, though the tracks started at different time intervals. At around ( $35^{\circ}$  E,  $23.5^{\circ}$  S), before TC Dineo made landfall over Inhambane (southern Mozambique), it was guided by an easterly steering flow over Mozambique channel induced by a 500 mb high pressure system that prevailed over most

southern Africa during the entire lifetime of TC Dineo (Moses and Ramotonto, 2018). At this location, all the tracks coincided, which means that the simulated data from MPAS, reanalysis and the observation data sets were all consistent in tracing the position of TC Dineo at that location.

From the same Figure 4.1, it can also be seen that Thompson consistently followed the same track as that of reanalysis over land, which indicated a much better performance than WSM6 in tracking the entire path of TC Dineo. From the Mozambique coast to eastern Botswana, the observation and WSM6 trajectories are positively correlated. However, the track from the simulated data set (WSM6) underestimated the track of TC Dineo as compared to the observation. Over the northern part of South Africa, WSM6 deviated a lot from the three data sets tracks, but all the tracks ended in eastern Botswana. It means that TC Dineo weakened and dissipated in the eastern part of Botswana. It is consistent with observations from Moses and Ramotonto (2018) that TC Dineo weakened to remnant low before hitting Botswana and dissipated on the same day after striking Botswana.

There is a consistency between the spatial distribution (Figure 4.1) of TC Dineo track and the time evolution of MSLP (Figure 4.2a). Looking at these two figures all data sets started at different periods and followed different trajectories as TC Dineo dragged southwards, however, on 15<sup>th</sup> February, all the data sets tracked the cyclone in different locations at different times. Furthermore, all data sets depicted that TC Dineo dissipated in Botswana on the same day (17<sup>th</sup> February) but in different locations. This is consistent with observations from Moses and Ramotonto (2018).

Generally, both MPAS (Thompson) and MPAS (WSM6) were able to produce a track that resembled the observations, however, Thompson track more closely resembled observations.

Figure 4.2a shows the time evolution of mean sea level pressure for the entire period of cyclone Dineo for four datasets: ERA5 reanalysis (red), RSMC observation (green), WSM6 (blue), and Thompson (orange). The data was obtained on a six-hourly interval on each TC Dineo occurred. TC Dineo reached its peak on 15<sup>th</sup> February and this is depicted by both observation and reanalysis data. Both WSM6 and Thompson almost underestimated the day at which TC Dineo matured because they are positively correlated, and their minimum point exits towards the end of 14<sup>th</sup>. The two MPAS schemes overestimated the peak intensity of MSLP compared to the observation and reanalysis. Looking at the trend of all the data sets, they are all gradually decreasing and then increasing again. This is so because TCs are low pressure systems, therefore during their entire lifetime, the pressure of the eye keeps on dropping until it matures and then increases again, making its sustainable conditions unfavourable hence dissipating as shown in Figure 4.2a.

Figure 4.2b illustrates the time evolution of wind speed. All the data set curves are increasing to their maximum and then decrease. This is because when a cyclone is growing, the winds in the eyewall keep on increasing rapidly to the maximum as the cyclone matures. On 14<sup>th</sup> at around 18:00 tending to 15<sup>th</sup>, WSM6 and Thompson had their maximum wind speeds, which implies that the storm matured by then, which is consistent with the results in Figure 4.2a, because at that same time is when WSM6 and Thompson had the lowest values of pressure. The observation data curve reached category 1, but the rest of other data sets remained in the tropical storm region. This implies that both MPAS schemes underestimated the peak intensity of WSPD in comparison with the observation data set. This may be due to terrain settings during model initialization which affected the magnitude of the wind speed for the two schemes.

Both in Figure 4.2a and 4.2b, the observation reached category one on the same day (15<sup>th</sup> February) and dropped back to tropical storm on the same day, however, the rest of the data sets remained tropical storms during the entire period which is consistent in both figures. This depicts a strong inverse relationship between MSLP and WSPD.

Figure 4.2c shows the wind pressure relationship. The plot illustrates that the data sets are well correlated to each other, and it depicts a strong relationship between WSPD and MSLP for all the data sets which is consistent with Figure 4.2a and 4.2b. This is because the trend in the two curves is opposite to each other, furthermore, the pressure in the eye is always very low, and the eyewall's wind speed is always at maximum. This relationship affected by several factors: TC size, latitude, environmental pressure and the intensification trend. For all the data sets there is a high concentration of the scatter points between 980-1010 hPa, this is because as the cyclone dragged southwards, the latitude kept on decreasing which decreased the Coriolis force, therefore requiring a

lesser pressure gradient force to counter balance the tangential wind. The WSM6, Thompson and reanalysis data distributions are almost all coinciding, this means that these data sets follow the same trend.

Figure 4.2d shows the time evolution of the total precipitation for the entire lifetime of TC Dineo, which formed on 11<sup>th</sup> February 2017. The total precipitation which is the sum of the accumulated convective and grid scale precipitation is obtained from four data sets. For the first five days, all the data sets almost produced the same total rainfall, however starting on 15<sup>th</sup> when TC Dineo reached its peak, the following days, the amount of rainfall that was produced was severe. WSM6 produced much rainfall on 16<sup>th</sup> at around mid day followed by Thompson, then reanalysis and finally the observation data set. Between 13<sup>th</sup> at 18:00 to 15<sup>th</sup> at around noon, WSM6 and Thompson are positively correlated. Moses and Ramotonto (2018) claimed that the maximum observed rainfall (mm) recorded on 16<sup>th</sup> by 12 stations was 17 mm, on 17<sup>th</sup> was 270 mm by 66 stations and on 18<sup>th</sup> was 55.5 mm from 49 stations. This claim is consistent with predictions produced by WSM6. Generally, WSM6 simulated the total precipitation better than Thompson, even though both schemes over estimated the total rainfall in comparison with the observation data from 15<sup>th</sup> till when TC Dineo dissipated (17<sup>th</sup>).




Figure 4.2: Time evolution of mean sea level pressure (MSLP), wind speed (WSPD), and Precipitation simulated by reanalysis, observation, Thompson, and WSM6 are depicted by (a), (b), and (d) respectively. (c) depicts the wind speed pressure relationship for the four data sets.

Figures 4.3a and 4.3b describe the average sea surface temperature (SST) for the entire period of TC Dineo for both Thompson and WSM6 respectively. According to the spatial plots display in Figure 4.3a and 4.3b, the SST (dark red) over the domain where TC Dineo formed and matured, is 302.1 K. This is a favourable condition for TC to develop, therefore both schemes: Thompson and WSM6 recognized SST data input during simulation. On both figures, near Inhambane were TC Dineo made landfall on 15<sup>th</sup>, both schemes estimated the average temperature of seawater that was necessary for TC Dineo to remain sustained.




Figure 4.3: Sea surface temperature (SST) for the entire period of TC Dineo simulated by the two data sets. (a) : MPAS (Thompson), (b) : MPAS (WSM6).

Figures 4.4a and 4.4b show the total accumulated precipitation for both MPAS (Thompson) and MPAS (WSM6), respectively, for the entire lifetime of TC Dineo. The total precipitation was obtained by summing the total precipitation produced by both microphysics (accumulated grid precipitation) and convective (accumulated convective precipitation) MPAS schemes. As observed from Figures 4.5a, 4.5b and 4.5c, TC Dineo was on its way to make landfall over Inhambane (southern Mozambique) and this is when it reached its peak strength (category 1) as shown in Figure 4.2b. It is clear from the graph that before this day both Thompson and WSM6 produced much rainfall between the southwest of Madagascar and Mozambique. However, Thompson consistently produced much rainfall as TC Dineo approached the Inhambane coast compared to WSM6. Even though the total rainfall simulated by Thompson was much more than WSM6 over the entire domain, WSM6 produced much higher rainfall in Inhambane after when TC Dineo had moved overland. It is consistent with the results obtained from the time series plot (Figure 4.2d). Over Madagascar, Thompson produced much more rainfall compared to WSM6. Moses and Ramotonto (2018) claimed that TC Dineo weakened to remnant low before hitting Botswana, and that this weakened storm triggered heavy rainfall over eastern Botswana. This claim contradicts with the simulations obtained from both schemes. However, Thompson produced a lot more rainfall in northern Botswana compared to WSM6.




Figure 4.4: Total Precipitation for the entire period of TC Dineo simulated by the two data sets. (a) : MPAS (Thompson), (b) : MPAS (WSM6).

## 4.2 Horizontal structure of TC Dineo

This section presents the horizontal structure of TC Dineo on 15<sup>th</sup> February 2017 (day when TC Dineo matured) for WSPD and MSLP obtained from three data sets: Reanalysis, MPAS (Thompson) and MPAS (WSM6).

Figures 4.5a, 4.5b and 4.5c illustrates the horizontal structure for the wind speed for the peak date of TC Dineo. The figures show a cyclonic flow around the eye (almost enclosed by the dark red region) with a high concentration in the east (dark red region) compared to the west. This is due to the converging of the cyclonic flow, the background flow in the eastern region and opposition of the background flow by the cyclonic flow in the western region as the cyclone drags southwards. This is depicted by the vector fields overlaid on the structures. In comparison to the reanalysis structure, both Thompson and WSM6 underestimated the location of the TC Dineo on reaching its peak. The reanalysis structure depicts that the cyclone was just away to make landfall over Inhambane with a maximum wind speed of 26.19 m/s. However, both Thompson and WSM6 simulated that TC Dineo was still a distance away from the coast, even though the maximum wind speed simulated by Thompson (24.17 m/s) is greater than of WSM6 (24.10 m/s). The eyewall (red dark region around the eye) for reanalysis structure is much thicker than that of both Thompson and WSM6 however Thompson's eyewall is thicker than that of WSM6, hence there is a negligible difference between Thompson and WSM6 in simulating the horizontal structure of the wind speed.




Figure 4.5: Horizontal structure of the wind speed (WSPD) for TC Dineo on its peak day (15<sup>th</sup> February 2017) for three data sets. (a) : Reanalysis, (b) : MPAS (Thompson), (c) : MPAS (WSM6).

Figures 4.6a, 4.6b and 4.6c represents the horizontal distribution of the MSLP in the eye (central purple colour) for three datasets: reanalysis, Thompson, and WSM6 for the peak date of TC Dineo. All the structures were able to predict the lowest pressure centre (eye). The eye in reanalysis is at a pressure of 98900 Pa, and 100200 Pa for both Thompson and WSM6. This implies that both simulated data sets: Thompson and WSM6 overestimated the minimum sea-level pressure. The position of the eye away from the coast is much shorter for reanalysis compared to Thompson and WSM6 even though Thompson is much at a closer distance compared WSM6. These results are consistent with results obtained in figures 4.5a, 4.5b, and 4.5c for the WSPD. The horizontal ascending increase of the MSLP from the eye to other parts of the cyclone is observed in all the figures, consistent with Roy and Kovordányi (2012) literature on the anatomy of TCs.




Figure 4.6: Horizontal structure of the mean sea level pressure (MSLP) for TC Dineo on its peak day (15<sup>th</sup> February 2017) for three data sets. (a) : Reanalysis, (b) : MPAS (Thompson), (c) : MPAS (WSM6).

### 4.3 Vertical cross section of TC Dineo

In this section, the vertical distribution of vorticity, warm core and WSPD on the day when TC Dineo reached its peak are discussed for three data sets: reanalysis, MPAS (Thompson) and MPAS (WSM6).



Based on the fact that TCs are warm core systems, Figures 4.7a, 4.7b, and 4.7c illustrate the vertical structure of temperature of TC Dineo for three datasets. The adiabatic (compressional warming) heat releases of subsiding air causes the eye to be a warm core temperature. This is depicted in all temperature distributions for the three data sets however, it can be seen that it is realistic in the reanalysis distribution compared to that in Thompson and WSM6. In all distributions, warmer temperatures (dark red) in the upper levels and lower in the lower levels are exhibited, this is consistent with the literature about the characteristics of the eye (Roy and Kovordányi, 2012) hence all data sets were able to simulate temperatures of TC Dineo aloft. However the distributions in Thompson and WSM6 slightly differ from that in reanalysis, this means that Thompson and WSM6 exhibited a skill in simulating the temperature structure of TC Dineo.




Figure 4.7: Vertical structure of the warm core temperature for TC Dineo on its peak day (15<sup>th</sup> February 2017) for three data sets. (a) : Reanalysis, (b) : MPAS (Thompson), (c) : MPAS (WSM6).

Figures 4.8a, 4.8b, and 4.8c show the vertical distribution of relative vorticity of TC Dineo on the 15<sup>th</sup> for three data sets. In all the distributions, relative vorticity in the centre of TC Dineo was well simulated with large negative values at lower levels that slowly increase to smaller values at upper levels. However, the feature is exhibited more in reanalysis and Thompson distributions compared to WSM6. Since TC Dineo formed in the southern hemisphere, negative low-level vorticity is associated with the spinning of a cyclonic rotating cyclone. Reanalysis show a wider area  $\approx 1^\circ$  of large values of negative vorticity compared Thompson and WSM6 at lower levels. However, Thompson  $\approx 0.5^\circ$  predicted a bigger value compared to WSM6  $\approx 0.2^\circ$ , hence Thompson simulated surface convergence better than WSM6, which is a condition favorable for a cyclone to be sustained. This is consistent with the intercontinental convergence zone (ITCZ) because large negative values cause surface convergence of easterly waves in the southern hemisphere. In all the relative vorticity distributions for all data sets, large areas (red) with positive values of vorticity depict the Southern Indian high pressure system regions (Abiodun, 2016). Therefore such regions with positive vorticity are associated with subsidence, which often constrains convective precipitation, therefore affecting results in Figures 4.4a and 4.4b (33 – 40°E).






Figure 4.8: Vertical structure of Vorticity for TC Dineo on its peak day (15<sup>th</sup> February 2017) for three data sets. (a) : Reanalysis, (b) : MPAS (Thompson), (c) : MPAS (WSM6).

The Wind speed of TC Dineo on the 15<sup>th</sup> (the day it matured) for three data sets: reanalysis, Thompson and WSM6 is shown in Figures 4.9a, 4.9b, and 4.9c. The actual structure of a TC is depicted in Figure 4.9a with a calm eye surrounded by an eyewall with a maximum sustained wind speed of approximately 14 m/s. The wind speed is higher in the east than in the west due to the combination of cyclonic flow, the background flow in the east and cancellation of these two flows in the west, as well as the fact that TC Dineo formed in the southern hemisphere. The two schemes failed to depict the actual structure of TC Dineo at its mature stage. However, both schemes simulated the central and eastern distribution of the wind speed even though the distribution simulated in WSM6 shows similar qualities to that depicted in the reanalysis, i.e., in both distributions, the eyewall diameter exhibits the same characteristics with  $\approx 0.2^\circ$  at the lower level and gradually increases to  $\approx 0.5^\circ$  in the upper levels. Furthermore, the centre of the eye is almost at the same position ( $37^\circ\text{E}$ ) in both reanalysis and WSM6 distributions. The eye structure in the WSM6 and reanalysis distributions are clear than that of Thompson (faint), though all are almost at the same position. Therefore WSM6 simulated the distribution better than Thompson. However, the storm is deeper and stronger in the east at lower levels in all data sets even though thicker for both WSM6 and Thompson compared to the reanalysis.




Figure 4.9: Vertical structure of the wind speed (WSPD) for TC Dineo on its peak day (15<sup>th</sup> February 2017) for three data sets. (a) : Reanalysis, (b) : MPAS (Thompson), (c) : MPAS (WSM6).

## 5. Conclusion

The study compared the capability of two MPAS microphysics Schemes: WSM6 and Thompson in simulating TCs over the SWIO basin. The simulation was performed over nine days (11<sup>th</sup>/02/2017 - 19<sup>th</sup>/02/2017) for TC Dineo (case study). The model simulation was compared to 6 hourly ERA5 reanalysis and Regional Specialised meteorological Centre (RSMC) observational data to assess the schemes' capability to simulate the spatial and temporal variations of TCs over the SWIO basin. TCs were detected and tracked using an unbiased tracking algorithm.

Thompson and WSM6 appropriately reproduced the fundamental dynamical structure of TC Dineo, which comprised of a warm core and a well defined eye ringed by cyclonic winds that vanish gradually with increasing distance from the centre. A similar structure was simulated by reanalysis with more advanced details in most aspects.

The reanalysis and the schemes cloned well the spatial and temporal distribution of TC Dineo track, eye, WSPD, warm core and vorticity over the SWIO basin. Thompson performed slightly better than WSM6 in almost all distributions in comparison to reanalysis. However, WSM6 did well in simulating the vertical cross section of the WSPD compared to Thompson. Overall, MPAS (Thompson) illustrated good approximations with the Observations and reanalysis in both spatial and temporal variations in simulating TCs over the SWIO basin for the study period.

Both WSM6 and Thompson cloned well the daily variation of TC Dineo although, in comparison to reanalysis and observation they underestimated the MSLP and WSPD on the peak day (15<sup>th</sup> of February). The two schemes both failed to represent the lowest MSLP of  $\approx 964$  hPa in the observation. However, in comparison with the reanalysis  $\approx 989.02$  hPa, Thompson  $\approx 1001.06$  hPa simulated MSLP better than WSM6  $\approx 1001.24$  hPa.

The result from this study depicted that Thompson has assurance for simulating TCs over the SWIO basin compared to WSM6. However, further studies will intend to improve the performance of the schemes in predicting TCs over the basin. The first development would be using a higher resolution (3km) rather than 25 km variable resolution in conducting the simulations, in order to compare the impacts of grid resolution while using the two schemes in simulating TCs over the basin. Furthermore, various parameterization schemes coupled with Thompson and WSM6 in the same suit for 3km (convective permitting suit) and 25km (mesoscale reference suit) will also be investigated. Since different coupled parameterizations of different schemes in different suits, impacts TC simulation differently. The second improvement would be comparing the capability of all the three microphysics schemes: WSM6 (default), Thompson and Kessler with two different resolutions: 3km and 25km in simulating TCs over the SWIO basin.

# Acknowledgements

I cannot express enough my thanks to the almighty God for aa successful completion of my project. I offer my sincere appreciation to Prof. Babatunde J. Abiodun, my project supervisor, for his guidance, mentoring, supportive idea, and encouragement. Because of his support I was able to complete my project successfully.

Special thanks goes to Dr. Stive Roussel Tankio Djio kap, my personal tutor for his guidance, inspirations, mentorship, suggestions, and useful criticisms which has helped me improve my work and Dr. Roland Ridick Takong who helped me with the computing facility on the Center for High Performance Computing (CHPC South Africa).

Furthermore I would like to thank my project colleagues (Romain, Yao, Remy) for putting ideas together during model simulations and Miss. Grace Aborah Adu for taking time to read my work and giving helpful suggestions. Last but not least I would like to thank AIMS community.

# References

- 2018 - South-West Indian Ocean Tropical Cyclones, n.d. 2018 - south-west indian ocean tropical cyclones. EUROPEAN COMMISSION JOINT RESEARCH CENTRE, <https://www.gdacs.org/Public/download.aspx?type=DC&id=161>, Accessed April 2020.
- Abiodun, B. Simulating the characteristics of tropical cyclones over the south west indian ocean using a stretched-grid global climate model. 2016. URL <https://www.ecmwf.int/node/16707>.
- Atmospheric physics. Mpas-atmosphere model user's guide. www2.mmm.ucar.edu , [https://www2.mmm.ucar.edu/projects/mpas/mpas\\_atmosphere\\_users\\_guide\\_7.0.pdf](https://www2.mmm.ucar.edu/projects/mpas/mpas_atmosphere_users_guide_7.0.pdf), Accessed April 2020.
- Cyclone Dineo. (2020, 3 15). Cyclone\_dineo. En.wikipedia.org, [https://en.wikipedia.org/wiki/Cyclone\\_Dineo](https://en.wikipedia.org/wiki/Cyclone_Dineo), Accessed April 2020.
- Doocy, S., Daniels, A., Packer, C., Dick, A., and Kirsch, T. D. The human impact of earthquakes: a historical review of events 1980-2009 and systematic literature review. *PLoS currents*, 5, 2013.
- Federico, S. Implementation of the wsm5 and wsm6 single moment microphysics scheme into the rams model: verification for the hymex-sop1. *Advances in Meteorology*, 2016, 2016.
- Fitchett, J. M. and Grab, S. W. A 66-year tropical cyclone record for south-east africa: temporal trends in a global context. *International Journal of Climatology*, 34(13):3604–3615, 2014.
- Harris, L. M. and Lin, S.-J. Global-to-regional nested grid climate simulations in the gfdl high resolution atmospheric model. *Journal of climate*, 27(13):4890–4910, 2014.
- Hashimoto, A., Done, J. M., Fowler, L. D., and Bruyère, C. L. Tropical cyclone activity in nested regional and global grid-refined simulations. *Climate Dynamics*, 47(1-2):497–508, 2016.
- Hong, S.-Y. Hongandlim-jkms-2006. *Journal of the Korean Meteorological Society*, 42:129–151, 06 2006.
- Hong, S.-Y., Kim, J.-H., Lim, J.-o., and Dudhia, J. The wrf single moment microphysics scheme (wsm). *Journal of the Korean Meteorological Society*, 42:129–151, 03 2006.
- JetStream. Tropical cyclone introduction. National Weather Service, <https://www.weather.gov/jetstream/tc>, Accessed April 2020.
- Joseph A. Zehnder. Tropical cyclone. Encyclopædia Britannica, <https://www.britannica.com/science/tropical-cyclone>, Accessed April 2020.
- Kleppek, S., Muccione, V., Raible, C., Bresch, D., Köllner-Heck, P., Stocker, T., Muccione, C., Raible, D., Bresch, P., Koellner-Heck, T., and Stocker. Tropical cyclones in era-40: A detection and tracking method. *Geophys. Res. Lett*, 35, 01 2008.
- Mavume, A. F., Rydberg, L., Rouault, M., and Lutjeharms, J. R. Climatology and landfall of tropical cyclones in the south-west indian ocean. *Western Indian Ocean Journal of Marine Science*, 8(1), 2009.
- Moses, O. and Ramotonto, S. Assessing forecasting models on prediction of the tropical cyclone dineo and the associated rainfall over botswana. *Weather and climate extremes*, 21:102–109, 2018.
- Mozambique. (n.d.). Tropical cyclone dineo hits mozambique. ReliefWeb, <https://reliefweb.int/report/mozambique/tropical-cyclone-dineo-hits-mozambique>, Accessed April 2020.
- MPAS. (2020). Model\_for\_prediction\_across\_scales. Mpas-dev.github.io, <https://mpas-dev.github.io/>, Accessed April 2020.

- NASA, 2017. Nasa examines ex-tropical cyclone dineo's rainfall. NASA, n.d., <https://www.nasa.gov/feature/goddard/2017/dineo-southern-indian-ocean>, 2017.
- Natural Disasters. Tropical cyclones. Tulane University, [https://www.tulane.edu/~sanelson/Natural\\_Disasters/tropical\\_cyclones.htm](https://www.tulane.edu/~sanelson/Natural_Disasters/tropical_cyclones.htm), Accessed April 2020.
- ncar.ucar.edu. (n.d.). Model\_for\_prediction\_across\_scales. wikipedia, [https://en.m.wikipedia.org/wiki/Model\\_for\\_Prediction\\_Across\\_Scales](https://en.m.wikipedia.org/wiki/Model_for_Prediction_Across_Scales), Accessed April 2020.
- NOAA Research. Wrf namelist.input file description. National Oceanic & Atmospheric Administration , [https://esrl.noaa.gov/gsd/wrfportal/namelist\\_input\\_options.html](https://esrl.noaa.gov/gsd/wrfportal/namelist_input_options.html), Accessed April 2020.
- Roy, C. and Kovordányi, R. Tropical cyclone track forecasting techniques-a review. *Atmospheric research*, 104: 40–69, 2012.
- Rutledge, S. and Hobbs, P. The mesoscale and microscale structure and organization of clouds and precipitation in midlatitude cyclones. xii: A diagnostic modeling study of precipitation development in narrow cold-frontal rainbands. *Journal of The Atmospheric Sciences - J ATMOS SCI*, 41:2949–2972, 10 1984. doi: 10.1175/1520-0469(1984)041<2949:TMAMSA>2.0.CO;2.
- Sciences, E. (n.d.). Nasa catches formation of tropical cyclone dineo. Phys.org, <https://phys.org/news/2017-02-nasa-formation-tropical-cyclone-dineo.html>, Accessed April 2020.
- Skamarock, e. a. A description of the advanced research wrf version 3, 2008. NCAR TECHNICAL NOTE.
- Su, L. and Fung, J. C. Investigating the role of dust in ice nucleation within clouds and further effects on the regional weather system over east asia–part 1: model development and validation. *Atmospheric Chemistry and Physics*, 18(12):8707–8725, 2018.
- ThoughtCo. The anatomy of a hurricane. ThoughtCo, <https://www.thoughtco.com/anatomy-of-a-hurricane-3443962>, Accessed April 2020.
- TROPICAL CYCLONES. Technical hazard sheet - natural disaster profiles. World Health Organization, [https://www.who.int/hac/techguidance/ems/tropical\\_cyclones/en/](https://www.who.int/hac/techguidance/ems/tropical_cyclones/en/), Accessed April 2020.
- Tropical cyclones in the South West Indian Ocean. Tropical cyclones in the south west indian ocean. ReliefWeb, <https://reliefweb.int/report/world/tropical-cyclones-south-west-indian-ocean-new-insights>, Accessed April 2020.
- tropmet. Tropical cyclone. Linter, <http://people.envsci.rutgers.edu/lintner/tropmet/Lecture17.pdf>, Accessed April 2020.
- ucar, n.d. mpas\_atmosphere\_users\_guide\_6.0. Www2.mmm.ucar.edu, n.d., [https://www2.mmm.ucar.edu/projects/mpas/mpas\\_atmosphere\\_users\\_guide\\_6.0.pdf](https://www2.mmm.ucar.edu/projects/mpas/mpas_atmosphere_users_guide_6.0.pdf), Accessed April 2020.