

Potential Theory in Multiply Connected Domains

Brian KYANJO Yao GAHOUNZO

*Department of Mathematics
Boise State University*

May 5, 2021

Outline

Introduction

Multiply Connected Domains

Interior and Exterior Problems in 2D

Proposed Solution

Introduction

Potential Theory

This is basically the study of harmonic function and its properties as a function according to mathematical physics (Garabedian and Schiffer, 1950).

Dirichlet problem

This is the problem of finding a function which solves a specified PDE in the interior of a given region that takes prescribed values on the boundary of the region.

Introduction

Potential Theory

This is basically the study of harmonic function and its properties as a function according to mathematical physics (Garabedian and Schiffer, 1950).

Dirichlet problem

This is the problem of finding a function which solves a specified PDE in the interior of a given region that takes prescribed values on the boundary of the region.

Multiple Connected Domains

Consider the Dirichlet problem in a finite open region D in the plane which is $(M + 1)$ -ply connected.




Figure: A bounded multiply connected domain D in the plane.
(Greenbaum et al., 1993)

The outer boundary of D is denoted by L_0 , while the interior boundary curves denoted by L_1, \dots, L_M .

Multiple Connected Domains

Consider the Dirichlet problem in a finite open region D in the plane which is $(M + 1)$ -ply connected.




Figure: A bounded multiply connected domain D in the plane.
(Greenbaum et al., 1993)

The outer boundary of D is denoted by L_0 , while the interior boundary curves denoted by L_1, \dots, L_M .

Cont.

In multiply connected domains, it is not so simple to reduce the cost of each iteration to $O(N \log N)$ or $O(N)$. Since the net cost of solving a linear system is $O(N^2)$.

Second Kind integral equation methods for the Dirichlet problem develop a nullspace of dimensions equal to the number of boundary components M .

The solution is obtained by:

- ▶ Projecting the Dirichlet data onto the range of operators (Subjecting data to certain capability constraints).
- ▶ Using a modified integral equation.

Cont.

In multiply connected domains, it is not so simple to reduce the cost of each iteration to $O(N \log N)$ or $O(N)$. Since the net cost of solving a linear system is $O(N^2)$.

Second Kind integral equation methods for the Dirichlet problem develop a nullspace of dimensions equal to the number of boundary components M .

The solution is obtained by:

- ▶ Projecting the Dirichlet data onto the range of operators (Subjecting data to certain capability constraints).
- ▶ Using a modified integral equation.

Cont.

In multiply connected domains, it is not so simple to reduce the cost of each iteration to $O(N \log N)$ or $O(N)$. Since the net cost of solving a linear system is $O(N^2)$.

Second Kind integral equation methods for the Dirichlet problem develop a nullspace of dimensions equal to the number of boundary components M .

The solution is obtained by:

- ▶ Projecting the Dirichlet data onto the range of operators (Subjecting data to certain capability constraints).
- ▶ Using a modified integral equation.

Interior and Exterior Problems in 2D

Consider a finite open region D in the plane with boundary L , which assumes to be smooth and to have continuous curvature.

The **Interior Dirichlet problem** involves the determination of a function $U(P)$ that satisfies;

$$\Delta U(P) = 0 \quad \text{for} \quad P \in D$$

with boundary condition

$$\lim_{\substack{P \rightarrow Q \\ P \in D}} U(P) = f(Q) \quad \text{for} \quad Q \in L$$

Interior and Exterior Problems in 2D

Consider a finite open region D in the plane with boundary L , which assumes to be smooth and to have continuous curvature.

The Interior Dirichlet problem involves the determination of a function $U(P)$ that satisfies;

$$\Delta U(P) = 0 \quad \text{for} \quad P \in D$$

with boundary condition

$$\lim_{\substack{P \rightarrow Q \\ P \in D}} U(P) = f(Q) \quad \text{for} \quad Q \in L$$

Cont.

Denote E to be the open region in the plane exterior to L .

The exterior Dirichlet problem is defined by:

$$\Delta U(P) = 0 \quad \text{for} \quad P \in E$$

with boundary condition

$$\lim_{\substack{P \rightarrow Q \\ P \in E}} U(P) = f(Q) \quad \text{for} \quad Q \in L$$

The double layer potential solution ($U(P)$):

$$U(P) = \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - P| dQ$$

where P is a point inside D , $\mu(Q)$ is the value value of the unknown dipole distribution at boundary point Q . $\frac{\partial}{\partial V_Q}$ is the outward normal derivative at the point Q .

Cont.

Denote E to be the open region in the plane exterior to L .

The exterior Dirichlet problem is defined by:

$$\Delta U(P) = 0 \quad \text{for} \quad P \in E$$

with boundary condition

$$\lim_{\substack{P \rightarrow Q \\ P \in E}} U(P) = f(Q) \quad \text{for} \quad Q \in L$$

The double layer potential solution ($U(P)$):

$$U(P) = \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - P| dQ$$

where P is a point inside D , $\mu(Q)$ is the value value of the unknown dipole distribution at boundary point Q . $\frac{\partial}{\partial V_Q}$ is the outward normal derivative at the point Q .

Cont.

Denote E to be the open region in the plane exterior to L .

The exterior Dirichlet problem is defined by:

$$\Delta U(P) = 0 \quad \text{for} \quad P \in E$$

with boundary condition

$$\lim_{\substack{P \rightarrow Q \\ P \in E}} U(P) = f(Q) \quad \text{for} \quad Q \in L$$

The double layer potential solution ($U(P)$):

$$U(P) = \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - P| dQ$$

where P is a point inside D , $\mu(Q)$ is the value value of the unknown dipole distribution at boundary point Q . $\frac{\partial}{\partial V_Q}$ is the outward normal derivative at the point Q .

Cont.

For any point Q_o on the boundary $U(P)$ satisfies the jump relation:

$$\lim_{\substack{P \rightarrow Q_o \\ P \in D}} U(P) = \frac{1}{2} \mu(Q_o) + \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - Q_o| dQ \quad (1)$$

$$\lim_{\substack{P \rightarrow Q_o \\ P \in E}} U(P) = -\frac{1}{2} \mu(Q_o) + \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - Q_o| dQ \quad (2)$$

Cont.

For any point Q_o on the boundary $U(P)$ satisfies the jump relation:

$$\lim_{\substack{P \rightarrow Q_o \\ P \in D}} U(P) = \frac{1}{2} \mu(Q_o) + \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - Q_o| dQ \quad (1)$$

$$\lim_{\substack{P \rightarrow Q_o \\ P \in E}} U(P) = -\frac{1}{2} \mu(Q_o) + \frac{1}{2\pi} \int_L \mu(Q) \frac{\partial}{\partial V_Q} \ln |Q - Q_o| dQ \quad (2)$$

Difference.

For interior problem, $\frac{\partial}{\partial v_Q}$ refers to the normal derivative in the direction outward from the domain D . Thus, whether the boundary point Q_0 lies on the outer boundary or one of the interior curves, the relevant jump condition is equation (1).

For exterior problem, $\frac{\partial}{\partial v_Q}$ refers to the inward normal derivative and the relevant jump condition is equation (2).

Difference.

For interior problem, $\frac{\partial}{\partial v_Q}$ refers to the normal derivative in the direction outward from the domain D . Thus, whether the boundary point Q_0 lies on the outer boundary or one of the interior curves, the relevant jump condition is equation (1).

For exterior problem, $\frac{\partial}{\partial v_Q}$ refers to the inward normal derivative and the relevant jump condition is equation (2).

Proposed Solution

The author proposed and presented a new integral equation method for the solution of the Dirichlet problem in multiply connected domains.

Fast multipole methods were combined with the new formulation to create algorithm capable of solving Laplace equations in domains of hundreds of distinct boundary components in minutes.

Proposed Solution

The author proposed and presented a new integral equation method for the solution of the Dirichlet problem in multiply connected domains.

Fast multipole methods were combined with the new formulation to create algorithm capable of solving Laplace equations in domains of hundreds of distinct boundary components in minutes.

Thank you!

briankyanjo@u.boisestate.edu
yaogahounzo@u.boisestate.edu

References

- Garabedian, P. and Schiffer, M. (1950). On existence theorems of potential theory and conformal mapping. *Annals of Mathematics*, pages 164–187.
- Greenbaum, A., Greengard, L., and McFadden, G. (1993). Laplace's equation and the dirichlet-neumann map in multiply connected domains. *Journal of Computational Physics*, 105(2):267–278.