

CS23331-DAA-2024-CSE / 5-Pair with Difference- $O(n^2)$ Time Complexity, $O(1)$  Space Complexity


## 5-Pair with Difference- $O(n^2)$ Time Complexity, $O(1)$ Space Complexity

Started on	Sunday, 12 October 2025, 7:11 PM
State	Finished
Completed on	Sunday, 12 October 2025, 7:12 PM
Time taken	52 secs
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

**Question 1** | Correct | Mark 1.00 out of 1.00 | [Flag question](#)

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[j] - A[i] = k$ ,  $i \neq j$ .

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as  $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3	1
1 3 5	
4	

**Answer:** (penalty regime: 0 %)

```
1 #include <stdio.h>
2
3 int main() {
4     int n;
5     scanf("%d", &n);
6
7     int arr[n];
8     for(int i = 0; i < n; i++) {
9         scanf("%d", &arr[i]);
10    }
11
12    int k;
13    scanf("%d", &k);
14
15    int i = 0, j = 1;
16    int found = 0;
17
18    while(j < n) {
19        if(i == j) j++;
20        if(j >= n) break;
21
22        int diff = arr[j] - arr[i];
23        if(diff == k) {
24            found = 1;
25            break;
26        } else if(diff < k) {
27            j++;
28        } else {
29            i++;
30        }
31    }
32
33    printf("%d\n", found);
34    return 0;
35 }
36
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

**Correct**

Marks for this submission: 1.00/1.00.

[Finish review](#)

[Back to Course](#)

[Data retention summary](#)