

CS 228 : Logic in Computer Science

S. Krishna

FOL : Notations

- ▶ Recall : A signature τ consists of a finite set of constants, relations and functions
- ▶ Recall : A formula φ is over some chosen signature τ
- ▶ Recall : A τ -structure (or a τ -model) consists of a universe (or domain), and gives meanings to the constants, functions and relations in τ .
- ▶ Recall notation : $\varphi(x_1, \dots, x_n)$
- ▶ Recall Assignments, satisfaction

FOL : Notations

- ▶ Recall : A signature τ consists of a finite set of constants, relations and functions
- ▶ Recall : A formula φ is over some chosen signature τ
- ▶ Recall : A τ -structure (or a τ -model) consists of a universe (or domain), and gives meanings to the constants, functions and relations in τ .
- ▶ Recall notation : $\varphi(x_1, \dots, x_n)$
- ▶ Recall Assignments, satisfaction
- ▶ Signature τ_E having a single binary relation E ,
- ▶ Signature τ_W having binary relations $S, <$ as well as unary relations Q_a for finitely many symbols a .

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$

Example of Satisfaction

$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
 $\alpha(y) = 1, \alpha(z) = 3$, and consider $\alpha[x \mapsto 2]$.

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
 $\alpha(y) = 1, \alpha(z) = 3$, and consider $\alpha[x \mapsto 2]$.
- ▶ Check this: $\mathcal{G} \not\models \exists x \forall y E(x, y)$,

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
 $\alpha(y) = 1, \alpha(z) = 3$, and consider $\alpha[x \mapsto 2]$.
- ▶ Check this: $\mathcal{G} \not\models \exists x \forall y E(x, y)$, $\mathcal{G} \models \forall x \exists y E(x, y)$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$
 - ▶ There is an assignment α for which
 $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$
 - ▶ There is an assignment α for which
 $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$
One possibility : $\alpha(x) = 2, \alpha(y) = 3$
 - ▶ There is no assignment α which satisfies
 $\exists x \exists y (Q_b(x) \wedge Q_b(y) \wedge x \neq y)$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$
 - ▶ There is an assignment α for which
 $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$
One possibility : $\alpha(x) = 2, \alpha(y) = 3$
 - ▶ There is no assignment α which satisfies
 $\exists x \exists y (Q_b(x) \wedge Q_b(y) \wedge x \neq y)$
 - ▶ Prove or disprove : $\mathcal{W} \models \exists x \forall y [Q_b(x) \wedge x < y \wedge Q_a(y)]$
 - ▶ Prove or disprove : $\mathcal{W} \models \exists x \forall y [Q_b(x) \wedge x < y \Rightarrow Q_a(y)]$

Satisfiability, Validity, Equivalence

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$

Satisfiability, Validity, Equivalence

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula φ over a signature τ is said to be **valid** iff for every τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$

Satisfiability, Validity, Equivalence

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula φ over a signature τ is said to be **valid** iff for every τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ Formulae $\varphi(x_1, \dots, x_n)$ and $\psi(x_1, \dots, x_n)$ are **equivalent** denoted $\varphi \equiv \psi$ iff for every \mathcal{A} and α , $\mathcal{A} \models_{\alpha} \varphi$ iff $\mathcal{A} \models_{\alpha} \psi$

Check SAT

- ▶ $\varphi = \exists x[(\forall y E(x, y)) \wedge \forall z[(\forall y E(z, y)) \rightarrow z = x]]$. Does φ evaluate to true under some graph structure?

Check SAT

- ▶ $\varphi = \exists x[(\forall y E(x, y)) \wedge \forall z[(\forall y E(z, y)) \rightarrow z = x]]$. Does φ evaluate to true under some graph structure?
- ▶ $\psi(z) = \exists x[Q_a(x) \wedge \forall y[(y < x \wedge Q_b(y)) \rightarrow (z < x \wedge y < z \wedge Q_a(z))]]$. Does ψ evaluate to true under some word structure?

Consider the formula

$$\varphi = \forall x. Q_a(x) \vee [\forall x. (Q_a(x) \Rightarrow \exists y. (Q_b(y) \wedge x < y))].$$

1. The word *aaa* is a model for φ
2. The word *b* is a model for φ
3. The word *ab* is a model for φ
4. The word *aba* is a model for φ
5. The word *bab* is a model for φ
6. The word *abab* is a model for φ
7. The word *baaaaa* is a model for φ
8. The word *bbb* is a model for φ , but *bb* is not
9. The word *abb* is not a model for φ , but *bba* is
10. Every word over *a, b* is a model for φ