

## QUESTION 1

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		

For the solution of this question please use only the front face and if necessary the back face of this page.

[12p] a) Find the length of the curve  $\vec{r}(t) = \frac{1}{2}(\sin t - t \cos t) \vec{i} + \frac{1}{2}(\cos t + t \sin t) \vec{j}$ ,  $1 \leq t \leq 3$   
and find its unit tangent vector for  $t = \frac{\pi}{3}$ .

[13p] b) Find the distance from the point  $P(1, 1, 2)$  to the plane passing through the points  $A(2, 2, -1)$ ,  $B(0, 3, 1)$  and  $C(-1, -1, -2)$ .

$$a) \quad L = \int_a^b |\vec{v}| dt \quad \vec{v} = \frac{1}{2} (\cos t - \cos t + t \sin t) \vec{i} + \frac{1}{2} (-\sin t + \sin t + t \cos t) \vec{j}$$

$$\vec{v} = \frac{t}{2} \sin t \vec{i} + \frac{t}{2} \cos t \vec{j}$$

$$|\vec{v}| = \sqrt{\left(\frac{t}{2} \sin t\right)^2 + \left(\frac{t}{2} \cos t\right)^2} = \frac{|t|}{2} = \frac{t}{2} \quad (1 \leq t \leq 3)$$

$$L = \int_1^3 \frac{t}{2} dt = \frac{1}{2} \frac{t^2}{2} \Big|_1^3 = 2 //$$

$$\vec{T} = \frac{1}{|\vec{v}|} \vec{v} = \frac{1}{t/2} \left( \frac{t}{2} \sin t \vec{i} + \frac{t}{2} \cos t \vec{j} \right) = \sin t \vec{i} + \cos t \vec{j}$$

$$t = \frac{\pi}{3} \Rightarrow \vec{T} = \frac{\sqrt{3}}{2} \vec{i} + \frac{1}{2} \vec{j}$$

$$b) \quad d = \frac{|\vec{PS} \cdot \vec{n}|}{|\vec{n}|}$$

$P$ : the point from which the distance is to be calculated

$S$ : a point on the plane

$\vec{n}$ : normal vector of the plane

$$\vec{BA} = (2-0)\vec{i} + (2-3)\vec{j} + (-1-1)\vec{k} = 2\vec{i} - \vec{j} - 2\vec{k}$$

$$\vec{BC} = (0-(-1))\vec{i} + (3-(-1))\vec{j} + (1-(-2))\vec{k} = \vec{i} + 4\vec{j} + 3\vec{k}$$

$$\vec{n} = \vec{BA} \times \vec{BC} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -1 & -2 \\ 1 & 4 & 3 \end{vmatrix} = 5\vec{i} - 8\vec{j} + 9\vec{k}, \quad |\vec{n}| = \sqrt{170}$$

$$\text{Let } S = A(2, 2, -1) : \vec{PS} = \vec{i} + \vec{j} - 3\vec{k} \Rightarrow d = \frac{|5-8-27|}{\sqrt{170}} = \frac{30}{\sqrt{170}} //$$

## QUESTION 2

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		

For the solution of this question please use only the front face and if necessary the back face of this page.

[12p] a) Investigate the continuity of the function

$$f(x, y) = \begin{cases} \frac{2x^2}{x^2 + 2xy + y} & (x, y) \neq (0, 0) \\ 1 & (x, y) = (0, 0) \end{cases}$$

at the point  $(x, y) = (0, 0)$ .[13p] b) Find the directional derivative of the function  $f(x, y, z) = x + x \cos z - y \sin z + y$  at the point  $(1, -2, \pi)$  in the direction of  $\vec{v} = 2\vec{i} - 2\vec{j} - 2\vec{k}$  and at this point determine the direction in which the function increases most rapidly.(a)  $f(0, 0) = 1$  exists

$$f(x, y) \Big|_{x=0, y \neq 0} = \frac{2 \cdot 0^2}{0^2 + 2 \cdot 0 \cdot y + y} = 0$$

$$\text{so } \lim_{\substack{(x, y) \rightarrow (0, 0) \\ \text{along } x=0, y \neq 0}} f(x, y) = \lim_{y \rightarrow 0} 0 = 0$$

$$f(x, y) \Big|_{y=0, x \neq 0} = \frac{2x^2}{x^2 + 2x \cdot 0 + 0} = 2$$

$$\text{so } \lim_{\substack{(x, y) \rightarrow (0, 0) \\ \text{along } y=0, x \neq 0}} f(x, y) = \lim_{x \rightarrow 0} 2 = 2$$

$\lim_{(x, y) \rightarrow (0, 0)} f(x, y)$   
does not exist  
by the two-path test.

$f$  is not cont.  
at  $(0, 0)$ .

$$(b) \quad \vec{v} = 2\vec{i} - 2\vec{j} - 2\vec{k}, \quad |\vec{v}| = 2\sqrt{3} \rightarrow \vec{u} = \frac{\vec{v}}{|\vec{v}|} = \frac{1}{\sqrt{3}}(\vec{i} - \vec{j} - \vec{k})$$

$$\vec{\nabla} f = f_x \vec{i} + f_y \vec{j} + f_z \vec{k} = (1 + \cos z) \vec{i} + (-\sin z + 1) \vec{j} - (x \sin z + y \cos z) \vec{k}$$

$$\vec{\nabla} f \Big|_{(1, -2, \pi)} = 0 \cdot \vec{i} + \vec{j} - 2\vec{k}$$

$$(D_{\vec{u}} f)_{P_0} = \vec{\nabla} f \Big|_{P_0} \cdot \vec{u} = (0 \cdot \vec{i} + \vec{j} - 2\vec{k}) \cdot \left( \frac{1}{\sqrt{3}}(\vec{i} - \vec{j} - \vec{k}) \right) = \frac{1}{\sqrt{3}}$$

$$\vec{\nabla} f \Big|_{(1, -2, \pi)} = \vec{j} - 2\vec{k} \quad \text{is the direction in which } f \text{ increases most rapidly.}$$

## QUESTION 3

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		

For the solution of this question please use only the front face and if necessary the back face of this page.

[10p] a) Find the equation of the tangent plane of the surface  $z = e^{x+y} + 2xy - 2$  at the point  $(1, -1, -3)$ .[15p] b) Using the method of Lagrange multipliers, find the maximum and minimum values of the function  $f(x, y) = x^2 + y^2$  subject to the constraint  $x^2 + x + y^2 + 2y = 0$ .

$$(a) \quad g(x, y, z) = e^{x+y} + 2xy - z - 2 = 0$$

$$\vec{\nabla} g = (e^{x+y} + 2y) \vec{i} + (e^{x+y} + 2x) \vec{j} - 1 \cdot \vec{k}$$

$$\vec{n} = \vec{\nabla} g \Big|_{(1, -1, -3)} = -\vec{i} + 3\vec{j} - \vec{k}$$

$$-1 \cdot (x-1) + 3 \cdot (y-(-1)) - 1 \cdot (z-(-3)) = 0 \Rightarrow -x + 3y - z = -1 //$$

$$(b) \quad g(x, y) = x^2 + x + y^2 + 2y = 0$$

$$\vec{\nabla} f = \lambda \vec{\nabla} g \Rightarrow 2x \vec{i} + 2y \vec{j} = \lambda [(2x+1) \vec{i} + (2y+2) \vec{j}]$$

$$2x = \lambda(2x+1) \quad \left. \begin{array}{l} 2x(1-\lambda) = \lambda \quad (1) \\ 2y(1-\lambda) = 2\lambda \quad (2) \\ x^2+x+y^2+2y=0 \quad (3) \end{array} \right\}$$

$$2y = \lambda(2y+2)$$

$$x^2+x+y^2+2y=0$$

$$(1) \&(2) : 2y(1-\lambda) = 2 \cdot 2x \cdot (1-\lambda) \rightarrow 2(y-2x)(1-\lambda) = 0$$

Two possibilities: If  $\lambda=1$ , (1) & (2) gives  $\lambda=0$ ; so not possible.

$$y=2x : (3) \text{ gives } x^2+x+4x^2+4x=0 \rightarrow 5x(x+1)=0$$

$$x=0, y=0 : f(0,0)=0 \text{ minimum value of } f$$

$$x=-1, y=-2 : f(-1,-2)=5 \text{ maximum value of } f$$

## QUESTION 4

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		


For the solution of this question please use only the front face and if necessary the back face of this page.

Lütfen bu soruyu bu kağıdın ön yüzünü ve gerekirse arka yüzünü kullanarak cevaplayınız.

[10p] a) Sketch the the region of integration of  $\int_0^1 \int_{x^2}^1 4xe^{y^2} dy dx$  and evaluate the integral.

[15p] b) Using cylindrical coordinates, find the volume of the solid bounded from below by the cone  $z = -\sqrt{x^2 + y^2}$ , above by the sphere  $x^2 + y^2 + z^2 = 1$  and laterally (from the sides) by the cylinder  $x^2 + y^2 = 1$ .

(a)




$$\int_0^1 \int_{x^2}^1 4xe^{y^2} dy dx = \int_{y=0}^1 \int_{x=0}^{x=\sqrt{y}} 4xe^{y^2} dx dy$$

$$= \int_{y=0}^1 \left( 2x^2 e^{y^2} \Big|_{x=0}^{x=\sqrt{y}} \right) dy = \int_0^1 2ye^{y^2} dy$$

$$= e^{y^2} \Big|_0^1 = e - 1$$

(b)



In Cylindrical coordinates:

$$z = -\sqrt{x^2 + y^2} \rightarrow z = -r$$

$$x^2 + y^2 + z^2 = 1, z \neq 0 \rightarrow z = -\sqrt{1-r^2}$$

$$x^2 + y^2 = 1 \rightarrow r = 1 \quad -\sqrt{x^2 + y^2} = -\sqrt{1-r^2} \rightarrow r = \frac{1}{\sqrt{2}}$$

$$\text{Volume} = \int_0^{2\pi} \int_{1/\sqrt{2}}^1 \int_{-r}^{-\sqrt{1-r^2}} dz r dr d\theta$$

$$= \int_0^{2\pi} \int_{1/\sqrt{2}}^1 (-\sqrt{1-r^2} + r) r dr d\theta$$

$$= \int_0^{2\pi} \left[ -\frac{1}{3}(1-r^2)^{3/2} + \frac{r^3}{3} \right]_{1/\sqrt{2}}^1 d\theta = \frac{2\pi}{3} \left( 1 - \frac{1}{\sqrt{2}} \right)$$