

# Notes on Real Analysis

Kai-Jyun Wang\*

Fall 2024

## Foreword

I took Real Analysis in the fall of 2024 with Professor Tien. This is my note on the course. I tried to include all the proofs and details that has or has not been covered in the class, in order to make this note as self-contained as possible. Some of the proofs might be taken from somewhere and some might be wrong. The following topics are covered in the lecture: measure theory, Lebesgue integration, Banach space, Hilbert space, and approximation theory.

Some funny things happened in the class. The professor had taught so fast that we had already reached the Banach space before our first midterm. Every student was wondering if the professor forgot that this is actually a one-year course. Time comes to the eleventh week, the professor walked into the class and said, “Few days ago, someone told me that we actually have two semesters for real analysis, and I didn’t know that before!” It turns out that our concern was right. The professor then said, “But that is also a good thing, because we can learn more advanced topics in the second semester, like the harmonic analysis, the Fourier analysis...”

---

\*National Taiwan University, Department of Economics.

# Contents

<b>1</b>	<b>Measure Theory and Integration</b>	<b>3</b>
1.1	Lebesgue Measure . . . . .	3
1.2	Lebesgue Integration . . . . .	7
1.3	Differentiation . . . . .	13
1.4	Radon-Nikodym Theorem . . . . .	16
1.5	Product Measure . . . . .	21
1.6	Convergence in Measure . . . . .	25
<b>2</b>	<b>Banach Space</b>	<b>31</b>
2.1	Banach Space and Bounded Linear Functional . . . . .	31
2.2	$\ell^p$ Space . . . . .	33
2.3	$\mathcal{L}^p$ Space . . . . .	35
2.4	Dual Space . . . . .	40
2.5	Weak Convergence . . . . .	54
2.6	Open Mapping Theorem and Closed Graph Theorem . . . . .	67
<b>3</b>	<b>Hilbert Space</b>	<b>71</b>
3.1	Cauchy-Schwarz Inequality . . . . .	71
3.2	Separability and Orthonormal Basis . . . . .	76
3.3	Riesz Representation and Bilinear Form . . . . .	80
3.4	Symmetric and Compact Operators . . . . .	85
<b>4</b>	<b>Approximation Theory and Fourier Theory</b>	<b>93</b>
4.1	Approximation by Polynomials . . . . .	93
4.2	Fourier Series . . . . .	101
4.3	Fourier Transform . . . . .	114

# 1. Measure Theory and Integration

## 1.1. Lebesgue Measure

### Definition 1.1

The **length** of an open interval  $(a, b) = I$  is  $b - a$  in the extended sense, denoted by  $\ell(I)$ .

### Remark

We define  $(a, a) = \emptyset$ .

### Definition 1.2

The **Lebesgue outer measure** (or in brief, **outer measure**) of a set  $E \subset \mathbb{R}$  is

$$\mu^*(E) = \inf \left\{ \sum_n \ell(I_n) \mid I_n \text{ are countable open intervals covering } E \right\}.$$

### Proposition 1.3

- (a) Countable sets are of outer measure zero.
- (b) If  $A \subset B$ , then  $\mu^*(A) \leq \mu^*(B)$ .
- (c) For  $x \in \mathbb{R}$  and  $A \subset \mathbb{R}$ ,  $\mu^*(A + x) = \mu^*(A)$ .
- (d) For countable  $A_n \subset \mathbb{R}$ ,  $\mu^*(\bigcup_n A_n) \leq \sum_n \mu^*(A_n)$ .

*Proof.* For (a), let  $x_n$  denumerate a countable set  $A$ . Then consider

$$I_n = (x_n - 2^{-n}\epsilon, x_n + 2^{-n}\epsilon)$$

for  $n \in \mathbb{N}$ . Then  $A \subset \bigcup_n I_n$  and  $\mu^*(A) \leq \sum_n 2 \cdot 2^{-n}\epsilon = 2\epsilon$ . Since  $\epsilon$  is arbitrary,  $\mu^*(A) = 0$ .

For (b), note that any cover of  $B$  must cover  $A$ . The result follows.

For (c), note that the translations of open intervals preserve their lengths.

For (d), let  $\{I_j^n\}$  cover  $A_n$  for each  $n$  such that  $\sum_j \ell(I_j^n) < \mu^*(A_n) + 2^{-n}\epsilon$ . Then we have that  $\bigcup_n \bigcup_j I_j^n$  covers  $\bigcup_n A_n$  and

$$\sum_n \sum_j \ell(I_j^n) < \sum_n \mu^*(A_n) + 2^{-n}\epsilon = \epsilon + \sum_n \mu^*(A_n).$$

Since  $\epsilon$  is arbitrary, it follows that  $\mu^*(\bigcup_n A_n) \leq \sum_n \mu^*(A_n)$ . ■

### Definition 1.4

A family of sets  $\mathcal{M}$  is called a  **$\sigma$ -algebra** if

- (a)  $\emptyset \in \mathcal{M}$ .
- (b)  $A \in \mathcal{M}$  implies  $A^c \in \mathcal{M}$ .
- (c) For countably many  $A_n \in \mathcal{M}$  we have  $\bigcup_n A_n \in \mathcal{M}$ .

The space  $(X, \mathcal{M})$  is called a **measurable space** and the sets in  $\mathcal{M}$  are called **measurable sets**.

**Proposition 1.5**

$\mathcal{M}$  is a  $\sigma$ -algebra if and only if the following hold:

- (a)  $X \in \mathcal{M}$ .
- (b)  $A, B \in \mathcal{M}$  implies  $A \cap B, A \cup B, A - B \in \mathcal{M}$ .
- (c) For countably many  $A_n \in \mathcal{M}$  we have  $\bigcap_n A_n \in \mathcal{M}$ .

*Proof.* Omitted. ■

**Proposition 1.6**

Let  $\mathcal{F}$  be a family of sets in  $X$ . Then there exists a unique smallest  $\sigma$ -algebra containing  $\mathcal{F}$ .

*Proof.* Let  $\mathcal{M}$  be the intersection of all  $\sigma$ -algebras containing  $\mathcal{F}$ . Since  $\mathcal{P}(X)$  must be such a  $\sigma$ -algebra,  $\mathcal{M}$  is non-empty. Now we verify that  $\mathcal{M}$  is a  $\sigma$ -algebra. First,  $\emptyset \in \mathcal{M}$  since  $\emptyset$  is in every  $\sigma$ -algebra. Second, if  $A \in \mathcal{F}$  then  $A$  must belong to every  $\sigma$ -algebra containing  $\mathcal{F}$  and so does  $A^c$ . Hence  $A^c \in \mathcal{M}$ . The closure under countable unions follows from a similar argument. We conclude that  $\mathcal{M}$  is the desired  $\sigma$ -algebra. ■

**Definition 1.7**

For a family of sets  $\mathcal{F}$ , we denote the smallest  $\sigma$ -algebra containing  $\mathcal{F}$  by  $\sigma(\mathcal{F})$ .

**Definition 1.8**

Let  $\mathcal{T}$  be the family of all open sets. The **Borel  $\sigma$ -algebra** is defined as  $\mathcal{B} = \sigma(\mathcal{T})$ . The sets in  $\mathcal{B}$  are called **Borel sets**.

**Definition 1.9**

A set  $E$  is called **Lebesgue measurable** if for  $\epsilon > 0$ , there exists an open set  $V$  such that  $E \subset V$  and  $\mu^*(V - E) \leq \epsilon$ .

**Remark**

The Lebesgue measurable sets form a  $\sigma$ -algebra.

**Remark**

The Borel sets are Lebesgue measurable.

**Remark**

Not all subsets in  $\mathbb{R}$  are Lebesgue measurable. Consider the Vitali set. For a Lebesgue measurable set that is not Borel, consider the preimage of a Vitali set of Cantor-Lebesgue function.

**Definition 1.10**

A function  $f : (X, \mathcal{M}) \rightarrow (\mathbb{R}, \mathcal{B})$  is called  **$\mathcal{M}$ -measurable** if  $f^{-1}(B) \in \mathcal{M}$  for all  $B \in \mathcal{B}$ .

**Proposition 1.11**

Let  $f : X \rightarrow Y$  and  $A$  be an index set. Then

- (a)  $f^{-1}(B^c) = f^{-1}(B)^c$ .
- (b)  $f^{-1}(\bigcup_{a \in A} B_a) = \bigcup_{a \in A} f^{-1}(B_a)$ .
- (c)  $f^{-1}(\bigcap_{a \in A} B_a) = \bigcap_{a \in A} f^{-1}(B_a)$

*Proof.* Omitted. ■

**Proposition 1.12**

$f : (X, \mathcal{M}) \rightarrow (\mathbb{R}, \mathcal{B})$  is  $\mathcal{M}$ -measurable if  $f^{-1}((a, \infty)) \in \mathcal{M}$ .

*Proof.* Observe that  $\{A \subset \mathbb{R} \mid f^{-1}(A) \in \mathcal{F}\}$  is a  $\sigma$ -algebra. By assumption,  $[a, b]$ ,  $(a, b]$ ,  $[a, b)$  and  $(a, b)$  are in this  $\sigma$ -algebra for  $a, b \in \overline{\mathbb{R}}$ . ■

**Proposition 1.13**

$f_n$  are measurable. Then  $\sup_n f_n$ ,  $\inf_n f_n$ ,  $\limsup f_n$  and  $\liminf f_n$  are measurable.

*Proof.* Note that  $\{\sup_n f_n > a\} = \bigcup_n \{f_n > a\}$  and  $\{\inf_n f_n < a\} = \bigcup_n \{f_n < a\}$  are measurable.  $\limsup_n f_n = \inf_k \sup_{n \geq k} f_n$  and  $\liminf_n f_n = \sup_k \inf_{n \geq k} f_n$  are measurable as well. ■

**Remark**

$\lim_n f_n = \limsup_n f_n = \liminf_n f_n$  is measurable.

**Definition 1.14**

Let  $(X, \mathcal{M})$  be a measurable space. A **measure** on  $X$  is a function  $\mu : \mathcal{M} \rightarrow [0, \infty]$  satisfying

- (a)  $\mu(\emptyset) = 0$ .
- (b)  $\mu(\bigcup_n A_n) = \sum_n \mu(A_n)$  for disjoint  $A_n$ .

The triple  $(X, \mathcal{M}, \mu)$  is called a **measure space**.

**Proposition 1.15**

Let  $(X, \mathcal{M}, \mu)$  be a measure space and  $A, B \in \mathcal{M}$ . Then

- (a)  $A \subset B$  implies  $\mu(A) \leq \mu(B)$ .
- (b)  $\mu(A - B) = \mu(A) - \mu(B)$  if  $B \subset A$  and  $\mu(B) < \infty$ .

*Proof.* Omitted. ■

**Proposition 1.16**

Let  $(X, \mathcal{M}, \mu)$  be a measure space and  $E_n$  be a sequence of measurable sets. Then

- (a) If  $E_n \nearrow E$ , then  $\mu(E_n) \rightarrow \mu(E)$  as  $n \rightarrow \infty$ .
- (b) If  $E_n \searrow E$  and  $\mu(E_1) < \infty$ , then  $\mu(E_n) \rightarrow \mu(E)$  as  $n \rightarrow \infty$ .

*Proof.* Suppose  $\mu(E_n) < \infty$  for all  $n$ . Consider  $S_n = E_n - E_{n-1}$  with  $E_0 = \emptyset$ . Then  $S_n$  are disjoint and  $\bigcup_n S_n = E$ . Then

$$\mu(E) = \mu\left(\bigcup_n S_n\right) = \sum_n \mu(S_n) = \sum_n \mu(E_n) - \mu(E_{n-1}) = \lim_n \mu(E_n).$$

If  $\mu(E_n) = \infty$  for some  $n$ , then  $\mu(E) = \infty$  and the result follows.

For the second part, note that  $E_1 - E_n \nearrow E_1 - E$ . Then

$$\mu(E_1) - \mu(E_n) = \mu(E_1 - E_n) \rightarrow \mu(E_1 - E) = \mu(E_1) - \mu(E).$$

Rearranging gives the desired result. ■

**Theorem 1.17** (Egorov)

Let  $E$  be a measurable set with  $\mu(E) < \infty$  and  $f_n : E \rightarrow \mathbb{R}$  are measurable functions. If  $f_n \rightarrow f$  a.e. on  $E$ , then for all  $\epsilon > 0$ , there exists a closed set  $A_\epsilon \subset E$  such that  $\mu(E - A_\epsilon) < \epsilon$  and  $f_n \rightarrow f$  uniformly on  $A_\epsilon$ .

*Proof.* Consider the case where  $f_n \rightarrow f$  everywhere on  $E$  since  $\{x \in E \mid f_n(x) \not\rightarrow f(x)\}$  is of measure zero. For each  $n, k \in \mathbb{N}$ , let  $E_k^n = \{x \in E \mid |f_j(x) - f(x)| < 1/n \text{ for all } j > k\}$ . Then fix  $n$  and note that  $E_k^n \nearrow E$  as  $k \rightarrow \infty$ . By [proposition 1.16](#), there exists  $k_n$  such that  $\mu(E - E_{k_n}^n) < 2^{-n}$ . Then we have  $|f_j(x) - f(x)| < 1/n$  for every  $j > k_n$  and  $x \in E_{k_n}^n$ . Choose  $N$  such that  $\sum_{n \geq N} 2^{-n} < \epsilon/2$  and let  $\hat{A}_\epsilon = \bigcap_{n \geq N} E_{k_n}^n$ . Then  $\mu(E - \hat{A}_\epsilon) \leq \sum_{n \geq N} \mu(E - E_{k_n}^n) < \epsilon/2$ . Also, for any  $\delta > 0$ , we may pick  $n > N$  with  $1/n < \delta$  and for  $x \in \hat{A}_\epsilon$ ,  $|f_j(x) - f(x)| < \delta$  whenever  $j > k_n$ . Hence  $f_n \rightarrow f$  uniformly on  $\hat{A}_\epsilon$ . We may further find a closed  $A_\epsilon \subset \hat{A}_\epsilon$  such that  $\mu(\hat{A}_\epsilon - A_\epsilon) < \epsilon/2$ . Then  $A_\epsilon$  is the desired set. ■

**Definition 1.18**

A sequence of measurable functions  $f_n$  is said to **converge almost uniformly** to a function  $f$  if for every  $\epsilon > 0$ , there exists a measurable set  $E_\epsilon$  such that  $\mu(E_\epsilon^c) < \epsilon$  and  $f_n \rightarrow f$  uniformly on  $E_\epsilon$ .

**Remark**

The Egorov theorem states that if the space is of finite measure, then converging almost everywhere implies converging almost uniformly.

**Definition 1.19**

A function  $s : X \rightarrow Y$  is called **simple** if it only takes finitely many values.

**Lemma 1.20**

$f : E \rightarrow [0, \infty]$  is measurable. Then there exists a sequence of simple functions  $s_n \nearrow f$ ; furthermore, if  $f$  is bounded, then  $s_n \rightarrow f$  uniformly.

*Proof.* Consider  $s_n = \sum_{k=0}^{2^n-1} k 2^{-n} \chi_{f^{-1}([k 2^{-n}, (k+1) 2^{-n}))} + n \chi_{f^{-1}([n, \infty))}$ . Then  $s_n$  are simple and  $s_n \nearrow f$ . If  $f$  is bounded, then  $f^{-1}([n, \infty)) = \emptyset$  for some  $n$  large enough and  $s_n \rightarrow f$  uniformly. ■

**Theorem 1.21** (Lusin)

Let  $E \subset \mathbb{R}$  be a set of finite measure and  $f : E \rightarrow \mathbb{R}$  be a measurable, finite-valued function. Then for all  $\epsilon > 0$ , there exists a closed set  $F_\epsilon \subset E$  such that  $\mu(E - F_\epsilon) < \epsilon$  and  $f|_{F_\epsilon}$  is continuous.

*Proof.* First we may partition  $E$  into  $E = \bigcup_{i \in \mathbb{N}} E_i$  where  $E_i = E \cap [-i, i]$ . We first prove the result for simple functions. Let  $f = \sum_{j=1}^N c_j \chi_{A_j}$  be a simple function with the stated properties. Then for each  $j$ , we may find a closed set  $F_j \subset A_j$  such that  $\mu(A_j - F_j) < \epsilon/N$ . Now since  $E_i$  are bounded,  $F_j \cap E_i$  are compact and hence  $f$  being constant on each  $F_j \cap E_i$  is continuous. Note that  $F_\epsilon = \bigcup_{i,j=1}^N F_j \cap E_i$  satisfies the desired properties. Next, for a general measurable function  $f$ , we may find a sequence of simple functions  $s_n \nearrow f$  by [lemma 1.20](#). Now by Egorov's theorem, we may find a closed set  $F_\epsilon \subset E$  such that  $\mu(E - F_\epsilon) < \epsilon$  and  $s_n \rightarrow f$  uniformly on  $F_\epsilon$ . Since  $s_n$  are continuous on  $F_\epsilon$ ,  $f$  is continuous on  $F_\epsilon$ . ■

### Remark

By Tietze's extension theorem,  $f$  can be extended to a continuous function on all of  $\mathbb{R}$ .

### Proposition 1.22

$E$  is Lebesgue measurable if and only if  $\mu(E \Delta B) = 0$  for some Borel set  $B$ .

*Proof.* Suppose  $E$  is Lebesgue measurable. Then for each  $n$ , there exists an open set  $V_n$  such that  $E \subset V_n$  and  $\mu(V_n - E) < 1/n$ . Let  $B = \bigcap_n V_n$ . Then  $B$  is a Borel set and  $\mu(E \Delta B) = 0$ . Conversely, if  $\mu(E \Delta B) = 0$  for some Borel set  $B$ , since  $B$  is measurable, there exists an open  $V \supset B$  such that  $\mu(V - B) < \epsilon$ . Then  $B = (E \cap B) \cup (B - E)$  and since the later set has outer measure zero,  $E \cap B$  is measurable. And since  $E - B$  is outer measure zero,  $E \cap B = E$  is measurable. ■

### Proposition 1.23

If  $f$  is Lebesgue measurable, then there exists a Borel measurable function  $g$  such that  $f = g$  a.e.

*Proof.* Let  $s_k \nearrow f$  be a sequence of simple functions with  $s_k = \sum_{i=1}^{n_k} c_i \chi_{E_i}$  where  $E_i$  are measurable. Then for each  $E_i$  we may find a Borel set  $B_i \subset E_i$  such that  $\mu(E_i - B_i) = 0$  by the previous proposition. Then  $t_k = \sum_{i=1}^{n_k} c_i \chi_{B_i}$  is a Borel measurable function. Let  $g = \lim_{k \rightarrow \infty} t_k$ . Then  $g$  is Borel measurable and  $f = g$  a.e. since  $\mu(E_i - B_i) = 0$  for countably many  $i$ . ■

## 1.2. Lebesgue Integration

### Definition 1.24

For a simple function  $s = \sum_{i=1}^n c_i \chi_{E_i}$ , its **Lebesgue integral** is defined as

$$\int s d\mu = \sum_{i=1}^n c_i \mu(E_i).$$

### Definition 1.25

For a non-negative measurable function  $f$ , its **Lebesgue integral** is defined as

$$\int f d\mu = \sup \left\{ \int s d\mu \mid s \text{ is simple and } 0 \leq s \leq f \right\}.$$

**Definition 1.26**

For a measurable function  $f : X \rightarrow [-\infty, \infty]$ , its **Lebesgue integral** is defined as

$$\int f d\mu = \int f^+ d\mu - \int f^- d\mu,$$

where  $f^+ = \max\{f, 0\}$  and  $f^- = \max\{-f, 0\}$  provided that

$$\int |f| d\mu = \int f^+ d\mu + \int f^- d\mu < \infty.$$

In such a case, we say that  $f$  is **integrable**.

**Proposition 1.27**

For  $f, g$  integrable and  $c \in \mathbb{R}$ ,

- (a)  $\int cf + gd\mu = c \int fd\mu + \int gd\mu.$
- (b) If  $f \leq g$  a.e., then  $\int fd\mu \leq \int gd\mu.$

*Proof.* Omitted. ■

**Theorem 1.28** (Lebesgue Monotone Convergence Theorem)

Let  $f_n : X \rightarrow [0, \infty]$  be a sequence of measurable functions with  $f_n \nearrow f$  a.e. Then

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$$

*Proof.* By the monotonicity we have

$$\int f_n d\mu \leq \int f d\mu$$

for all  $n$  and hence

$$\lim_{n \rightarrow \infty} \int f_n d\mu \leq \int f d\mu.$$

To obtain the reverse inequality, note that for any  $c \in (0, 1)$ , there exists  $N$  such that  $f_n \geq cf$  a.e. for all  $n \geq N$ . Then

$$\int f_n d\mu \geq c \int f d\mu$$

for all  $n \geq N$ . Letting  $n \rightarrow \infty$ ,

$$\lim_{n \rightarrow \infty} \int f_n d\mu \geq c \int f d\mu.$$

Taking  $c \rightarrow 1^-$  then

$$\lim_{n \rightarrow \infty} \int f_n d\mu \geq \int f d\mu \implies \lim_{n \rightarrow \infty} \int f_n d\mu = \int f d\mu.$$
■



**Remark**

As a consequence,

$$\int \sum_n f_n d\mu = \sum_n \int f_n d\mu.$$

**Theorem 1.29** (Bounded Convergence Theorem)

Suppose  $\mu(X) < \infty$ . Let  $f_n : X \rightarrow \mathbb{R}_+$  be a sequence measurable functions such that  $f_n \leq M$  a.e. for some  $M \in \mathbb{R}$ . If  $f_n \rightarrow f$  a.e., then  $f$  is integrable and

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$$

*Proof.* For any  $\epsilon > 0$ , by Egorov's theorem, there exists  $F \subset X$  such that  $\mu(X - F) < \epsilon$  and  $f_n \rightarrow f$  uniformly on  $F$ . Then there exists  $N$  such that  $|f_n - f| < \epsilon$  on  $F$  for all  $n \geq N$ . We have

$$\begin{aligned} \left| \int f_n d\mu - \int f d\mu \right| &\leq \int_X |f_n - f| d\mu \\ &= \int_F |f_n - f| d\mu + \int_{X-F} |f_n - f| d\mu \\ &\leq \epsilon \mu(F) + 2M \mu(X - F) = \epsilon(\mu(F) + 2M\epsilon). \end{aligned}$$

Since  $\mu(X) < \infty$  and  $\epsilon$  is arbitrary, we may conclude that

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$$

■

**Lemma 1.30** (Fatou)

$f_n : X \rightarrow [0, \infty]$  are measurable. Then

$$\int \liminf_n f_n d\mu \leq \liminf_n \int f_n d\mu.$$

*Proof.* Let  $g_n = \inf_{k \geq n} f_k$ . Then  $g_n \nearrow g = \liminf_n f_n$ . By LMCT,

$$\int g_n d\mu \rightarrow \int g d\mu = \int \liminf_n f_n d\mu.$$

Note that  $f_n \geq g_n$  and thus  $\int f_n d\mu \geq \int g_n d\mu$ . Hence

$$\liminf_n \int f_n d\mu \geq \liminf_n \int g_n d\mu = \int g d\mu = \int \liminf_n f_n d\mu.$$

■

**Theorem 1.31** (Lebesgue Dominated Convergence Theorem)

Let  $f_n : X \rightarrow [-\infty, \infty]$  be a sequence of measurable functions such that  $f_n \rightarrow f$  a.e. and there

exists an integrable function  $g$  such that  $|f_n| \leq g$  a.e. for all  $n$ . Then

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$$

*Proof.* Since  $|f_n| \leq g$  a.e.,  $|f| \leq g$  a.e. Now  $|f_n - f| \leq |f_n| + |f| \leq 2g$  a.e. Let  $h_n = 2g - |f_n - f| \geq 0$  a.e. By Fatou's lemma,

$$\begin{aligned} \int 2g d\mu &= \int \liminf_n h_n d\mu \leq \liminf_n \int h_n d\mu = \liminf_n \int 2g - |f_n - f| d\mu \\ &= \int 2g d\mu - \limsup_n \int |f_n - f| d\mu. \end{aligned}$$

It follows that

$$0 \leq \liminf_n \int |f_n - f| d\mu \leq \limsup_n \int |f_n - f| d\mu \leq 0.$$

Hence

$$\lim_{n \rightarrow \infty} \int |f_n - f| d\mu = 0.$$

By the triangle inequality,

$$\left| \int f d\mu - \int f_n d\mu \right| \leq \int |f - f_n| d\mu \rightarrow 0.$$

So

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$$

■

### Remark

If  $\text{supp}(f)$  has finite measure and  $f$  is bounded, then

$$\int f = \inf_{s \geq f} \int s d\mu,$$

where  $s$  is simple.

### Definition 1.32

$\mathcal{L}^1 = \{f : X \rightarrow \mathbb{R} \mid f \text{ is integrable}\}$  with the norm  $\|f\|_{\mathcal{L}^1} = \int |f| d\mu$  is called the  $\mathcal{L}^1$  **space**.

### Remark

The elements in  $\mathcal{L}^1$  are in fact equivalence classes of functions that are equal a.e.

### Proposition 1.33

Let  $f \in \mathcal{L}^1$  be a nonnegative function. Then for every  $\epsilon > 0$ , there is some  $\delta > 0$  such that for any measurable  $E$  with  $\mu(E) \leq \delta$ ,

$$\int_E f d\mu \leq \epsilon.$$

*Proof.* Let  $E_n = \{x \in X \mid f(x) > n\}$ . Then by Lebesgue dominated convergence theorem, since  $f\chi_{E_n} \leq f$ ,

$$\int_{E_n} f d\mu \rightarrow 0.$$

For any  $\epsilon > 0$ , there exists  $n$  such that

$$\int_{E_n} f d\mu \leq \frac{\epsilon}{2}.$$

Pick  $\delta \leq \epsilon/(2n)$ . Then for any measurable  $E$  with  $\mu(E) \leq \delta$ ,

$$\int_E f d\mu = \int_{E \cap E_n} f d\mu + \int_{E \cap E_n^c} f d\mu \leq \int_{E_n} f d\mu + n\mu(E) \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

since  $f \leq n$  on  $E_n^c$ . This completes the proof. ■

**Theorem 1.34** (Lebesgue-Vitali)

$f : X \rightarrow \mathbb{R}$  is Riemann integrable if and only if the discontinuity set of  $f$  has Lebesgue measure zero. Furthermore, if  $f$  is Riemann integrable, then the Riemann integral and the Lebesgue integral agrees.

*Proof.* Define the oscillation of  $f$  at  $x$  as

$$\text{osc}(f, x) = \inf_{U: x \in U} \text{diam}(f(U)),$$

where  $U$  is open.

We first claim that  $f$  is continuous at  $x$  if and only if  $\text{osc}(f, x) = 0$ . Indeed, if  $f$  is continuous at  $x$ , then  $\forall \epsilon > 0, \exists \delta > 0$  such that  $|f(x) - f(y)| < \epsilon$  for all  $y \in B_\delta(x)$ . Then  $\text{diam}(f(B_\delta(x))) \leq 2\epsilon$ . Since  $\epsilon$  is arbitrary,  $\text{osc}(f, x) = 0$ . Conversely, if  $\text{osc}(f, x) = 0$ , then  $\forall \epsilon > 0, \exists$  open  $U$  containing  $x$  such that  $\text{diam}(f(U)) < \epsilon$ . This implies that  $|f(x) - f(y)| < \epsilon$  for all  $y \in U$  and hence  $f$  is continuous at  $x$ .

Next, let  $D_\epsilon$  collect all points  $x$  such that  $\text{osc}(f, x) \geq \epsilon > 0$ . We claim that  $D_\epsilon$  is closed. For any convergent sequence  $x_k \in D_\epsilon$ , let  $x_k \rightarrow x$ . For any open  $U$  containing  $x$ ,  $\exists N$  such that  $x_k \in U$  for all  $k \geq N$ . Then  $\exists$  an open neighborhood of  $x_N$ ,  $U'$ , such that  $U' \subset U$  and  $\text{diam}(f(U')) \geq \epsilon$ . Hence  $\text{osc}(f, x) \geq \epsilon$  and  $x \in D_\epsilon$ , showing that  $D_\epsilon$  is closed. Observe that  $D = \bigcup_{n=1}^{\infty} D_{1/n}$ .

Now suppose that  $f$  is Riemann integrable. Then for any  $\epsilon > 0, \exists \mathcal{P}$  such that  $U(f, \mathcal{P}) - L(f, \mathcal{P}) < \frac{1}{n}$  and  $\|\mathcal{P}\| < \frac{1}{n}$ . Then

$$\begin{aligned} & \sum_{\substack{Q \in \mathcal{P}, \\ Q \cap D_{\frac{1}{n}} \neq \emptyset}} (\sup_Q f - \inf_Q f) |Q| + \sum_{\substack{Q \in \mathcal{P}, \\ Q \cap D_{\frac{1}{n}} = \emptyset}} (\sup_Q f - \inf_Q f) |Q| \\ &= \sum_{Q \in \mathcal{P}} (\sup_Q f - \inf_Q f) |Q| = U(f, \mathcal{P}) - L(f, \mathcal{P}) < \epsilon. \end{aligned}$$

Note that  $\sup_Q f - \inf_Q f = \text{diam}(f(Q))$ . This gives that  $2M\mu^*(D_{\frac{1}{n}}) < \epsilon$  for every  $n$ . Since  $\epsilon$  is arbitrary, we conclude that  $\mu^*(D_{\frac{1}{n}}) = 0$  for each  $n$ . Thus  $D$  is an union of sets of measure zero and hence also has measure zero.

For the converse, suppose that  $m(D) = 0$ . Then  $D_\epsilon$  also has measure zero. Let  $\mathcal{P}$  be a partition on  $E$  with  $\|\mathcal{P}\| < \delta$  for some  $\delta > 0$ , which will be determined later. Then

$$\begin{aligned} U(f, \mathcal{P}) - L(f, \mathcal{P}) &= \sum_{Q \in \mathcal{P}} (\sup_Q f - \inf_Q f) |Q| \\ &= \sum_{\substack{Q \in \mathcal{P}, \\ Q \cap D_\epsilon = \emptyset}} (\sup_Q f - \inf_Q f) |Q| + \sum_{\substack{Q \in \mathcal{P}, \\ Q \cap D_\epsilon \neq \emptyset}} (\sup_Q f - \inf_Q f) |Q| \end{aligned}$$

For the first term,  $\sup_Q f - \inf_Q f < \epsilon$  for  $\|\mathcal{P}\| < \delta_1$  for some  $\delta_1 > 0$ . And thus the first term is bounded by  $\epsilon m(E)$ . For the second term,  $\sup_Q f - \inf_Q f < 2M$  and since  $D_\epsilon$  has measure zero,  $\exists Q_k$  cubic cover of  $D_\epsilon$  such that  $\sum_k |Q_k| < \epsilon$ . Now if  $\text{diam}(Q) < \delta_2$  for some  $\delta_2 > 0$ , then those  $Q$  intersecting  $D_\epsilon$  nonempty are subset of  $\bigcup_k Q_k$ . Thus the second term is bounded by  $2M\epsilon$ . Choosing  $\delta = \min\{\delta_1, \delta_2\}$  yields that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \epsilon m(E) + 2M\epsilon$$

whenever  $\|\mathcal{P}\| < \delta$ . Since  $\epsilon$  is arbitrary,  $f$  is Riemann integrable. ■

### Proposition 1.35

- (a) Step functions are dense in  $\mathcal{L}^1$ .
- (b) Continuous functions with compact support are dense in  $\mathcal{L}^1$ .

*Proof.* Let  $f \in \mathcal{L}^1$ . By [lemma 1.20](#), we already know that simple functions are dense in  $\mathcal{L}^1$ . It now remains to show that step functions can approximate simple functions. Since simple functions are linear combinations of finitely many characteristic functions, it suffices to show that characteristic functions can be approximated by step functions. Now for any measurable  $E$ , there is a family of almost disjoint cubes  $Q_i$  such that  $\mu(E \triangle \bigcup_{i=1}^M Q_i) \leq 2\epsilon$ , and thus we may set the step function to be  $\phi = \sum_{i=1}^M \chi_{Q_i}$ , with  $\|\chi_E - \phi\|_{\mathcal{L}^1} \leq 2\epsilon$ .

For the second part, let it now suffices to show that continuous functions with compact support can approximate characteristic functions of a rectangle, say  $[a, b]$ . Then set

$$g(x) = \begin{cases} 0 & x \leq a - \epsilon, \\ \frac{x-a+\epsilon}{\epsilon} & a - \epsilon \leq x \leq a, \\ 1 & a \leq x \leq b, \\ 1 - \frac{x-b}{\epsilon} & b \leq x \leq b + \epsilon, \\ 0 & x \geq b + \epsilon. \end{cases}$$

Then  $g$  is continuous with compact support and  $\|\chi_{[a,b]} - g\|_{\mathcal{L}^1} \leq \epsilon/2 + \epsilon/2 = \epsilon$ . ■

### 1.3. Differentiation

#### Definition 1.36

Let  $f \in \mathcal{L}^1(\mathbb{R}^d)$ . The **Hardy-Littlewood maximal function** is defined as

$$f^*(x) = \sup_{B: x \in B} \frac{1}{\mu(B)} \int_B |f(y)| dy,$$

where the supremum is taken over all balls containing  $x$ .

#### Proposition 1.37

$f^*$  is measurable.

*Proof.* Let  $E_\alpha = \{x \mid f^*(x) > \alpha\}$ . We claim that it is an open set. Indeed, if  $p \in E_\alpha$ , there exists a ball  $B$  containing  $p$  such that

$$\frac{1}{\mu(B)} \int_B |f(y)| dy > \alpha.$$

Now any  $x$  close enough to  $p$  will be contained in  $B$  and hence in  $E_\alpha$ . Thus  $E_\alpha$  is open. Hence  $f^*$  is measurable. ■

#### Lemma 1.38

[Vitali Covering Lemma] Suppose  $\{B_1, \dots, B_N\}$  is a finite collection of open balls in  $\mathbb{R}^d$ . Then there exists a disjoint subcollection  $\{B_{i_1}, \dots, B_{i_k}\}$  such that

$$\mu\left(\bigcup_{j=1}^N B_j\right) \leq 3^d \sum_{j=1}^k \mu(B_{i_j}).$$

*Proof.* First we make an observation that if  $B$  and  $B'$  are balls intersecting with, say, the radius of  $B$  is greater than the radius of  $B'$ , then  $B'$  is contained in the ball  $\tilde{B}$  that is concentric with  $B$  but with 3 times the radius.

The construction of the subcollection is proceeded as follows. First, pick a ball  $B_{i_1}$  with the largest radius. Then remove all balls intersecting with  $\tilde{B}_{i_1}$ , the ball concentric with  $B_{i_1}$  but with 3 times the radius. Among the remaining balls, we repeat the process and pick  $B_{i_2}$ . The process terminates when no more balls can be picked, after at most  $N$  steps and we obtain a disjoint subcollection of balls  $\{B_{i_1}, \dots, B_{i_k}\}$ .

Lastly, we verify the inequality. By the construction, we know that  $\bigcup_{j=1}^N B_j \subset \bigcup_{j=1}^k \tilde{B}_{i_j}$  and thus

$$\mu\left(\bigcup_{j=1}^N B_j\right) \leq \mu\left(\bigcup_{j=1}^k \tilde{B}_{i_j}\right) \leq \sum_{j=1}^k \mu(\tilde{B}_{i_j}) = \sum_{j=1}^k 3^d \mu(B_{i_j}).$$

■

#### Theorem 1.39 (Weak-Type Inequality)

Let  $f \in \mathcal{L}^1(\mathbb{R}^d)$ . Then for all  $\alpha > 0$ ,

$$\mu\left(\{x \in \mathbb{R}^d \mid f^*(x) > \alpha\}\right) \leq \frac{A}{\alpha} \|f\|_{\mathcal{L}^1(\mathbb{R}^d)},$$

where  $A = 3^d$ .

*Proof.* Let  $E_\alpha = \{x \mid f^*(x) > \alpha\}$ . For each  $x \in E_\alpha$  there exists a ball  $B_x$  containing  $x$  such that

$$\frac{1}{\mu(B_x)} \int_{B_x} |f(y)| dy > \alpha \quad \Rightarrow \quad \mu(B_x) < \frac{1}{\alpha} \int_{B_x} |f(y)| dy.$$

Now for any fixed compact  $K \subset E_\alpha$ ,  $K$  is covered by  $\bigcup_{x \in E_\alpha} B_x$ , and hence there exists a finite subcover  $\{B_1, \dots, B_N\}$  of  $K$ . By the Vitali covering lemma, there exists a disjoint subcollection  $\{B_{i_1}, \dots, B_{i_k}\}$  with

$$\mu\left(\bigcup_{j=1}^N B_j\right) \leq 3^d \sum_{j=1}^k \mu(B_{i_j}).$$

As a result,

$$\begin{aligned} \mu(K) &\leq \mu\left(\bigcup_{j=1}^N B_j\right) \leq 3^d \sum_{j=1}^k \mu(B_{i_j}) \leq \frac{3^d}{\alpha} \sum_{j=1}^k \int_{B_{i_j}} |f(y)| dy \\ &\leq \frac{3^d}{\alpha} \int_{\bigcup_{j=1}^k B_{i_j}} |f(y)| dy \leq \frac{3^d}{\alpha} \int_{\mathbb{R}^d} |f(y)| dy. \end{aligned}$$

Since the inequality holds for all compact subset  $K$  of  $E_\alpha$ , the proof is complete. ■

### Remark

Note that  $\{x \mid f^*(x) = \infty\} \subset \{x \mid f^*(x) > \alpha\}$  for every  $\alpha > 0$ . Taking  $\alpha \rightarrow \infty$  yields

$$\mu(\{x \mid f^*(x) = \infty\}) = 0.$$

Hence  $f^*(x) < \infty$  a.e.

### Theorem 1.40 (Lebesgue Differentiation Theorem)

Let  $f \in \mathcal{L}^1(\mathbb{R}^d)$ . Then for almost every  $x \in \mathbb{R}^d$ ,

$$\lim_{m(B) \rightarrow 0, x \in B} \frac{1}{m(B)} \int_B f(y) dy = f(x).$$

*Proof.* Since continuous functions are dense in  $\mathcal{L}^1$ , we may find a continuous  $g$  such that  $\|f - g\|_{\mathcal{L}^1} < \epsilon$ . For such  $g$ , by the continuity, there exists a ball such that  $|g(y) - g(x)| < \epsilon$

for all  $x, y \in B$ . Thus

$$\begin{aligned} \left| \frac{1}{m(B)} \int_B f(y) dy - f(x) \right| &= \left| \frac{1}{m(B)} \int_B (f(y) - g(y)) dy + \frac{1}{m(B)} \int_B g(y) - g(x) dy + g(x) - f(x) \right| \\ &\leq \frac{1}{m(B)} \int_B |(f(y) - g(y))| dy + \frac{1}{m(B)} \int_B |g(y) - g(x)| dy + |g(x) - f(x)| \\ &\leq (f - g)^*(x) + \epsilon + |g(x) - f(x)|. \end{aligned}$$

Since  $\epsilon$  can be arbitrary small, we have

$$\left| \frac{1}{m(B)} \int_B f(y) dy - f(x) \right| \leq (f - g)^*(x) + |g(x) - f(x)|.$$

Now we let

$$E_\alpha = \left\{ x \mid \limsup_{m(B) \rightarrow 0, x \in B} \left| \frac{1}{m(B)} \int_B f(y) dy - f(x) \right| > 2\alpha \right\}.$$

We claim that  $E_\alpha$  has measure zero. Set

$$F_\alpha = \{x \mid (f - g)^*(x) > \alpha\} \quad \text{and} \quad G_\alpha = \{x \mid |g(x) - f(x)| > \alpha\}.$$

Then we have  $E_\alpha \subset F_\alpha \cup G_\alpha$ . By the weak-type inequality and Tchebyshev's inequality,

$$\mu(F_\alpha) \leq \frac{A}{\alpha} \|f - g\|_{\mathcal{L}^1} < \frac{A}{\alpha} \epsilon \quad \text{and} \quad \mu(G_\alpha) \leq \frac{1}{\alpha} \|f - g\|_{\mathcal{L}^1} < \frac{1}{\alpha} \epsilon.$$

Thus  $\mu(E_\alpha) \leq \mu(F_\alpha \cup G_\alpha) < \frac{A+1}{\alpha} \epsilon$ . Since  $\epsilon$  is arbitrary, we have  $\mu(E_\alpha) = 0$  and the proof is complete. ■

### Remark

For  $f \in \mathcal{L}^1(\mathbb{R})$ , and  $F(x) = \int_{-\infty}^x f(y) dy$ , we have  $F'(x) = f(x)$  a.e. Indeed,

$$\begin{aligned} \left| \frac{F(x+h) - F(x)}{h} - f(x) \right| &= \frac{1}{h} \left| \int_x^{x+h} f(y) - f(x) dy \right| \leq \frac{1}{h} \int_x^{x+h} |f(y) - f(x)| dy \\ &\leq \frac{1}{h} \int_{x-h}^{x+h} |f(y) - f(x)| dy \leq 2 \frac{1}{2h} \int_{x-h}^{x+h} |f(y) - f(x)| dy \rightarrow 0 \end{aligned}$$

as  $h \rightarrow 0$  a.e.  $x$ .

### Remark

In fact, the requirement that  $f \in \mathcal{L}^1$  can be relaxed to  $f \in \mathcal{L}_{loc}^1$ , which is defined as the set of all locally integrable functions, i.e.,  $f \chi_B \in \mathcal{L}^1$  for all finite balls  $B$  since the proof only requires  $B$  to be a ball near  $x$ .

## 1.4. Radon-Nikodym Theorem

### Definition 1.41

Let  $(X, \mathcal{A})$  be a measurable space. A **signed measure** is a function  $\mu : \mathcal{A} \rightarrow [-\infty, \infty]$  such that  $\mu(\emptyset) = 0$  and for any countable disjoint collection  $\{A_i\}_{i \in \mathbb{N}}$ ,

$$\mu\left(\bigcup_{i \in \mathbb{N}} A_i\right) = \sum_{i \in \mathbb{N}} \mu(A_i).$$

### Remark

The range of  $\mu$  can only include one of  $\pm\infty$ .

### Definition 1.42

Let  $(X, \mathcal{A}, \mu)$  be a measure space.  $\mu$  is called  **$\sigma$ -finite** if  $X$  can be covered by countably many  $A_n \in \mathcal{A}$  such that  $\mu(A_n) < \infty$  for all  $n$ . In this case, we also call  $X$   **$\sigma$ -finite**.

### Definition 1.43

Let  $\nu, \lambda$  be two measures defined on a measurable space.  $\nu$  is said to be **absolutely continuous** with respect to  $\lambda$  if  $\lambda(A) = 0$  implies that  $\nu(A) = 0$  for all measurable  $A$ , denoted as  $\nu \ll \lambda$ .

### Example

Let

$$\nu(A) = \int_A f d\lambda$$

where  $f \geq 0$  is measurable. Then  $\lambda(A) = 0$  implies  $\nu(A) = 0$ .  $\nu \ll \lambda$ .

### Definition 1.44

Let  $\nu, \lambda$  be two measures defined on a measurable space.  $\nu$  is said to be **singular** with respect to  $\lambda$  if there exists a measurable set  $A$  such that  $\lambda(A) = 0$  and  $\nu(A^c) = 0$ , denoted as  $\nu \perp \lambda$ .

### Example

Let  $\lambda$  be the Lebesgue measure on  $[0, 1]$  and

$$\nu(A) = \sum_i c_i \delta_{q_i}(A), \quad \text{with} \quad \sum_i c_i < \infty, \quad \delta_{q_i}(A) = \mathbf{1}\{q_i \in A\},$$

where  $q_i$  enumerates the rationals in  $[0, 1]$  and  $\mathbf{1}$  is the indicator function. Then  $\nu \perp \lambda$ .

### Definition 1.45

$\nu$  and  $\lambda$  are said to be **equivalent** if  $\nu \ll \lambda$  and  $\lambda \ll \nu$ .

### Definition 1.46

Let  $(X, \mathcal{A}, \mu)$  be a measure space. A set  $P \in \mathcal{A}$  is said to be **positive** if  $\mu(A) \geq 0$  for all measurable  $A \subset P$ ; a set  $N \in \mathcal{A}$  is said to be **negative** if  $\mu(A) \leq 0$  for all measurable  $A \subset N$ .



**Theorem 1.47** (Hahn Decomposition)

Let  $\mu$  be a signed measure on a measurable space  $(X, \mathcal{A})$ . Then  $X$  can be partitioned into a positive set  $P$  and a negative set  $N$ . Furthermore, if  $P', N'$  form another such partition, then  $P \Delta P'$  and  $N \Delta N'$  are measure zero.

*Proof.* We may consider the case where  $\mu(A) \neq -\infty$  for all  $A \in \mathcal{A}$ . The other case is similar. We first claim that every measurable set  $A$  contains a positive set  $P$  such that  $\mu(P) \geq \mu(A)$ .

To prove the claim, we first show that for every  $\epsilon > 0$ , there exists  $A_\epsilon \subset A$  such that  $\mu(A_\epsilon) \geq \mu(A)$  and  $B \subset A_\epsilon$  implies  $\mu(B) > -\epsilon$ . Otherwise, we can pick a sequence of set  $B_k$  inductively, such that  $B_1 \subset A, \dots, B_k \subset A - (B_1 \cup \dots \cup B_{k-1}), \dots$  with  $\mu(B_k) \leq -\epsilon$ . Put  $B = \cup_k B_k$ . Since  $B_k$  are disjoint,  $\mu(B) = -\infty$ . Also,  $\mu(A - B) = \mu(A) - \mu(B) = \infty$ , contradicting to the remark that  $\mu$  cannot take both  $\pm\infty$ . Now choose  $\epsilon_n \rightarrow 0$  and let  $P = \cap_n A_{\epsilon_n}$ .  $A_{\epsilon_n} \searrow P$  and then  $\mu(A_{\epsilon_n}) \rightarrow \mu(P)$  by [proposition 1.16](#). Thus  $\mu(P) \geq \mu(A)$ .

Next, let  $s = \sup \{\mu(A) \mid A \in \mathcal{A}\}$ . There is a sequence  $P_n$  such that  $\mu(P_n) \rightarrow s$ . Note that  $s \geq 0$  since  $\emptyset \in \mathcal{A}$ . By the claim, we may assume that  $P_n$  are positive. Putting  $P = \cup_n P_n$ , we have  $\mu(P) = s$  and  $P$  is positive. Now let  $N = X - P$ .  $N$  is negative; otherwise if  $E \subset N$  and  $\mu(E) > 0$ , then  $\mu(P \cup E) = \mu(P) + \mu(E) > s$ , which contradicts to the definition of  $s$ .

Finally, suppose  $P'$  and  $N'$  are another such partition. Then  $P \cap N'$  and  $N \cap P'$  are both negative and positive, implying that they are measure zero.  $\mu(P \Delta P') = \mu(P \cap N') + \mu(N \cap P') = 0$ . This furnishes the proof. ■

**Corollary 1.48** (Hahn-Jordan Decomposition)

If  $\nu$  is a signed measure on a measurable space  $(X, \mathcal{A})$ , then there exists a unique pair of positive measures  $\nu^+$  and  $\nu^-$  such that  $\nu = \nu^+ - \nu^-$ .

*Proof.* By the Hahn decomposition,  $X$  can be partitioned into a positive set  $P$  and a negative set  $N$ . Define  $\nu^+(A) = \nu(A \cap P)$  and  $\nu^-(A) = -\nu(A \cap N)$ . Then  $\nu^+$  and  $\nu^-$  are positive measures and  $\nu = \nu^+ - \nu^-$ . The uniqueness follows from the uniqueness of the Hahn decomposition. ■

**Theorem 1.49** (Radon-Nikodym)

Let  $(X, \mathcal{A})$  be a measurable space and  $\nu, \lambda$  are  $\sigma$ -finite measures on  $(X, \mathcal{A})$ . If  $\nu \ll \lambda$ , then there exists an  $\mathcal{A}$ -measurable function  $f : X \rightarrow [0, \infty)$  such that for every  $A \in \mathcal{A}$ ,

$$\nu(A) = \int_A f d\lambda.$$

Furthermore, if  $f$  and  $f'$  are two such functions, then  $f = f'$  a.e.

*Proof.* We first consider the case where  $\nu$  and  $\lambda$  are finite. Let

$$F = \left\{ f : X \rightarrow [0, \infty) \mid \int_A f d\lambda \leq \nu(A) \text{ for all } A \in \mathcal{A} \right\}.$$

$F \neq \emptyset$  since  $f = 0$  is in  $F$ . Now let  $f_1, f_2 \in F$  and  $A \in \mathcal{A}$  and define

$$A_1 = \{x \in A \mid f_1(x) > f_2(x)\}, \quad A_2 = \{x \in A \mid f_1(x) \leq f_2(x)\}.$$

Then

$$\int_A \max \{f_1, f_2\} d\lambda = \int_{A_1} f_1 d\lambda + \int_{A_2} f_2 d\lambda \leq \nu(A_1) + \nu(A_2) = \nu(A).$$

Thus  $\max \{f_1, f_2\} \in F$ . Next, for any sequence of functions  $f_n \in F$  such that

$$\lim_{n \rightarrow \infty} \int_X f_n d\lambda = \sup_{f \in F} \int_X f d\lambda,$$

we may assume that  $f_n \nearrow$  by replacing  $f_n$  with the maximum among  $f_1, \dots, f_n$ . Let  $g$  be the pointwise limit of  $f_n$ . By Lebesgue's monotone convergence theorem,

$$\int_A g d\lambda = \lim_{n \rightarrow \infty} \int_A f_n d\lambda \leq \nu(A),$$

so  $g \in F$ . Also, by construction,

$$\int_X g d\lambda = \sup_{f \in F} \int_X f d\lambda.$$

Now define

$$\nu_0(A) = \nu(A) - \int_A g d\lambda.$$

Since  $g \in F$ ,  $\nu_0$  is a nonnegative measure. To prove the equality, we need to show that  $\nu_0(A) = 0$  for all  $A \in \mathcal{A}$ . Suppose  $\nu_0 > 0$ . Then there exists  $\epsilon > 0$  such that  $\nu_0(X) > \epsilon\lambda(X)$ . By the Hahn decomposition theorem, we can find a positive set  $P$  such that  $\nu_0(A) \geq \epsilon\lambda(A)$  for each  $A \subset P$ . Thus

$$\nu(A) = \int_A g d\lambda + \nu_0(A) \geq \int_A g d\lambda + \nu_0(P \cap A) \geq \int_A g d\lambda + \epsilon\lambda(P \cap A) = \int_A (g + \epsilon\chi_P) d\lambda.$$

Note that  $\lambda(P) > 0$ , for otherwise  $\lambda(P) = 0$  and  $\nu_0(P) \leq \nu(P) = 0 \implies \nu(P) = 0$  by the absolute continuity and hence

$$\nu_0(X) - \epsilon\lambda(X) = (\nu_0 - \epsilon\lambda)(N) \leq 0,$$

posing a contradiction. Meanwhile,

$$\int_X (g + \epsilon\chi_P) d\lambda \leq \nu(X) < \infty \implies g + \epsilon\chi_P \in F,$$

and

$$\int_X (g + \epsilon\chi_P) d\lambda > \int_X g d\lambda = \sup_{f \in F} \int_X f d\lambda.$$

This violates the definition of the supremum. Thus  $\nu_0 = 0$  and we obtain that

$$\nu(A) = \int_A g d\lambda.$$

Finally, if we define

$$f(x) = \begin{cases} g(x) & \text{if } g(x) < \infty, \\ 0 & \text{if } g(x) = \infty, \end{cases}$$

since  $g$  is  $\lambda$ -integrable,  $f = g$   $\lambda$ -a.e. and  $f$  is the desired function.

For the uniqueness, suppose  $f$  and  $f'$  are two such functions. Then

$$\nu(A) = \int_A f d\lambda = \int_A f' d\lambda \implies \int_A (f - f') d\lambda = 0$$

for every  $A$ . In particular, letting  $A = \{x \in X \mid f(x) \leq f'(x)\}$  or  $A = \{x \in X \mid f(x) \geq f'(x)\}$  gives

$$\int_X (f - f')^+ d\lambda = \int_X (f - f')^- d\lambda = 0.$$

Thus  $f = f'$   $\lambda$ -a.e.

For the general case where  $\nu$  and  $\lambda$  are  $\sigma$ -finite, we can write  $X = \cup_n X_n$  such that  $\lambda(X_n) < \infty$  and  $X_n$  are disjoint. For each  $n$  we can find  $f_n$  such that

$$\nu(A) = \int_A f_n d\lambda.$$

for every  $\mathcal{A}$ -measurable  $A \subset X_n$ . Let  $f = \sum_n f_n \chi_{X_n}$ .

$$\int_A f d\lambda = \sum_n \int_{A \cap X_n} f_n d\lambda = \sum_n \nu(A \cap X_n) = \nu(A),$$

for every  $A \in \mathcal{A}$ . The uniqueness follows from the uniqueness of  $f_n$ . ■

### Remark

The function  $f$  can be chosen in  $\mathcal{L}^1(X, \lambda)$  if  $\nu$  is finite.

### Definition 1.50

The function  $f$  in the Radon-Nikodym theorem is called the **Radon-Nikodym derivative** of  $\nu$  with respect to  $\lambda$ , denoted as  $f = \frac{d\nu}{d\lambda}$ .

### Proposition 1.51

Let  $\nu$ ,  $\mu$  and  $\lambda$  be  $\sigma$ -finite measures defined on measurable space  $(X, \mathcal{A})$ . If  $\nu \ll \lambda$  and  $\mu \ll \lambda$ , then

(a)  $\frac{d(\nu+\mu)}{d\lambda} = \frac{d\nu}{d\lambda} + \frac{d\mu}{d\lambda}$   $\lambda$ -a.e.

(b) If  $\nu \ll \mu \ll \lambda$ , then  $\frac{d\nu}{d\lambda} = \frac{d\nu}{d\mu} \frac{d\mu}{d\lambda}$   $\lambda$ -a.e.

(c) If  $\nu$  and  $\mu$  are equivalent, then  $\frac{d\nu}{d\mu} = \left(\frac{d\mu}{d\nu}\right)^{-1}$   $\mu$ -a.e.

(d) If  $g$  is  $\nu$ -integrable, then

$$\int_X g d\nu = \int_X g \frac{d\nu}{d\lambda} d\lambda.$$

*Proof.* For (a), note that  $\nu + \mu \ll \lambda$  as well. Let  $f = \frac{d\nu}{d\lambda}$  and  $g = \frac{d\mu}{d\lambda}$ . Then

$$\int_A (f + g) d\lambda = \int_A f d\lambda + \int_A g d\lambda = \nu(A) + \mu(A) = (\nu + \mu)(A) = \int_A \frac{d(\nu + \mu)}{d\lambda} d\lambda \quad \text{for all } A \in \mathcal{A}.$$

Thus  $\frac{d\nu}{d\lambda} + \frac{d\mu}{d\lambda} = f + g = \frac{d(\nu + \mu)}{d\lambda}$   $\lambda$ -a.e.

Next, we jump to (d). We start by considering the case where  $g = \chi_A$  with  $A \in \mathcal{A}$ . By the Radon-Nikodym theorem,

$$\int_X g d\nu = \int_X \chi_A d\nu = \nu(A) = \int_A \frac{d\nu}{d\lambda} d\lambda = \int_X \chi_A \frac{d\nu}{d\lambda} d\lambda = \int_X g \frac{d\nu}{d\lambda} d\lambda.$$

By linearity, the result holds for simple functions. For a nonnegative  $g \in \mathcal{L}^1(\nu)$ , we can find a sequence of simple functions  $g_n \nearrow g$  so that

$$\int_X g d\nu = \lim_{n \rightarrow \infty} \int_X g_n d\nu = \lim_{n \rightarrow \infty} \int_X g_n \frac{d\nu}{d\lambda} d\lambda = \int_X g \frac{d\nu}{d\lambda} d\lambda$$

by Lebesgue's monotone convergence theorem. For general  $g \in \mathcal{L}^1(\nu)$ , we can write  $g = g^+ - g^-$  and apply the result to  $g^+$  and  $g^-$ .

$$\int_X g d\nu = \int_X g^+ d\nu - \int_X g^- d\nu = \int_X g^+ \frac{d\nu}{d\lambda} d\lambda - \int_X g^- \frac{d\nu}{d\lambda} d\lambda = \int_X g \frac{d\nu}{d\lambda} d\lambda.$$

With (d) established, we can now prove (b). By the Radon-Nikodym theorem,

$$\int_A \frac{d\nu}{d\mu} \frac{d\mu}{d\lambda} d\lambda = \int_A \frac{d\nu}{d\mu} d\mu = \int_A d\nu = \nu(A) = \int_A \frac{d\nu}{d\lambda} d\lambda.$$

Finally, for (c), letting  $\lambda = \nu$  and applying (b) gives  $1 = \frac{d\nu}{d\nu} = \frac{d\nu}{d\mu} \frac{d\mu}{d\nu}$   $\nu$ -a.e. and thus  $\mu$ -a.e. by the equivalence of  $\nu$  and  $\mu$ . Hence  $\frac{d\nu}{d\mu} = \left(\frac{d\mu}{d\nu}\right)^{-1}$   $\mu$ -a.e. ■

**Theorem 1.52** (Lebesgue Decomposition)

Let  $\nu, \lambda$  be two  $\sigma$ -finite measures defined on a measurable space  $(X, \mathcal{A})$ . Then  $\nu$  can be decomposed uniquely into  $\nu = \nu_a + \nu_s$  where  $\nu_a \ll \lambda$  and  $\nu_s \perp \lambda$ .

*Proof.* We first assume that  $\nu, \lambda$  are finite measures. Let  $\mu = \nu + \lambda$ . Then clearly  $\lambda \ll \mu$  and  $\mu$  is  $\sigma$ -finite. By the Radon-Nikodym theorem, there exists a Radon-Nikodym derivative  $f$  such that

$$\lambda(A) = \int_A f d\mu.$$

Denote  $\{x \in X \mid f(x) = 0\}$  by  $E$ . Define

$$\nu_a(A) = \nu(A \cap E^c), \quad \nu_s(A) = \nu(A \cap E)$$

for each  $A \in \mathcal{A}$ . Then clearly  $\nu_a(A) + \nu_s(A) = \nu(A \cap E^c) + \nu(A \cap E) = \nu(A)$  for all  $A \in \mathcal{A}$ .

Also, suppose  $\lambda(A) = 0$ . Then by [proposition 1.51](#),

$$0 = \lambda(A) = \int_A f d\mu = \int_A f d\lambda + \int_A f d\nu = \int_A f d\nu.$$

Hence  $f(x) = 0$   $\nu$ -a.e. on  $A$ . This implies that  $\nu(A) = \nu(A \cap E)$  and thus  $\nu_a(A) = \nu(A \cap E^c) = \nu(A) - \nu(A \cap E) = 0$ , so  $\nu_a \ll \lambda$ . Also, since  $\lambda(E) = 0$  and  $\nu_s(E^c) = \nu(\emptyset) = 0$ ,  $\nu_s \perp \lambda$ . For the uniqueness, suppose  $\nu = \nu_a + \nu_s = \nu'_a + \nu'_s$  both satisfy the conditions. Since  $\nu_a \ll \lambda$  and  $\nu'_a \ll \lambda$ , by the uniqueness of the Radon-Nikodym derivative,  $\nu_a = \nu'_a$  and hence  $\nu_s = \nu'_s$  as well.

Finally, for the general case where  $\nu, \lambda$  are  $\sigma$ -finite, write  $X = \cup_n X_n$  where  $\lambda(X_n) < \infty$  and  $X_n$  are disjoint. For each  $n$  we can find the corresponding decomposition  $\nu_a^n$  and  $\nu_s^n$ . Let  $\nu_a = \sum_n \nu_a^n$  and  $\nu_s = \sum_n \nu_s^n$ . Then  $\nu_a \ll \lambda$  and  $\nu_s \perp \lambda$ . The uniqueness follows from the uniqueness of the decompositions in each  $X_n$ . This establishes the proof. ■

### Corollary 1.53

*Let  $\nu$  be a signed measure and  $\lambda$  be a measure defined on a measurable space  $(X, \mathcal{A})$ . Suppose both  $\nu$  and  $\lambda$  are finite and  $\nu \ll \lambda$ . Then there exists a unique  $f \in \mathcal{L}^1(X, \lambda)$  such that*

$$\nu(A) = \int_A f d\lambda.$$

*Proof.* By Hahn decomposition, there exists a positive set  $P$  and a negative set  $N$  such that  $P \cup N = X$ . Define

$$\nu_P(A) = \nu(A \cap P), \quad \nu_N(A) = -\nu(A \cap N).$$

Then clearly  $\nu_P - \nu_N = \nu$  and  $|\nu| = \nu_P + \nu_N$ . Note that  $\nu_P$  and  $\nu_N$  are both positive measures. Also, by assumption, if  $\lambda(A) = 0$  then  $\nu(A) = 0$  and hence so are  $\nu_P$  and  $\nu_N$ . Thus  $\nu_P \ll \lambda$  and  $\nu_N \ll \lambda$ . By the Radon-Nikodym theorem, there exists  $f_P, f_N \in \mathcal{L}^1(X, \lambda)$  such that

$$\nu_P(A) = \int_A f_P d\lambda, \quad \nu_N(A) = \int_A f_N d\lambda.$$

Hence

$$\nu(A) = \nu_P(A) - \nu_N(A) = \int_A f_P d\lambda - \int_A f_N d\lambda = \int_A (f_P - f_N) d\lambda.$$

By setting  $f = f_P - f_N$ , we obtain the desired function. Uniqueness follows from the uniqueness of the Radon-Nikodym derivative. ■

## 1.5. Product Measure

### Definition 1.54

*Let  $\mathcal{S}, \mathcal{T}$  be two  $\sigma$ -algebra on  $X$  and  $Y$  respectively. The smallest  $\sigma$ -algebra on  $X \times Y$  containing the collection  $\{S \times T \mid S \in \mathcal{S}, T \in \mathcal{T}\}$  is called the **product  $\sigma$ -algebra** of  $\mathcal{S}$  and  $\mathcal{T}$ , denoted by  $\mathcal{S} \otimes \mathcal{T}$ .*

**Definition 1.55**

Suppose  $X$  is an arbitrary set and  $\mathcal{M}$  is a collection of subsets of  $X$ . We say that  $\mathcal{M}$  is a **monotone class** if

- (a) If  $E_i \subset E_{i+1}$  for countably many  $E_i \in \mathcal{M}$ , then  $\bigcup_{i=1}^{\infty} E_i \in \mathcal{M}$ .
- (b) If  $E_i \supset E_{i+1}$  for countably many  $E_i \in \mathcal{M}$ , then  $\bigcap_{i=1}^{\infty} E_i \in \mathcal{M}$ .

**Definition 1.56**

A collection  $\mathcal{A}$  of subsets in  $X$  is called an **algebra** on  $X$  if

- (a)  $\emptyset \in \mathcal{A}$ .
- (b) If  $A \in \mathcal{A}$ , then  $A^c \in \mathcal{A}$ .
- (c) If  $A_1, A_2 \in \mathcal{A}$ , then  $A_1 \cup A_2 \in \mathcal{A}$ .

**Remark**

The condition (c) implies that for finitely many  $A_i \in \mathcal{A}$ ,  $\bigcup_{i=1}^n A_i \in \mathcal{A}$ .

**Theorem 1.57** (Monotone Class Theorem)

Suppose  $\mathcal{A}$  is an algebra on  $X$ . Then the smallest  $\sigma$ -algebra containing  $\mathcal{A}$  is the smallest monotone class containing  $\mathcal{A}$ .

*Proof.* Let  $\mathcal{M}$  be the smallest monotone class containing  $\mathcal{A}$ . The theorem can be written as  $\sigma(\mathcal{A}) = \mathcal{M}$ . First we show that  $\mathcal{M} \subset \sigma(\mathcal{A})$ . To see this, we claim first that a  $\sigma$ -algebra is automatically a monotone class. Indeed, let  $\mathcal{S}$  be a  $\sigma$ -algebra. Then for any countably many  $E_i \in \mathcal{S}$  with  $E_i \nearrow E$ , we have  $E = \bigcup_{i=1}^{\infty} E_i \in \mathcal{S}$ . Also, for any countably many  $E_i \in \mathcal{S}$  with  $E_i \searrow E$ , we have  $E_i^c \nearrow E^c$ . Thus  $E^c \in \mathcal{S}$  and  $E \in \mathcal{S}$ . Therefore  $\mathcal{S}$  is a monotone class. It follows that  $\sigma(\mathcal{A})$  is a monotone class and hence  $\mathcal{M} \subset \sigma(\mathcal{A})$  by the minimality of  $\mathcal{M}$ .

Next, we claim that  $\mathcal{M}$  is a  $\sigma$ -algebra. By definition, we already have  $\emptyset \in \mathcal{M}$ . Let  $E \in \mathcal{M}$ . Then there is a sequence of sets  $E_i \in \mathcal{A}$  such that either  $E_i \nearrow E$  or  $E_i \searrow E$ . In the former case, we have  $E^c = \bigcap_{i=1}^{\infty} E_i^c \in \mathcal{M}$ ; in the latter case, we have  $E^c = \bigcup_{i=1}^{\infty} E_i^c \in \mathcal{M}$ . Thus  $E^c \in \mathcal{M}$ . Lastly, we need to show that  $\mathcal{M}$  is closed under countable unions. We start by showing that it is closed under finite unions. Consider  $A \in \mathcal{A}$ . Define  $\mathcal{D}_1 = \{D \in \mathcal{M} \mid D \cup A \in \mathcal{M}\}$ . It is clear that  $\mathcal{D}_1$  is a monotone class and  $\mathcal{A} \subset \mathcal{D}_1$ . Consider also  $\mathcal{D}_2 = \{D \in \mathcal{M} \mid D \cup E \in \mathcal{M} \text{ for all } E \in \mathcal{M}\}$ . Then  $\mathcal{D}_2$  is also a monotone class and  $\mathcal{A} \subset \mathcal{D}_2$ . By the minimality of  $\mathcal{M}$ , we have  $\mathcal{M} \subset \mathcal{D}_1 \cap \mathcal{D}_2$  and hence  $\mathcal{M}$  is closed under finite unions. Now let  $E_i \in \mathcal{M}$  be countably many sets. Put  $F_n = \bigcup_{i=1}^n E_i$ . Then  $F_n \nearrow E = \bigcup_i E_i$ . By the closure of  $\mathcal{M}$  under countable unions,  $F_n \in \mathcal{M}$ ; by the definition of  $\mathcal{M}$ ,  $E \in \mathcal{M}$ . We conclude that  $\mathcal{M}$  is closed under countable unions. Thus  $\mathcal{M}$  forms a  $\sigma$ -algebra. It now follows by the minimality of  $\sigma(\mathcal{A})$  that  $\sigma(\mathcal{A}) \subset \mathcal{M}$ . We conclude that  $\sigma(\mathcal{A}) = \mathcal{M}$ . ■

**Lemma 1.58**

Suppose  $(X, \mathcal{S}, \mu)$  and  $(Y, \mathcal{T}, \nu)$  are two finite measure spaces. Let

$$\mathcal{F} = \left\{ E \subset X \times Y \mid \int \int \chi_E(x, y) d\nu(y) d\mu(x) = \int \int \chi_E(x, y) d\mu(x) d\nu(y) \right\}.$$

Then  $\mathcal{S} \otimes \mathcal{T} \subset \mathcal{F}$ .

*Proof.* Since  $\emptyset \in \mathcal{F}$ ,  $\mathcal{F}$  is non-empty. Let  $E = A \times B$  for some  $A \in \mathcal{S}$  and  $B \in \mathcal{T}$ . Then

$$\begin{aligned} \int \int \chi_E(x, y) d\nu(y) d\mu(x) &= \int_A \int_B d\nu(y) d\mu(x) = \nu(B) \int_A d\mu(x) \\ &= \nu(B) \mu(A) = \int_B \mu(A) d\nu(y) \\ &= \int_B \int_A d\mu(x) d\nu(y) = \int \int \chi_E(x, y) d\mu(x) d\nu(y). \end{aligned}$$

Now let  $\mathcal{R}$  be the collection of all rectangles on  $X \times Y$ , i.e.,  $\mathcal{R} = \{A \times B \mid A \in \mathcal{S}, B \in \mathcal{T}\}$ . For  $R_1, R_2 \in \mathcal{R}$ ,  $R_1 \cap R_2 = \emptyset$  implies  $\chi_{R_1 \cup R_2} = \chi_{R_1} + \chi_{R_2}$ . By the above calculation, we know that  $\mathcal{R} \subset \mathcal{F}$ . Consider a sequence of sets  $E_i \in \mathcal{F}$ . If  $E_i \nearrow E$ , then

$$\begin{aligned} \int \int \chi_E(x, y) d\nu(y) d\mu(x) &= \lim_{i \rightarrow \infty} \int \int \chi_{E_i}(x, y) d\nu(y) d\mu(x) \\ &= \lim_{i \rightarrow \infty} \int \int \chi_{E_i}(x, y) d\mu(x) d\nu(y) = \int \int \chi_E(x, y) d\mu(x) d\nu(y). \end{aligned}$$

Also, if  $E_i \searrow E$ , then

$$\begin{aligned} \int \int \chi_E(x, y) d\nu(y) d\mu(x) &= \lim_{i \rightarrow \infty} \int \int \chi_{E_i}(x, y) d\nu(y) d\mu(x) \\ &= \lim_{i \rightarrow \infty} \int \int \chi_{E_i}(x, y) d\mu(x) d\nu(y) = \int \int \chi_E(x, y) d\mu(x) d\nu(y). \end{aligned}$$

Hence  $\mathcal{F}$  is a monotone class containing  $\mathcal{R}$ . By the monotone class theorem,  $\mathcal{S} \otimes \mathcal{T} \subset \mathcal{F}$ . ■

**Theorem 1.59** (Existence and Uniqueness of Product Measure)

Let  $(X, \mathcal{S}, \mu)$  and  $(Y, \mathcal{T}, \nu)$  be two  $\sigma$ -finite measure spaces. Let  $\omega$  be a set function on  $\mathcal{S} \otimes \mathcal{T}$ . For  $A \in \mathcal{S}$  and  $B \in \mathcal{T}$ , define

$$\omega(A \times B) = \mu(A)\nu(B).$$

Then,  $\omega$  extends uniquely to a measure on  $(X \times Y, \mathcal{S} \otimes \mathcal{T})$  such that for every  $E \in \mathcal{S} \otimes \mathcal{T}$ ,

$$\omega(E) = (\mu \times \nu)(E) = \int \int \chi_E(x, y) d\nu(y) d\mu(x) = \int \int \chi_E(x, y) d\mu(x) d\nu(y).$$

*Proof.* If we consider  $\mu$  and  $\nu$  to be  $\sigma$ -finite measures, [lemma 1.58](#) gives us that  $\omega(A \times B) = \mu(A)\nu(B)$ . We want to extend  $\omega$  to a set function

$$\omega(E) = \int \int \chi_E(x, y) d\nu(y) d\mu(x) = \int \int \chi_E(x, y) d\mu(x) d\nu(y).$$

Since integrals are linear,  $\omega$  is finitely additive. Applying the monotone class theorem,  $\omega$  becomes  $\sigma$ -additive. Hence  $\omega$  becomes a measure on  $(X \times Y, \mathcal{S} \otimes \mathcal{T})$ . To see the uniqueness, let  $\rho$  be another measure on  $(X \times Y, \mathcal{S} \otimes \mathcal{T})$  such that  $\rho(A \times B) = \mu(A)\nu(B)$ . Let  $\mathcal{M} =$

$\{E \subset X \times Y \mid \omega(E) = \rho(E)\}$ . For countably many  $E_i \in \mathcal{M}$  with  $E_i \nearrow E$ , we can write  $E = \cup_{i=1}^{\infty} D_i$  where  $D_i = E_{i+1} - E_i$  and  $E_0 = \emptyset$  are disjoint. The  $\sigma$ -additivity gives  $\omega(E) = \rho(E)$ . Thus  $E \in \mathcal{M}$ . A similar argument gives us that  $E_i \searrow E$  implies  $E \in \mathcal{M}$ . Hence  $\mathcal{M}$  is a monotone class. By the monotone class theorem,  $\mathcal{S} \otimes \mathcal{T} \subset \mathcal{M}$ . Thus  $\omega = \rho$ .

For the case  $\mu, \nu$  being  $\sigma$ -finite, consider  $\{A_i\}$  and  $\{B_j\}$  to be two disjoint partitions of  $X$  and  $Y$  respectively with  $\mu(A_i) < \infty$  and  $\nu(B_j) < \infty$  for all  $i, j$ . Let  $E_{ij} = E \cap (A_i \times B_j)$ . By the established result for finite measures,

$$\int \int \chi_{E_{ij}} d\mu d\nu = \int \int \chi_{E_{ij}} d\nu d\mu.$$

Taking the sum over  $i, j$  and applying Lebesgue monotone convergence theorem gives us

$$\omega(E) = \int \int \chi_E d\nu d\mu = \int \int \chi_E d\mu d\nu$$

for any  $E \in \mathcal{S} \otimes \mathcal{T}$ . Applying Lebesgue monotone convergence theorem again results in that  $\omega$  is  $\sigma$ -additive. Hence  $\omega$  is a measure on  $(X \times Y, \mathcal{S} \otimes \mathcal{T})$ . To see the uniqueness, let  $\rho$  be another measure on  $(X \times Y, \mathcal{S} \otimes \mathcal{T})$  such that  $\rho(A \times B) = \mu(A)\nu(B)$ . By the  $\sigma$ -additivity and the uniqueness of the finite measure case,

$$\omega(E) = \sum_{i,j} \omega(E_{ij}) = \sum_{i,j} \rho(E_{ij}) = \rho(E)$$

for all  $E \in \mathcal{S} \otimes \mathcal{T}$ . Thus  $\omega = \rho$ . ■

**Theorem 1.60** (Fubini-Tonelli)

Let  $(X, \mathcal{S}, \mu)$  and  $(Y, \mathcal{T}, \nu)$  be two  $\sigma$ -finite measure spaces. Let  $F : X \times Y \rightarrow \mathbb{R}$  be a  $\mathcal{S} \otimes \mathcal{T}$ -measurable function such that one of the following conditions holds:

- (a)  $F \geq 0$  a.e. (Tonelli);
- (b)  $F$  is integrable (Fubini).

Then

$$\int F(x, y) d(\mu \times \nu) = \int \int F d\mu d\nu = \int \int F d\nu d\mu$$

and furthermore,

$$\begin{cases} y \mapsto \int F(x, y) d\mu(x) & \text{is } \mathcal{T}\text{-measurable,} \\ x \mapsto \int F(x, y) d\nu(y) & \text{is } \mathcal{S}\text{-measurable.} \end{cases}$$

*Proof.* By [theorem 1.59](#), the statement holds for indicator functions and hence for simple functions. By Lebesgue monotone convergence theorem, the non-negative case (Tonelli) is proved. For the integrable case (Fubini), write  $F = F^+ - F^-$ . We also have that  $y \mapsto \int F^\pm(x, y) d\mu(x)$  and  $x \mapsto \int F^\pm(x, y) d\nu(y)$  are  $\mathcal{S}$ -measurable and  $\mathcal{T}$ -measurable by the [theorem 1.59](#). Furthermore,  $y \mapsto \int F^\pm(x, y) d\mu(x)$  and  $x \mapsto \int F^\pm(x, y) d\nu(y)$  are integrable a.e. or



the condition (b) is violated. Thus

$$\begin{aligned}\int F d(\mu \times \nu) &= \int \int F^+ d(\mu \times \nu) - \int \int F^- d(\mu \times \nu) \\ &= \int \int F^+ d\mu d\nu - \int \int F^- d\mu d\nu \\ &= \int \int F^+ d\nu d\mu - \int \int F^- d\nu d\mu.\end{aligned}$$

The proof is complete. ■

### Remark

By induction, one can extend the Lebesgue measure to any  $\mathbb{R}^d$ ,  $d \in \mathbb{N}$ .  $\mathcal{B}_m \otimes \mathcal{B}_n = \mathcal{B}_{m+n}$ , where  $\mathcal{B}_n$  is the Borel  $\sigma$ -algebra on  $\mathbb{R}^n$ . Extension to  $\mathbb{R}^\infty$  is also possible.

## 1.6. Convergence in Measure

### Definition 1.61

Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space. We say that a sequence of function  $f_n$  on  $\Omega$  **converges to a function  $f$  in measure** if for every  $\epsilon > 0$ ,

$$\mu(\{x \in \Omega \mid |f_n(x) - f(x)| \geq \epsilon\}) \rightarrow 0$$

as  $n \rightarrow \infty$ . We write  $f_n \xrightarrow{m} f$ .

### Theorem 1.62 (Markov Inequality)

Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space. For any non-negative measurable function  $f$  on  $\Omega$ ,

$$\mu(\{x \in \Omega \mid f \geq t\}) \leq \frac{1}{t} \int_{\Omega} f d\mu.$$

*Proof.* Let  $E_t = \{x \in \Omega \mid f(x) \geq t\}$ . Then

$$\mu(E_t) = \int \chi_{E_t} d\mu \leq \int \frac{f}{t} d\mu = \frac{1}{t} \int f d\mu.$$

■

### Corollary 1.63 (Chebyshev Inequality)

Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space. For any measurable function  $f$  on  $\Omega$ , and  $\alpha \in \mathbb{R}$ ,

$$\mu(\{x \in \Omega \mid |f(x) - \alpha| \geq t\}) \leq \frac{1}{t^2} \int_{\Omega} (f - \alpha)^2 d\mu.$$

*Proof.* Let  $g = |f - \alpha|^2$ . Apply Markov inequality,

$$\mu(\{x \in \Omega \mid |f(x) - \alpha| \geq t\}) = \mu(\{x \in \Omega \mid g \geq t^2\}) \leq \frac{1}{t^2} \int_{\Omega} g d\mu = \frac{1}{t^2} \int_{\Omega} (f - \alpha)^2 d\mu.$$

■

**Corollary 1.64** (Chernoff Bound)

Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space. For any measurable function  $f$  on  $\Omega$ , and  $\eta \in \mathbb{R}$ ,

$$\mu(\{x \in \Omega \mid f(x) \geq t\}) \leq e^{-\eta t} \int_{\Omega} e^{\eta f} d\mu$$

for all  $t \in \mathbb{R}$ .

*Proof.* Let  $g = e^{\eta f}$ . Then by Markov inequality,

$$\mu(\{x \in \Omega \mid f(x) \geq t\}) = \mu(\{x \in \Omega \mid g \geq e^{\eta t}\}) \leq \frac{1}{e^{\eta t}} \int_{\Omega} g d\mu = e^{-\eta t} \int_{\Omega} e^{\eta f} d\mu.$$

■

**Corollary 1.65**

If  $f_n \rightarrow f$  in  $\mathcal{L}^1$ , then  $f_n \xrightarrow{m} f$ .

*Proof.* Let  $\epsilon > 0$ . By Markov inequality, we have

$$\mu(\{x \in \Omega \mid |f_n(x) - f(x)| \geq \epsilon\}) \leq \frac{1}{\epsilon} \int_{\Omega} |f_n - f| d\mu \rightarrow 0$$

as  $n \rightarrow \infty$ . Thus  $f_n \xrightarrow{m} f$ .

■

**Remark**

The converse is not true. Simply find a sequence of functions converging in  $L^1$  but not almost everywhere will do. However, even stronger, we can actually find a sequence converging in measure but neither in  $\mathcal{L}^1$  nor almost everywhere. For example, let  $\Omega = [0, 1]$  with usual measure. Then let  $f_{k,j} = k^2 \chi_{[\frac{j}{k}, \frac{j+1}{k}]}$  for  $j = 0, 1, \dots, k-1$  and  $k \in \mathbb{N}$ . Reindex the sequence recursively by letting  $g_0 = f_{1,0}$  and

$$g_{n+1} = \begin{cases} f_{k,j+1} & \text{if } g_n = f_{k,j} \text{ with } j \neq k-1, \\ f_{k+1,0} & \text{if } g_n = f_{k,j} \text{ with } j = k-1. \end{cases}$$

This also defines a injective function  $\phi : n \mapsto (k_n, j_n)$ . Then  $g_n \rightarrow 0$  in measure because for any  $\epsilon > 0$ ,

$$\mu(\{x \mid |g_n| \geq \epsilon\}) = \frac{1}{k_n} \rightarrow 0.$$

But

$$\int_0^1 |g_n| d\mu = k_n \rightarrow \infty$$

and since  $[\frac{j_n}{k_n}, \frac{j_n+1}{k_n}]$  includes  $x$  infinitely many times for any  $x \in [0, 1]$ ,  $g_n$  converges nowhere in  $[0, 1]$ .

**Theorem 1.66**

Let  $(\Omega, \mathcal{S}, \mu)$  be a  $\sigma$ -finite measure space. If  $f_n \xrightarrow{m} f$ , then there exists a subsequence  $f_{n_k}$  such that  $f_{n_k} \rightarrow f$  almost everywhere.

*Proof.* Since  $f_n \xrightarrow{m} f$ , we can choose  $n_k$  such that

$$\mu\left(\left\{x \in \Omega \mid |f_n(x) - f(x)| \geq \frac{1}{k}\right\}\right) \leq \frac{1}{2^k} \quad \text{for all } n \geq n_k.$$

Let  $E_k = \{x \in \Omega \mid |f_n(x) - f(x)| \geq \frac{1}{k} \text{ for all } n \geq n_k\}$ . Then  $\mu(E_k) \leq 2^{-k}$ . Put  $H_m = \bigcup_{k=m}^{\infty} E_k$ . We have

$$\mu(H_m) \leq \sum_{k \geq m} \mu(E_k) \leq \sum_{k \geq m} 2^{-k} = 2^{-m+1}.$$

Put  $H = \bigcap_{m=1}^{\infty} H_m$ ,  $H_m \searrow H$ . Then

$$\mu(H) = \lim_{m \rightarrow \infty} \mu(H_m) = 0.$$

If  $x \notin H$ , then  $x \notin H_m$  for some  $m$ . Then

$$|f_{n_k}(x) - f(x)| < \frac{1}{k} \text{ for all } k \geq m.$$

Thus  $f_{n_k}(x) \rightarrow f(x)$  almost everywhere as  $k \rightarrow \infty$ . ■

**Definition 1.67**

Let  $f_n$  be a sequence of measurable functions on  $(\Omega, \mathcal{S}, \mu)$ . We say that  $f_n$  is **Cauchy in measure** if for every  $\epsilon > 0$ ,

$$\mu(\{x \in \Omega \mid |f_n(x) - f_m(x)| \geq \epsilon\}) \rightarrow 0$$

as  $n, m \rightarrow \infty$ .

**Theorem 1.68 (Cauchy Criterion for Convergence in Measure)**

Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space. A sequence of measurable functions  $f_n$  on  $\Omega$  converges in measure if and only if it is Cauchy in measure.

*Proof.* Suppose that  $f_n \xrightarrow{m} f$ . Let  $\epsilon > 0$  be given. We have

$$\mu(\{|f_n - f| \geq \epsilon\}) \rightarrow 0$$

as  $n \rightarrow \infty$ . Then since  $\{|f_n - f_m| \geq \epsilon\} \subset \{|f_n - f| \geq \epsilon/2\} \cup \{|f_m - f| \geq \epsilon/2\}$ ,

$$\mu(\{|f_n - f_m| \geq \epsilon\}) \leq \mu(\{|f_n - f| \geq \epsilon/2\}) + \mu(\{|f_m - f| \geq \epsilon/2\}) \rightarrow 0$$

as  $n, m \rightarrow \infty$ . Thus  $f_n$  is Cauchy in measure.

Conversely, suppose that  $f_n$  is Cauchy in measure. We can take a subsequence  $f_{n_j}$  such that

$$\mu(E_j) = \mu(\{|f_{n_j} - f_{n_{j+1}}| \geq 2^{-j}\}) \leq 2^{-j}.$$

Put  $F_k = \cup_{j=k}^{\infty} E_j$ . Then  $\mu(F_k) \leq \sum_{j=k}^{\infty} \mu(E_j) \leq 2^{-k+1}$ . For  $x \notin F_k, i > j$ ,

$$|f_{n_i} - f_{n_j}| \leq \sum_{l=j}^{i-1} |f_{n_{l+1}} - f_{n_l}| \leq \sum_{l=j}^{i-1} 2^{-l} \leq 2^{-j+1}.$$

Hence  $f_{n_j}$  is Cauchy on  $F_k^c$ . By the completeness of  $\mathbb{R}$ ,  $f_{n_j}$  converges pointwise on  $F_k^c$  for each  $k$ . Put  $F = \cap_{k=1}^{\infty} F_k$ . Then  $\mu(F) = 0$ . Let

$$f(x) = \begin{cases} \lim_{j \rightarrow \infty} f_{n_j}(x) & \text{if } x \notin F, \\ 0 & \text{if } x \in F. \end{cases}$$

Since  $f_{n_j}$  are measurable,  $f$  is measurable. Also,  $f_{n_j} \rightarrow f$  pointwisely almost everywhere. Thus  $f_{n_j} \xrightarrow{m} f$ . Observe that  $\{|f_n - f| \geq \epsilon\} \subset \{|f_n - f_{n_j}| \geq \epsilon/2\} \cup \{|f_{n_j} - f| \geq \epsilon/2\}$ . Hence

$$\mu(\{|f_n - f| \geq \epsilon\}) \leq \mu(\{|f_n - f_{n_j}| \geq \epsilon/2\}) + \mu(\{|f_{n_j} - f| \geq \epsilon/2\}) \rightarrow 0$$

as  $n \rightarrow \infty$ . Thus  $f_n \xrightarrow{m} f$ . ■

### Definition 1.69

A function  $\phi : (a, b) \rightarrow \mathbb{R}$ , where  $-\infty \leq a < b \leq \infty$ , is **convex** if for any  $x, y \in (a, b)$  and  $\lambda \in [0, 1]$ ,

$$\phi(\lambda x + (1 - \lambda)y) \leq \lambda\phi(x) + (1 - \lambda)\phi(y).$$

### Remark

Every convex function is continuous.

### Remark

The definition of convexity can also be written as

$$\frac{\phi(t) - \phi(s)}{t - s} \leq \frac{\phi(u) - \phi(t)}{u - t},$$

whenever  $a < s < t < u < b$ .

### Theorem 1.70 (Jensen's Inequality)

Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space with  $\mu(\Omega) = 1$ . Suppose that  $f : \Omega \rightarrow I$ ,  $f \in \mathcal{L}^1(\Omega)$  and  $\phi : I \rightarrow \mathbb{R}$  is a convex function on an interval  $I$ . Then

$$\phi\left(\int_{\Omega} f d\mu\right) \leq \int_{\Omega} \phi(f) d\mu.$$

*Proof.* Put  $t = \int_{\Omega} f d\mu$ . Then  $a < t < b$ . Let

$$\beta = \sup_{s \in (a, t)} \frac{\phi(t) - \phi(s)}{t - s}.$$

By the convexity,

$$\beta \leq \frac{\phi(u) - \phi(t)}{u - t}$$

for any  $u \in (t, b)$ . Thus

$$\phi(y) \geq \phi(t) + \beta(y - t)$$

for all  $y \in (a, b)$ . Hence

$$\phi(f(x)) - \phi(t) - \beta(f(x) - t) \geq 0$$

for every  $x \in \Omega$ . Since  $\phi$  is continuous,  $\phi \circ f$  is measurable. Thus

$$\int_{\Omega} \phi(f) d\mu - \phi(t) = \int_{\Omega} \phi(f) d\mu - \phi(t) - \beta \left( \int_{\Omega} f d\mu - t \right) = \int_{\Omega} \phi(f) d\mu - \phi(t) - \beta \int_{\Omega} (f - t) d\mu \geq 0.$$

Since  $t = \int_{\Omega} f d\mu$ , we have

$$\phi \left( \int_{\Omega} f d\mu \right) \leq \int_{\Omega} \phi(f) d\mu.$$

■

### Definition 1.71

A family of measure  $\{\nu_{\alpha}\}$  is said to be **equicontinuous at  $\emptyset$**  if for any  $\epsilon > 0$  and  $B_k \searrow \emptyset$ , there exists  $k_0$  such that

$$\sup_{\alpha} \nu_{\alpha}(B_k) < \epsilon$$

for all  $k \geq k_0$ .

### Definition 1.72

A family of measure  $\{\nu_{\alpha}\}$  is said to be **uniformly absolutely continuous** with respect to  $\mu$  if for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that for any  $B$  with  $\mu(B) < \delta$ ,

$$\sup_{\alpha} \nu_{\alpha}(B) < \epsilon.$$

### Lemma 1.73

If  $\{\nu_{\alpha}\}$  is equicontinuous at  $\emptyset$  and  $\nu_{\alpha} \ll \mu$  for all  $\alpha$ , then  $\{\nu_{\alpha}\}$  is uniformly absolutely continuous with respect to  $\mu$ .

*Proof.* Suppose that  $\{\nu_{\alpha}\}$  is not uniformly absolutely continuous with respect to  $\mu$ . Then there exists  $\epsilon > 0$  such that for any  $n$ , we can find  $B_n$  with  $\mu(B_n) \leq 2^{-n}$  and some  $\alpha_n$  with  $\nu_{\alpha_n}(B_n) \geq \epsilon$ . Put  $A_k = \cup_{n=k}^{\infty} B_n$ . Then  $\mu(A_k) \leq 2^{-k+1}$ . Set  $A = \cap_{k=1}^{\infty} A_k$ . Then  $A_k \searrow A$  and  $\mu(A) = 0$ . This implies  $\nu_{\alpha}(A) = 0$  for all  $\alpha$  since  $\nu_{\alpha} \ll \mu$ . Observe now that

$$\nu_{\alpha_n}(A_k - A) = \nu_{\alpha_n}(A_k) \geq \nu_{\alpha_n}(B_n) \geq \epsilon$$

for all  $n \geq k$ . But  $\nu_{\alpha_n}(A_k - A) \rightarrow 0$  as  $k \rightarrow \infty$ , a contradiction. Thus  $\{\nu_{\alpha}\}$  is uniformly absolutely continuous with respect to  $\mu$ . ■

**Theorem 1.74**

Let  $(\Omega, \mathcal{S}, \mu)$  be a  $\sigma$ -finite measure space. Suppose  $f_n \in \mathcal{L}^p(\Omega)$ . Consider a family of measures  $\nu_n$  defined by

$$\nu_n(A) = \int_A |f_n|^p d\mu.$$

If  $\nu_n$  is equicontinuous at  $\emptyset$  and  $f_n \xrightarrow{m} f$ , then  $f_n \rightarrow f$  in  $\mathcal{L}^p(\Omega)$ .

*Proof.* Since  $(\Omega, \mathcal{S}, \mu)$  is  $\sigma$ -finite, we can write  $\Omega = \cup_k E_k$  with  $\mu(E_k) < \infty$  for all  $k$ . Then  $E_k^c \searrow \emptyset$  and  $\nu_n(E_k^c) \rightarrow 0$  as  $k \rightarrow \infty$ . Also, since  $\nu_n$  is equicontinuous at  $\emptyset$ , for any  $\epsilon > 0$ , there exists  $k_0$  such that

$$\sup_n \nu_n(E_k^c) < \epsilon$$

for all  $k \geq k_0$ .

We claim that  $f_n$  is Cauchy in  $\mathcal{L}^p$ . Indeed,

$$\begin{aligned} \int |f_n - f_m|^p d\mu &= \int_{E_{k_0}^c} |f_n - f_m|^p d\mu + \int_{E_{k_0} \cap \{|f_n - f_m| \leq \epsilon/\mu(E_{k_0})\}} |f_n - f_m|^p d\mu \\ &\quad + \int_{E_{k_0} \cap \{|f_n - f_m| > \epsilon/\mu(E_{k_0})\}} |f_n - f_m|^p d\mu. \end{aligned}$$

Estimate from Jensen's inequality,

$$\int_{E_{k_0}^c} |f_n - f_m|^p d\mu \leq 2^p \int_{E_{k_0}^c} |f_n|^p d\mu + 2^p \int_{E_{k_0}^c} |f_m|^p d\mu = 2^p \nu_n(E_{k_0}^c) + 2^p \nu_m(E_{k_0}^c) \rightarrow 0,$$

$$\int_{E_{k_0} \cap \{|f_n - f_m| \leq \epsilon/\mu(E_{k_0})\}} |f_n - f_m|^p d\mu \leq \frac{\epsilon}{\mu(E_{k_0})} \mu(E_{k_0}) \rightarrow 0.$$

For the last term, since  $\nu_n \ll \mu$  for all  $n$ , [lemma 1.73](#) gives that  $\nu_n$  is uniformly absolutely continuous with respect to  $\mu$ . Given any  $\epsilon > 0$ , there is  $\delta > 0$  such that for all  $B$  with  $\mu(B) \leq \delta$ ,  $\nu_n(B) \leq \epsilon$  for all  $n$ . Thus

$$\mu\left(\left\{|f_j - f| \geq \frac{\epsilon}{\mu(E_{k_0})}\right\}\right) \rightarrow 0$$

as  $j \rightarrow \infty$ . Hence we obtain that  $f_n$  is Cauchy in  $\mathcal{L}^p$ . It follows from the Riesz-Fischer theorem that  $f_n \rightarrow g$  in  $\mathcal{L}^p(\Omega)$  for some  $g \in \mathcal{L}^p(\Omega)$ . Since  $f_n \xrightarrow{m} f$ ,  $f = g$  almost everywhere. Thus  $f_n \rightarrow f$  in  $\mathcal{L}^p(\Omega)$ . ■

## 2. Banach Space

### 2.1. Banach Space and Bounded Linear Functional

#### Definition 2.1

A space  $X$  is called a **Banach space** if it is a complete normed vector space.

#### Remark

$\mathcal{L}^1$  is a Banach space with the norm

$$\|f\|_{\mathcal{L}^1} = \int |f| d\mu.$$

We treat  $f = g$  a.e. as the same element in  $\mathcal{L}^1$ .

#### Definition 2.2

Let  $V, W$  be vector spaces. A map  $T : V \rightarrow W$  is **linear** if for every  $c \in \mathbb{R}$ ,  $f, g \in V$ ,  $T(cf + g) = cT(f) + T(g)$ .

#### Definition 2.3

A linear map  $T : V \rightarrow W$  has **operator norm** defined by

$$\|T\| = \sup_{\|f\|_V=1} \|T(f)\|_W.$$

$T$  is **bounded** if  $\|T\| < \infty$ . We denote the set of all bounded linear operators from  $V$  to  $W$  by  $B(V, W)$ .

#### Proposition 2.4

Suppose  $W$  is a Banach space. Then  $B(V, W)$  is a Banach space with the operator norm.

*Proof.* It suffices to show that  $B(V, W)$  is complete. Let  $\{T_i\} \subset B(V, W)$  be a Cauchy sequence. Then for  $f \in V$ ,

$$\|T_i(f) - T_j(f)\|_W \leq \|T_i - T_j\| \|f\|_V.$$

Hence  $\{T_i(f)\}$  is a Cauchy sequence in  $W$ . By the completeness of  $W$ , we may define  $Tf$  as the limit of  $T_i(f)$  as  $i \rightarrow \infty$ . Now,

$$\|Tf\| \leq \sup_i \|T_i(f)\| \leq \sup_i \|T_i\| \|f\|.$$

Since Cauchy sequences are bounded,  $\|Tf\| < \infty$  for all  $f \in V$  and  $T \in B(V, W)$ . It remains to show that  $T_i$  converges to  $T$  in the operator norm. For any  $f \in V$ , pick  $N$  such that  $\|T_i(f) - T_j(f)\| \leq \epsilon$  for all  $i, j \geq N$ . Then for fixed  $i$ ,

$$\|(T_i - T_j)f\| \leq \|T_i - T_j\| \|f\| \leq \epsilon \|f\|$$

for every  $f \in V$  and  $j \geq N$ . Hence  $\|T_i - T\| \leq \epsilon$  for all  $i \geq N$  and the proof is complete. ■

**Definition 2.5**

$T$  is continuous if  $f_i \rightarrow f$  in  $V$  implies that  $T(f_i) \rightarrow T(f)$  in  $W$ .

**Proposition 2.6**

Suppose  $T : V \rightarrow W$  is linear. Then  $T$  is continuous if and only if  $T$  is bounded.

*Proof.* Suppose  $T$  is not bounded. Then there exists  $f_i \in V$  with  $\|f_i\| \leq 1$  for all  $i$  and  $\|Tf_i\| \rightarrow \infty$ . Thus

$$\frac{f_i}{\|Tf_i\|} \rightarrow 0, \quad \text{but} \quad T\left(\frac{f_i}{\|Tf_i\|}\right) = \frac{Tf_i}{\|Tf_i\|} \not\rightarrow 0 \quad \text{as} \quad \frac{\|Tf_i\|}{\|Tf_i\|} = 1.$$

Hence  $T$  is not continuous.

Conversely, suppose  $T$  is bounded. Let  $f_i \rightarrow f$  in  $V$ . Then

$$\|Tf_i - Tf\| = \|T(f_i - f)\| \leq \|T\| \|f_i - f\| \rightarrow 0.$$

Hence  $T$  is continuous. ■

**Definition 2.7**

A **linear functional**  $T$  is a linear map  $T : V \rightarrow \mathbb{F}$ , where  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$  is the scalar field of  $V$ .

**Definition 2.8**

Let  $V, W$  be vector spaces.  $T : V \rightarrow W$  is linear. The **kernel** of  $T$  is defined as

$$\ker(T) = \{f \in V \mid T(f) = 0\}.$$

**Proposition 2.9**

Let  $X$  be a normed vector space and  $T \in X'$ . Then

- (a)  $\ker(T)$  is a closed subspace of  $X$ .
- (b) If  $T \neq 0$ , there exists  $x \in X$  such that  $T(x) \neq 0$ . Then for any  $y \in X$ , there exists  $c \in \mathbb{R}$  and  $z \in \ker(T)$  such that  $y = cx + z$ .

*Proof.* For (a), let  $x, y \in \ker(T)$  and  $c \in \mathbb{R}$ .

$$T(cx + y) = cT(x) + T(y) = 0. \implies cx + y \in \ker(T).$$

Also, let  $x_i \rightarrow x$  in  $X$ . Then since  $T$  is continuous,

$$T(x) = \lim_{n \rightarrow \infty} T(x_n) = 0. \implies x \in \ker(T).$$

Hence  $\ker(T)$  is a closed subspace of  $X$ .

For the rest part, fix  $x \in X$  and  $f(x) \neq 0$ . For each  $y \in X$ , let  $\alpha = T(y)/T(x)$  and  $z = y - T(y)x/T(x)$ . Then

$$\alpha x + z = \frac{T(y)}{T(x)}x + y - \frac{T(y)}{T(x)}x = y - \frac{T(y)}{T(x)}x = y.$$



■

### Definition 2.10

The **dual space** of  $V$  is defined as  $V' = B(V, \mathbb{F})$ , where  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .

### Remark

The dual space is a Banach space.

### Remark

$T : X \rightarrow Y$  is bounded and linear. Then

$$\|T\| = \inf \{c \in [0, \infty) \mid \|Tx\|_Y \leq c \|x\|_X \text{ for all } x \in X\}.$$

### Example

Let  $X = C([0, 1])$  with the supremum norm and  $Y = \mathbb{R}$  with the usual norm. For  $g \in X$ ,  $g(t) \neq 0$  on  $[0, 1]$ , define  $Tg : X \rightarrow \mathbb{R}$  by

$$Tg(f) = \int_0^1 f(t)g(t)dt.$$

Now for  $\|f\|_\infty \leq 1$ ,

$$\begin{aligned} |Tg(f)| &= \left| \int_0^1 f(t)g(t)dt \right| \leq \int_0^1 |f(t)g(t)| dt \leq \int_0^1 |g(t)| \sup_{[0,1]} |f(t)| dt \\ &= \|f\|_\infty \int_0^1 |g(t)| dt \leq \int_0^1 |g(t)| dt. \end{aligned}$$

Take  $f = g/|g|$ ,

$$|Tgf| = \left| \int_0^1 \frac{g^2(t)}{|g(t)|} dt \right| = \int_0^1 |g(t)| dt. \implies \|Tg\| = \int_0^1 |g(t)| dt.$$

### Example

Consider  $X = Y = C([0, 1])$  with the supremum norm. Define  $T : C^1([0, 1]) \rightarrow Y$  by  $Tf = f'$ . Then consider the sequence  $f_n(x) = e^{-n(x-1/2)^2}$ ,  $f'_n(x) = e^{-n(x-1/2)^2}(-2n(x-1/2))$ . Hence  $\|Tf_n\| / \|f_n\| = \sqrt{2n}e^{-1/2} \rightarrow \infty$  as  $n \rightarrow \infty$ . Thus  $T$  is not bounded.

## 2.2. $\ell^p$ Space

### Definition 2.11

$\ell^p = \{\{x_i\}_{i \in I} \mid \|x\|_p < \infty\}$ , where  $I$  is an countable index set and  $\|x\|_p = (\sum_i |x_i|^p)^{1/p}$ ,  $1 \leq p < \infty$ , is called the  $\ell^p$  **space**. For  $p = \infty$ , the norm is defined as  $\|x\|_\infty = \sup_i |x_i|$ .

### Definition 2.12

$f : X \rightarrow Y$  is called a **homomorphism** if it preserves the algebraic structure. In particular, for  $X, Y$  being vector spaces,  $f$  is a homomorphism if  $f(cx + y) = cf(x) + f(y)$ .

**Definition 2.13**

$f : X \rightarrow Y$  is called an **isomorphism** if it is a bijective homomorphism.

**Definition 2.14**

$f : X \rightarrow Y$  is called an **isometry** if  $\|f(x)\|_Y = \|x\|_X$  for all  $x \in X$ .

**Example**

A rightward shift operator  $S_R : \ell^p(\mathbb{N}) \rightarrow \ell^p(\mathbb{N})$  is not an isomorphism, but  $S_R : \ell^p(\mathbb{Z}) \rightarrow \ell^p(\mathbb{Z})$  is.

**Lemma 2.15** (Young's Inequality)

Let  $1 < p, p' < \infty$  with  $\frac{1}{p} + \frac{1}{p'} = 1$ . Then for all  $a, b \geq 0$ ,

$$ab \leq \frac{a^p}{p} + \frac{b^{p'}}{p'}.$$

Furthermore, the equality holds if and only if  $a^p = b^{p'}$ .

*Proof.* If  $a = 0$  or  $b = 0$ , the inequality is trivial. Suppose  $a, b > 0$ . Let  $t = 1/p$  and we can write

$$\log(ab) = \log(a) + \log(b) = t \log(a^p) + (1-t) \log(b^{p'}) \leq \log\left(ta^p + (1-t)b^{p'}\right)$$

by the concavity of logarithm and Jensen's inequality. Exponentiating both sides yields the desired inequality. The equality holds if and only if  $a^p = b^{p'}$  by the Jensen's inequality. ■

**Theorem 2.16** (Hölder's Inequality in  $\ell^p$ )

Let  $1 \leq p, p' \leq \infty$  with  $\frac{1}{p} + \frac{1}{p'} = 1$ . Then for all  $f \in \ell^p$  and  $g \in \ell^{p'}$ ,

$$\|fg\|_1 \leq \|f\|_p \|g\|_{p'}.$$

Moreover, the equality holds if and only if  $f = cg$  for some constant  $c$ .

*Proof.* If one of  $f$  or  $g$  is zero, the inequality is trivial. If  $p = 1$  and  $p' = \infty$ ,  $|f_i g_i| \leq \|g\|_\infty |f_i|$ . Summing over  $i$  yields the desired inequality. For the case  $p = \infty$  and  $p' = 1$  the proof is similar. Now suppose  $1 < p < \infty$  and  $1 < p' < \infty$ . Without loss of generality, we may assume that  $\|f\|_p = \|g\|_{p'} = 1$ . By Young's inequality,

$$|f_i g_i| \leq \frac{|f_i|^p}{p} + \frac{|g_i|^{p'}}{p'}.$$

Thus

$$\|fg\|_1 = \sum_i |f_i g_i| \leq \sum_i \frac{|f_i|^p}{p} + \sum_i \frac{|g_i|^{p'}}{p'} = \frac{1}{p} \|f\|_p^p + \frac{1}{p'} \|g\|_{p'}^{p'} = 1.$$

Hence we obtain the desired inequality. The equality holds if and only if  $|f_i|^p = |g_i|^{p'}$  for all  $i$  by the Young's inequality. In general, the equality holds if and only if  $f = cg$  for some constant  $c$  after scaling the both sides of the inequality by  $c$ . ■

### Remark

We call  $p'$  the **conjugate exponent** of  $p$  for  $1/p + 1/p' = 1$ .

### Theorem 2.17 (Minkowski's Inequality in $\ell^p$ )

Let  $1 \leq p \leq \infty$ . Then for all  $f, g \in \ell^p$ ,

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

*Proof.* If  $p = 1$ , the inequality comes from the triangle inequality. For  $1 < p < \infty$ ,

$$\begin{aligned} \|f + g\|_p^p &= \sum_i |f_i + g_i| |f_i + g_i|^{p-1} \\ &\leq \sum_i |f_i| |f_i + g_i|^{p-1} + \sum_i |g_i| |f_i + g_i|^{p-1} \\ &\leq \|f\|_p \left( \sum_i |f_i + g_i|^{(p-1)p'} \right)^{1/p'} + \|g\|_p \left( \sum_i |f_i + g_i|^{(p-1)p'} \right)^{1/p'} \\ &= \|f\|_p \|f + g\|_p^{p/p'} + \|g\|_p \|f + g\|_p^{p/p'} \end{aligned}$$

by the Hölder's inequality. Rearranging the inequality yields

$$\|f + g\|_p = \|f + g\|_p^{p-p/p'} \leq \|f\|_p + \|g\|_p.$$

For  $p = \infty$ ,

$$\|f + g\|_\infty = \sup_i |f_i + g_i| \leq \sup_i |f_i| + \sup_i |g_i| = \|f\|_\infty + \|g\|_\infty.$$

The proof is complete. ■

### Remark

The Minkowski's inequality is exactly the triangle inequality in  $\ell^p$  spaces. We can thus confirm that  $\ell^p$  norms are indeed norms.

## 2.3. $\mathcal{L}^p$ Space

### Definition 2.18

Let  $(X, \mathcal{A}, \mu)$  be a measure space and  $1 \leq p < \infty$ . The space  $\mathcal{L}^p(X)$  consists of all equivalence classes of measurable functions  $f : X \rightarrow \mathbb{R}$  such that

$$\|f\|_{\mathcal{L}^p} = \left( \int_X |f|^p d\mu \right)^{1/p} < \infty,$$

where  $f \sim g$  if  $f = g$  a.e. and the norm is defined on a representative of the equivalence class.

**Definition 2.19**

$f : X \rightarrow \mathbb{R}$  is measurable. The **essential supremum** of  $f$  on  $X$  is defined as

$$\operatorname{ess\,sup}_X f = \inf \left\{ \sup_X g \mid g = f \mu\text{-a.e.} \right\} = \inf \{ c \in \mathbb{R} \mid \mu(\{x \mid f(x) > c\}) = 0 \}.$$

We called  $f$  **essentially bounded** if  $\operatorname{ess\,sup}_X f < \infty$ . The space  $\mathcal{L}^\infty(X)$  consists of all equivalence classes of essentially bounded measurable functions with the norm

$$\|f\|_{\mathcal{L}^\infty} = \operatorname{ess\,sup}_X |f|.$$

**Theorem 2.20** (Hölder's Inequality in  $\mathcal{L}^p$ )

Let  $1 \leq p, p' \leq \infty$  with  $\frac{1}{p} + \frac{1}{p'} = 1$ . Then for all  $f \in \mathcal{L}^p$  and  $g \in \mathcal{L}^{p'}$ ,

$$\|fg\|_1 \leq \|f\|_p \|g\|_{p'}.$$

Moreover, the equality holds if and only if  $f = cg$  for some constant  $c$ .

*Proof.* For the case  $p = 1$  and  $p' = \infty$ , notice that

$$|fg| \leq |f| \operatorname{ess\,sup} |g| \implies \|fg\|_1 = \int |fg| d\mu \leq \int |f| \operatorname{ess\,sup} |g| d\mu = \|f\|_1 \|g\|_\infty.$$

For the case  $p = \infty$  and  $p' = 1$ , the proof is similar. Now suppose  $1 < p < \infty$  and  $1 < p' < \infty$ . If one of  $f$  or  $g$  is zero, the inequality is trivial. Without loss of generality, we may assume that  $\|f\|_p = \|g\|_{p'} = 1$ . By the Young's inequality,

$$|fg| \leq \frac{|f|^p}{p} + \frac{|g|^{p'}}{p'}.$$

Integrating both sides yields

$$\|fg\|_1 = \int |fg| d\mu \leq \int \frac{|f|^p}{p} d\mu + \int \frac{|g|^{p'}}{p'} d\mu = \frac{1}{p} + \frac{1}{p'} = 1.$$

Hence we obtain the desired inequality. The equality holds if and only if  $|f|^p = |g|^{p'}$  a.e. by the Young's inequality. In general, the equality holds if and only if  $f = cg$  a.e. for some constant  $c$  after scaling the both sides of the inequality by  $c$ . ■

**Theorem 2.21** (Minkowski's Inequality in  $\mathcal{L}^p$ )

Let  $1 \leq p \leq \infty$ . Then for all  $f, g \in \mathcal{L}^p$ ,

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

*Proof.* If  $p = 1$ , the inequality comes from the triangle inequality. For  $1 < p < \infty$ ,

$$\begin{aligned}
\|f + g\|_p^p &= \int |f + g|^p d\mu = \int |f + g| |f + g|^{p-1} d\mu \\
&\leq \int |f| |f + g|^{p-1} d\mu + \int |g| |f + g|^{p-1} d\mu \\
&\leq \|f\|_p \left( \int |f + g|^{(p-1)p'} d\mu \right)^{1/p'} + \|g\|_p \left( \int |f + g|^{(p-1)p'} d\mu \right)^{1/p'} \\
&= \|f\|_p \|f + g\|_p^{p/p'} + \|g\|_p \|f + g\|_p^{p/p'}.
\end{aligned}$$

Rearranging the inequality yields

$$\|f + g\|_p = \|f + g\|_p^{p-p/p'} \leq \|f\|_p + \|g\|_p.$$

For  $p = \infty$ ,

$$\|f + g\|_\infty = \text{ess sup } |f + g| \leq \text{ess sup } |f| + \text{ess sup } |g| = \|f\|_\infty + \|g\|_\infty.$$

The proof is complete. ■

### Theorem 2.22

$1 \leq p \leq \infty$ . Simple functions are dense in  $\mathcal{L}^p$ .

*Proof.* For  $p < \infty$ , consider  $f \geq 0$  and  $f \in \mathcal{L}^1$ . There exists a sequence of simple functions  $f_n \nearrow f$  a.e. Note that  $|f - f_n|^p \leq |f|^p \in \mathcal{L}^1$ . By Lebesgue's dominated convergence theorem,  $\|f_n - f\|_p \rightarrow 0$  as  $n \rightarrow \infty$ . For  $p = \infty$ , pick an  $f$  in the  $f$ -equivalent class such that  $f$  is bounded. Then since the approximation of simple functions can be done uniformly, the result follows. ■

### Remark

A simple function  $s = \sum_{i=1}^n c_i \chi_{A_i} \in \mathcal{L}^p$  must have  $\mu(A_i) < \infty$  for every  $i$  such that  $c_i > 0$ . Since continuous functions can approximate simple functions, they are dense in  $\mathcal{L}^p$  as well.

### Remark

Step functions and continuous functions with compact supports are dense in  $\mathcal{L}^p$  for  $1 \leq p < \infty$ . This can be seen by a slight modification of the proof of [proposition 1.35](#). Let  $\epsilon > 0$  be given. First, for  $f \in \mathcal{L}^p$ , we can find some  $M > 0$  such that

$$\int_{|x|>M} |f|^p d\mu < \epsilon.$$

Next, since  $\mathcal{L}^p([-M, M]) \subset \mathcal{L}^1([-M, M])$ , the result from [proposition 1.35](#) applies, and we can find a step function  $s$  such that  $\|f - s\|_\infty < \epsilon$  on  $[-M, M]$ . This implies

$$\int_{|x|\leq M} |f - s|^p d\mu \leq \int_{|x|\leq M} \epsilon^p d\mu = \epsilon^p \mu([-M, M]).$$

Thus

$$\|f - s\|_p^p = \int_{|x|>M} |f|^p d\mu + \int_{|x|\leq M} |f - s|^p d\mu \leq \epsilon + \epsilon^p \mu([-M, M]).$$

Hence step functions with compact supports are dense in  $\mathcal{L}^p$ . Using the same approximation technique in [proposition 1.35](#), we can find a continuous function  $g$  such that  $\|f - g\|_p < \epsilon$  as well.

**Lemma 2.23**

$1 \leq p < \infty$ .  $g_k \in \mathcal{L}^p$  and  $\sum_k \|g_k\|_p < \infty$ . Then there exists  $f \in \mathcal{L}^p$  such that  $\sum_k g_k = f$  pointwise a.e. and in  $\mathcal{L}^p$ .

*Proof.* Define  $h_n$  and  $h$  by  $h_n = \sum_{k=1}^n |g_k|$  and  $h = \sum_k |g_k|$ . Then  $h_n \nearrow h$ . By Lebesgue's monotone convergence theorem,

$$\lim_{n \rightarrow \infty} \int h_n^p d\mu = \int h^p d\mu.$$

By Minkowski's inequality,

$$\left( \int h_n^p d\mu \right)^{1/p} = \left( \int \left( \sum_{k=1}^n |g_k| \right)^p d\mu \right)^{1/p} \leq \sum_{k=1}^n \left( \int |g_k|^p d\mu \right)^{1/p} \leq \sum_{k=1}^n \|g_k\|_p < \infty$$

for every  $n$ , so  $h \in \mathcal{L}^p$  and  $\|h\|_p \leq M$  for some  $M$  bounding  $\sum_k \|g_k\|_p$ . Now since  $\sum_k g_k$  converges absolutely to some  $f$  pointwisely a.e. and  $|f| \leq h$ ,

$$\left| f - \sum_{k=1}^n g_k \right|^p \leq \left( |f| + \sum_{k=1}^n |g_k| \right)^p \leq (2h)^p \in \mathcal{L}^1.$$

By Lebesgue's dominated convergence theorem,  $\|f - \sum_{k=1}^n g_k\|_p \rightarrow 0$  as  $n \rightarrow \infty$ . Thus the proof is complete. ■

**Theorem 2.24 (Riesz-Fischer)**

$\mathcal{L}^p$  spaces are complete.

*Proof.* First, we focus on the case where  $1 \leq p < \infty$ . Let  $f_k$  be a Cauchy sequence in  $\mathcal{L}^p$ . Take a subsequence  $f_{k_j}$  such that  $\|f_{k_{j+1}} - f_{k_j}\| \leq 2^{-j}$ . Let  $g_j = f_{k_{j+1}} - f_{k_j} \in \mathcal{L}^p$  and we have  $\sum_j \|g_j\|_p < \infty$ . By the [lemma 2.23](#), there exists  $f \in \mathcal{L}^p$  such that  $f = \sum_j g_j$  a.e. and

$$\lim_{j \rightarrow \infty} f_{k_j} = \lim_{j \rightarrow \infty} f_{k_1} + \sum_{i=1}^{j-1} g_i = f_{k_1} + f \in \mathcal{L}^p.$$

Since  $f_k$  is Cauchy and a subsequence converges, the original sequence  $f_k$  converges to  $f_{k_1} + f \in \mathcal{L}^p$  as well. We now consider the case where  $p = \infty$ . Let  $f_k$  be a Cauchy sequence in  $\mathcal{L}^\infty$ . Then for almost every  $x$ ,  $\{f_k(x)\}$  is a Cauchy sequence in  $\mathbb{R}$ . Thus we can define  $f(x)$  as the limit of  $f_k(x)$  as  $k \rightarrow \infty$ . On the set where  $f_k(x)$  does not converge, we let  $f(x)$  be zero. Then

$f \in \mathcal{L}^\infty$  since  $\{f_k\}$  is Cauchy and has an uniform bound except on a measure zero set. Also, for any  $\epsilon > 0$ , we can find  $N$  such that  $\|f_k - f_j\|_\infty < \epsilon$  for all  $k, j \geq N$ . Hence  $\|f_k - f\|_\infty < \epsilon$  for all  $k \geq N$ . Thus  $f_k \rightarrow f$  in  $\mathcal{L}^\infty$ . We conclude that  $\mathcal{L}^p$  spaces are complete. ■

### Theorem 2.25

Let  $1 \leq p < \infty$ . Let  $f_n \in \mathcal{L}^p$  be a sequence of measurable functions on a  $\sigma$ -finite measure space  $X$ . If  $f_n \rightarrow f$  in  $\mathcal{L}^p$ , then there exists a subsequence  $f_{n_k}$  such that  $f_{n_k} \rightarrow f$  a.e. on  $X$ .

*Proof.* Using the Markov inequality,

$$\begin{aligned} \mu(\{x \in X \mid |f_n(x) - f(x)| \geq \epsilon\}) &= \mu(\{x \in X \mid |f_n(x) - f(x)|^p \geq \epsilon^p\}) \\ &\leq \frac{1}{\epsilon^p} \int |f_n(x) - f(x)|^p d\mu = \frac{1}{\epsilon^p} \|f_n - f\|_p^p \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ . Thus  $f_n \xrightarrow{m} f$ . By [theorem 1.66](#), there is a subsequence  $f_{n_k}$  such that  $f_{n_k} \rightarrow f$  a.e. on  $X$ . ■

### Definition 2.26

A metric space  $(X, d)$  is **separable** if there exists a countable dense subset.

### Theorem 2.27

Let  $1 \leq p < \infty$ .  $\mathcal{L}^p(\mathbb{R})$  is separable.

*Proof.* Consider the collection of sets  $\mathcal{I} = \{(q, r) \mid q < r \in \mathbb{Q}\}$ . Then the family of functions  $F = \{\sum_{i=1}^n c_i \chi_{I_i} \mid I_i \in \mathcal{I}, c_i \in \mathbb{Q}, n \in \mathbb{N}\}$  is countable. We claim that  $F$  is dense in  $\mathcal{L}^p(\mathbb{R})$ . Indeed, since the continuous functions with compact supports are dense in  $\mathcal{L}^p(\mathbb{R})$ , it suffices to show that any such function can be approximated by functions in  $F$ . Let  $f \in \mathcal{L}^p(\mathbb{R})$  be a continuous function with compact support. By the uniform continuity, there exists  $\delta > 0$  such that for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta$ ,  $|f(x) - f(y)| < \epsilon$ .

Consider  $\mathcal{I}' = \{I \in \mathcal{I} \mid I \cap \text{supp}(f) \neq \emptyset, \mu I < \delta\}$ , an open cover of  $\text{supp}(f)$ . By the compactness of  $\text{supp}(f)$ , we can find a finite subcover  $\mathcal{I}'' = \{I_i \mid i = 1, \dots, n\}$  such that  $\text{supp}(f) \subset \cup_{i=1}^n I_i$ . By the density of  $\mathbb{Q}$  in  $\mathbb{R}$ , we can find  $c_i \in \mathbb{Q}$  such that  $|f(x) - c_i| < \epsilon$  for all  $x \in I_i$ , for  $i = 1, \dots, n$ . Let  $g = \sum_{i=1}^n c_i \chi_{I_i} \in F$ . Then  $\|f - g\|_\infty < \epsilon$ .

$$\|f - g\|_p^p = \int |f - g|^p d\mu \leq \int_{\text{supp}(f-g)} \epsilon^p d\mu = \epsilon^p \mu(\text{supp}(f - g)).$$

Since  $\epsilon$  is arbitrary, we conclude that  $F$  is dense in  $\mathcal{L}^p(\mathbb{R})$ . Thus  $\mathcal{L}^p(\mathbb{R})$  is separable. ■

### Remark

$\mathcal{L}^\infty(\Omega, \mu)$  is not separable in general. For example, let  $\Omega = [a, b]$ . Suppose that  $\{f_n\}$  is a countable dense subset of  $\mathcal{L}^\infty(\Omega)$ . Define  $\eta : [a, b] \rightarrow \mathbb{N}$  such that  $\|\chi_{[a,b]} - f_{\eta(x)}\| < \frac{1}{2}$ . Then if  $x_1 \neq x_2$ ,  $\|\chi_{[a,x_1]} - \chi_{[a,x_2]}\|_\infty = 1$ . This implies that  $f_{\eta(x_1)} \neq f_{\eta(x_2)}$  and  $\eta(x_1) \neq \eta(x_2)$ . Thus  $\eta$  is injective. But  $[a, b]$  is uncountable, a contradiction. Hence  $\mathcal{L}^\infty(\Omega)$  is not separable.

## 2.4. Dual Space

**Theorem 2.28** (Dualities of  $\ell^p$  Spaces)

Let  $1 < p < \infty$ . Then  $(\ell^p)' \cong \ell^{p'}$ , where  $p'$  is the conjugate exponent of  $p$ .

*Proof.* We need to prove that there exists an isometric isomorphism  $\psi : \ell^{p'} \rightarrow (\ell^p)'$  such that  $\psi g f = \sum_i f_i g_i$  for all  $g \in \ell^{p'}$  and  $f \in \ell^p$ . We show that  $\psi$  is well-defined, linear, bounded, bijective, and isometric.

First, we show that  $\psi$  is well-defined. For  $f \in \ell^p$  and  $g \in \ell^{p'}$ ,

$$|\psi g f| \leq \sum_i |f_i g_i| \leq \|f\|_p \|g\|_{p'} < \infty$$

by the Hölder's inequality. Thus  $\psi g \in (\ell^p)'$  is well-defined.

Next,  $\psi$  is linear since for  $g_1, g_2 \in \ell^{p'}$  and  $c \in \mathbb{R}$ ,

$$\psi(cg_1 + g_2)(f) = \sum_i f_i(cg_{1i} + g_{2i}) = c \sum_i f_i g_{1i} + \sum_i f_i g_{2i} = c\psi g_1(f) + \psi g_2(f)$$

for all  $f \in \ell^p$ . Hence  $\psi(cg_1 + g_2) = c\psi g_1 + \psi g_2$ .

Now, to show that  $\psi$  is bounded,

$$\begin{aligned} \|\psi g\| &= \sup \{ |\psi g f| \mid \|f\|_p = 1 \} = \sup \left\{ \left| \sum_i f_i g_i \right| \mid \|f\|_p = 1 \right\} \\ &\leq \sup_{\|f\|_p=1} \{ \|g\|_{p'} \} \leq \|g\|_{p'}. \end{aligned}$$

We see that  $\|\psi\| \leq 1$ . Next, let  $h \in (\ell^p)'$  and define  $g$  by  $g_i = h(e_i)$ . Then

$$\|g\|_{p'} = \left( \sum_i |g_i|^{p'} \right)^{1/p'} = \left( \sum_i |h(e_i)|^{p'} \right)^{1/p'} \leq \left( \sum_i \|h\|^{p'} \right)^{1/p'} = \|h\|.$$

Then  $g \in \ell^{p'}$ . Furthermore, for such  $g$ ,

$$\psi g(f) = \sum_i f_i g_i = \sum_i f_i h(e_i) = h\left(\sum_i f_i e_i\right) = h(f)$$

for every  $f \in \ell^p$ . Hence  $\psi$  is surjective and  $\|\psi g\| = \|h\|$ . The isometry of  $\psi$  is immediate from that

$$\|\psi g\| \leq \|g\|_{p'} \leq \|h\| = \|\psi g\|.$$

Finally,  $\psi$  is injective since otherwise there exists  $g \neq 0$  such that  $\psi g = 0$ . Then  $\|g\|_{p'} = 0$  by the isometry of  $\psi$ , which implies that  $g = 0$ , a contradiction. We conclude that  $\psi$  is an isometric isomorphism and the proof is complete. ■



**Proposition 2.29**

$1 \leq p < \infty$ ,  $1/p + 1/p' = 1$ . Let  $g \in \mathcal{L}^{p'}(X, \mu)$ . Then the mapping  $Tg : \mathcal{L}^p(X, \mu) \rightarrow \mathbb{R}$  defined by

$$Tg(f) = \int_X fg d\mu$$

is a bounded linear functional. Furthermore,  $\|Tg\|_{\mathcal{L}^p \rightarrow \mathbb{R}} = \|g\|_{p'}$ .

*Proof.* We start by checking that  $Tg$  is well-defined. For  $f \in \mathcal{L}^p$ ,

$$|Tg(f)| = \left| \int fg d\mu \right| \leq \int |fg| d\mu \leq \|f\|_p \|g\|_{p'}$$

by Hölder's inequality. Thus  $Tg(f) \in \mathbb{R}$ . Also, we obtain that  $\|Tg\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \leq \|g\|_{p'}$ . For the linearity, let  $c \in \mathbb{R}$  and  $f_1, f_2 \in \mathcal{L}^p$ .

$$Tg(cf_1 + f_2) = \int (cf_1 + f_2)g d\mu = c \int f_1 g d\mu + \int f_2 g d\mu = cTg(f_1) + Tg(f_2).$$

Lastly, to furnish the isometry, let  $g \neq 0$  and define

$$f = \text{sgn}(g) \left( \frac{|g|}{\|g\|_{p'}} \right)^{p'/p} \implies \int |f|^p d\mu = \int \left( \frac{|g|}{\|g\|_{p'}} \right)^{p'} d\mu < \infty.$$

Then  $f \in \mathcal{L}^p$  and  $\|f\|_p = 1$ . Also,

$$Tg(f) = \int \text{sgn}(g) \left( \frac{|g|}{\|g\|_{p'}} \right)^{p'/p} g d\mu = \|g\|_{p'}.$$

It follows that  $\|Tg\|_{\mathcal{L}^p \rightarrow \mathbb{R}} = \|g\|_{p'}$ . ■

**Theorem 2.30 (Riesz Representation)**

Let  $(X, \mathcal{A}, \mu)$  be a  $\sigma$ -finite measure space and  $1 \leq p < \infty$ . Then the mapping  $T : \mathcal{L}^{p'}(X, \mu) \rightarrow (\mathcal{L}^p(X, \mu))'$  defined by  $Tg \in \mathcal{L}^p(X, \mu)$ ,

$$Tg(f) = \int fg d\mu,$$

is an isometric isomorphism.

*Proof.* By **proposition 2.29**,  $Tg$  is a bounded linear functional. Besides, let  $c \in \mathbb{R}$  and  $g_1, g_2 \in \mathcal{L}^{p'}$ ,

$$T(cg_1 + g_2)(f) = \int (cg_1 + g_2)f d\mu = c \int g_1 f d\mu + \int g_2 f d\mu = cTg_1(f) + Tg_2(f) = (cTg_1 + Tg_2)(f)$$

for all  $f \in \mathcal{L}^p$ . Thus  $T$  is linear. It remains to show that  $T$  is a bijection. We first verify that  $T$  is surjective.

Consider the case where  $p > 1$  and  $\mu(X) < \infty$ . Let  $h \in (\mathcal{L}^p)'$ . Define  $\nu : \mathcal{A} \rightarrow \mathbb{R}$  by  $\nu(A) = h(\chi_A)$ . We claim that  $\nu$  is a finite measure and  $\nu \ll \mu$ . Since

$$|\nu(A)| = |h(\chi(A))| \leq \|h\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \|\chi_A\|_p = \|h\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \mu(A)^{1/p},$$

we see that  $\nu$  is finite since so is  $\mu$ . Also, if  $\mu(A) = 0$ , then  $|\nu(A)| = 0$  and hence  $\nu(A) = 0$ . Thus  $\nu \ll \mu$ . For finite additivity, let  $A_1, A_2 \in \mathcal{A}$  be disjoint.

$$\nu(A_1 \cup A_2) = h(\chi_{A_1 \cup A_2}) = h(\chi_{A_1} + \chi_{A_2}) = h(\chi_{A_1}) + h(\chi_{A_2}) = \nu(A_1) + \nu(A_2).$$

To show the  $\sigma$ -additivity, let  $A_j \in \mathcal{A}$  be countably many disjoint sets. Put  $A = \cup_j A_j$ ,  $A = B_n + C_n$  where  $B_n = \cup_{j=1}^n A_j$  and  $C_n = \cup_{j=n+1}^\infty A_j$ . Then since  $B_n \cap C_n = \emptyset$ ,

$$\nu(A) = \nu(B_n + C_n) = \nu(B_n) + \nu(C_n) = \sum_{j=1}^n \nu(A_j) + \nu(C_n)$$

for all  $n$ . Since  $\mu(X) < \infty$ ,  $\sum_j \mu(A_j) < \infty$  and  $\mu(C_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Thus

$$|\nu(C_n)| = |h(C_n)| \leq \|h\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \mu(C_n)^{1/p} \rightarrow 0.$$

We conclude that  $\nu(A) = \sum_j \nu(A_j)$  and  $\nu$  is a measure.

Next, since  $\nu \ll \mu$ , by the Radon-Nikodym theorem, there exists a unique  $g \in \mathcal{L}^1(X, \mu)$  such that

$$h(\chi_A) = \nu(A) = \int_A g d\mu = \int_X \chi_A g d\mu = Tg(\chi_A).$$

for arbitrary  $A \in \mathcal{A}$ . Extend by linearity to  $p$ -integrable simple functions, say  $s = \sum_{i=1}^n c_i \chi_{A_i}$ .

$$h(s) = \sum_{i=1}^n c_i h(\chi_{A_i}) = \sum_{i=1}^n c_i \int_X \chi_{A_i} g d\mu = \int_X \sum_{i=1}^n c_i \chi_{A_i} g d\mu = \int_X s g d\mu = Tg(s).$$

For a general  $f \in \mathcal{L}^p$ , by separating  $f = f^+ - f^-$  if necessary, we may assume that  $f \geq 0$ . By [lemma 1.20](#), there exists a sequence of simple functions  $s_n \nearrow f$ . Then by Lebesgue's monotone convergence theorem,  $\|f - s_n\|_p \rightarrow 0$ . Since  $h$  is a bounded linear functional, it is continuous, and hence  $h(s_n) \rightarrow h(f)$  as  $n \rightarrow \infty$ . We obtain that

$$h(f) = \lim_{n \rightarrow \infty} h(s_n) = \lim_{n \rightarrow \infty} \int_X s_n g d\mu = \int_X f g d\mu = Tg(f)$$

for all  $f \in \mathcal{L}^p$ . Thus  $Tg = h$ . It remains to check that  $g \in \mathcal{L}^{p'}$ . Let

$$f_n = \begin{cases} |g|^{p'-1} \operatorname{sgn}(g) & \text{if } |g(x)|^{p'-1} \leq n, \\ n \operatorname{sgn}(g) & \text{otherwise.} \end{cases}$$

Then  $f_n \in \mathcal{L}^p$  and  $f_n g \nearrow |g|^{p'}$ .

$$|Tg(f_n)| = \left| \int f_n g d\mu \right| \leq \|Tg\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \|f_n\|_p.$$

Also,  $f_n g = |f_n| |g| \geq |f_n| |f_n|^{1/(p'-1)} = |f_n|^p$  and

$$\|f_n\|_p^p = \int |f_n|^p d\mu \leq \int f_n g d\mu \leq \|Tg\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \|f_n\|_p.$$

As a result,

$$\|g\|_{p'}^{p'} = \int |g|^{p'} d\mu = \lim_{n \rightarrow \infty} \int f_n g d\mu \leq \|Tg\|_{\mathcal{L}^p \rightarrow \mathbb{R}} \|f_n\|_p < \infty.$$

Hence  $g \in \mathcal{L}^{p'}$  and  $T$  is indeed surjective. Furthermore, such  $g$  is unique by the uniqueness of the Radon-Nikodym derivative. We also conclude that  $T$  is injective.

For the case where  $p = 1$  and  $\mu(X) < \infty$ ,  $p' = \infty$ . We consider the same mapping  $T$  with  $Tg(f) = \int f g d\mu$ . We claim that  $g \in \mathcal{L}^\infty$ . Suppose  $g \notin \mathcal{L}^\infty$ . Then for every  $K$ , the set  $A_K = \{x \in X \mid |g(x)| > K\}$  has positive measure. Define  $f_K = \text{sgn}(g) \chi_{A_K} / \mu(A_K)$ . Note that  $\|f_K\|_1 = 1$ . If  $g \geq 0$ , then

$$|Tg(f_K)| = \int f_K g d\mu > K$$

for all  $K$ . But  $Tg$  is a bounded linear functional, which is a contradiction. Thus  $g \in \mathcal{L}^\infty$ .

Finally, we prove the case where  $X$  is  $\sigma$ -finite. Write  $X = \cup_n X_n$  where  $\mu(X_n) < \infty$  and  $X_n \subset X_{n+1}$ . For every  $f \in \mathcal{L}^p(X_k, \mu)$ , consider  $\hat{f} \in \mathcal{L}^p(X, \mu)$  defined by  $\hat{f} = f$  on  $X_k$  and  $\hat{f} = 0$  on  $X - X_k$ . Then  $\|f\|_{\mathcal{L}^p(X_k)} = \|\hat{f}\|_{\mathcal{L}^p(X)}$ . Let  $h \in (\mathcal{L}^p(X))'$  and consider  $h_k \in (\mathcal{L}^p(X_k))'$  by  $h_k(f) = h(\hat{f})$ . Then  $\|h_k\| \leq \|h\|$ . By the previous result, we can find a unique  $g_k \in \mathcal{L}^{p'}(X_k, \mu)$  such that

$$h_k(f) = \int f g_k d\mu, \|g_k\|_{\mathcal{L}^{p'}(X_k)} \leq \|h_k\| \leq \|h\|.$$

Since  $X_n \subset X_{n+1}$ , for  $f \in \mathcal{L}^p(X_k)$ , we have  $h_k(f) = h(\hat{f}) = h_{k+1}(f)$  and  $g_k = g_{k+1}$   $\mu$ -a.e. in  $X_k$ . Define  $g = g_k$  on  $X_k$  with  $\|g\|_{\mathcal{L}^{p'}(X)} \leq \|h\|$ . Let  $f \in \mathcal{L}^p(X, \mu)$ . Hölder's inequality implies that  $f g \in \mathcal{L}^1(X, \mu)$  and

$$h(f \chi_{X_k}) = h_k(f) = \int f \chi_{X_k} g_k d\mu$$

Since  $f \chi_{X_k} \leq |f|$ ,  $f \chi_{X_k} \rightarrow f \in \mathcal{L}^p(X, \mu)$  by Lebesgue's dominated convergence theorem. Also,

$$h_k(f) = \int f \chi_{X_k} g_k d\mu \rightarrow \int f g d\mu = Tg(f)$$

by Lebesgue's dominated convergence theorem. Thus  $T$  is indeed the desired isometric isomorphism. ■

### Remark

$(\mathcal{L}^\infty)' \not\cong \mathcal{L}^1$ . Consider  $C^\infty([-1, 1])$ , a subspace of  $\mathcal{L}^\infty$ . Define a linear functional  $\delta : C^\infty([-1, 1]) \rightarrow \mathbb{R}$  by  $\delta(f) = f(0)$ . Clearly  $\delta \in (\mathcal{L}^\infty)'$ . Now suppose there exists  $g \in \mathcal{L}^1$  such that  $\delta(f) =$

$\int_{-1}^1 fg dx$ . Let  $f = \chi_A$  where  $A$  is measurable. Then  $f \in \mathcal{L}^\infty$  and by definition,

$$0 = f(0) = \delta(f) = \int_{-1}^1 fg dx = \int_A g dx.$$

Thus  $g = 0$  a.e. and  $\delta = 0$ , a contradiction.

### Definition 2.31

$M(X)$  is a space consisting of all finite signed measures. For  $\nu \in M(X)$ , the total variation norm of  $\nu$  is defined by  $\|\nu\| = \nu^+(X) + \nu^-(X)$ , where  $\nu^+$  and  $\nu^-$  are the Hahn-Jordan decompositions of  $\nu$ .

### Proposition 2.32

$M(X)$  with the total variation norm forms a Banach space.

*Proof.* Clearly,  $M(X)$  forms a vector space. We check that  $\|\cdot\|$  is indeed a norm. For  $\nu \in M(X)$ , clearly  $\|\nu\| \geq 0$ . If  $\|\nu\| = 0$ , then  $\nu^+(X) = \nu^-(X) = 0$ ,  $\nu^+(A)$  and  $\nu^-(A)$  are zero for all  $A \in \mathcal{A}$ , and hence  $\nu = 0$ . Conversely, if  $\nu = 0$ , then so are  $\nu^+$  and  $\nu^-$  and hence  $\|\nu\| = 0$ . For  $c \in \mathbb{R}$ ,

$$\|c\nu\| = |c| \nu^+(X) + |c| \nu^-(X) = |c| (\nu^+(X) + \nu^-(X)) = |c| \|\nu\|.$$

Lastly, let  $\nu, \mu \in M(X)$ . Notice that  $(\nu + \mu)^+ \leq \nu^+ + \mu^+$  and  $(\nu + \mu)^- \leq \nu^- + \mu^-$ . Thus

$$\|\nu + \mu\| = (\nu + \mu)^+(X) + (\nu + \mu)^-(X) \leq \nu^+(X) + \mu^+(X) + \nu^-(X) + \mu^-(X) = \|\nu\| + \|\mu\|,$$

proving that  $\|\cdot\|$  is indeed a norm.

For the completeness, let  $\nu_n$  be a Cauchy sequence in  $M(X)$ . We define a measure  $\nu$  by  $\nu(A) = \lim_{n \rightarrow \infty} \nu_n(A)$  for all  $A \in \mathcal{A}$ . We claim that the limit exists and  $\nu$  is indeed a finite signed measure. Since the sequence is Cauchy, for every  $\epsilon > 0$ , there exists  $N$  such that

$$(\nu_m - \nu_n)^+(X) + (\nu_m - \nu_n)^-(X) = \|\nu_m - \nu_n\| \leq \epsilon$$

for all  $m, n \geq N$ . Since both  $(\nu_m - \nu_n)^+$  and  $(\nu_m - \nu_n)^-$  are positive measures, we have

$$(\nu_m - \nu_n)^+(A) \leq (\nu_m - \nu_n)^+(X) \leq \epsilon, \quad \text{and} \quad (\nu_m - \nu_n)^-(A) \leq (\nu_m - \nu_n)^-(X) \leq \epsilon$$

for every  $A \in \mathcal{A}$ . Thus

$$|\nu_m(A) - \nu_n(A)| = |(\nu_m - \nu_n)^+(A) - (\nu_m - \nu_n)^-(A)| \leq \epsilon.$$

It follows that for any fixed  $A \in \mathcal{A}$ ,  $\nu_n(A)$  is a Cauchy sequence in  $\mathbb{R}$  and hence the limit exists. Also, taking  $A = X$ , we see that  $\nu(X)$  is finite. To show that  $\nu$  is a measure, first note that  $\nu(\emptyset) = 0$ . For finite additivity, let  $A_1, A_2 \in \mathcal{A}$  be disjoint. Then

$$\nu(A_1 \cup A_2) = \lim_{n \rightarrow \infty} \nu_n(A_1 \cup A_2) = \lim_{n \rightarrow \infty} \nu_n(A_1) + \nu_n(A_2) = \nu(A_1) + \nu(A_2).$$

For the  $\sigma$ -additivity, let  $A_n \in \mathcal{A}$  be countably many disjoint sets. Put  $A = \cup_n A_n$ ,  $A = B_n \cup C_n$  where  $B_n = \cup_{j=1}^n A_j$  and  $C_n = \cup_{j=n+1}^\infty A_j$ . Since  $\nu(X) < \infty$ ,  $\sum_j \nu(A_j) < \infty$  and hence  $\nu(C_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Thus

$$\nu(A) = \nu(B_n) + \nu(C_n) = \sum_{j=1}^n \nu(A_j) + \nu(C_n)$$

for every  $n$  and by letting  $n \rightarrow \infty$ , we obtain  $\nu(A) = \sum_j \nu(A_j)$ . Finally, fix  $n$  and let  $m \rightarrow \infty$ ,

$$\|\nu - \nu_n\| = \lim_{m \rightarrow \infty} \|\nu_m - \nu_n\| = \lim_{m \rightarrow \infty} |\nu_m(X) - \nu_n(X)| = |\nu(X) - \nu_n(X)| \leq \epsilon$$

for all  $n \geq N$ . Thus  $\nu_n \rightarrow \nu$  in norm and  $M(X)$  is complete. ■

### Definition 2.33

Let  $f : [a, b] \rightarrow \mathbb{R}$ . The **variation** of  $f$  is defined by

$$V_{\mathcal{P}}(f) = \sum_{i=0}^{n-1} |f(t_{i+1}) - f(t_i)|,$$

where  $\mathcal{P} = \{a = t_0 < t_1 < \dots < t_n = b\}$  is a partition of  $[a, b]$ . The **total variation** of  $f$  on  $[a, b]$  is defined by

$$V(f) = \sup_{\mathcal{P}} V_{\mathcal{P}}(f).$$

### Definition 2.34

The **bounded variation space**  $BV([a, b])$  consists of all functions  $f : [a, b] \rightarrow \mathbb{R}$  such that  $V(f) < \infty$ . For  $f \in BV([a, b])$ , the **total variation norm** is defined by  $\|f\|_{TV} = |f(a)| + V(f)$ .

### Proposition 2.35

$BV([a, b])$  with the total variation norm forms a Banach space.

*Proof.* It clearly forms a vector space. We check that  $\|\cdot\|_{TV}$  is indeed a norm. First, clearly  $\|f\|_{TV} \geq 0$ . If  $\|f\|_{TV} = 0$ , then  $f(a) = 0$  and  $f(t) = f(t')$  for all  $t, t' \in [a, b]$ . Hence  $f = 0$ ; if  $f = 0$ , then  $V(f) = 0$  and  $f(a) = 0$  and  $\|f\|_{TV} = 0$ . Next, for  $c \in \mathbb{R}$ ,

$$\|cf\|_{TV} = |cf(a)| + \sum_{i=0}^{n-1} |cf(t_{i+1}) - cf(t_i)| = |c| \left( |f(a)| + \sum_{i=0}^{n-1} |f(t_{i+1}) - f(t_i)| \right) = |c| \|f\|_{TV}.$$

Lastly, let  $f, g \in BV([a, b])$ . Then

$$\begin{aligned} \|f + g\|_{TV} &= \sup_{\mathcal{P}} |(f + g)(a)| + \sum_{i=0}^{n-1} |(f + g)(t_{i+1}) - (f + g)(t_i)| \\ &\leq \sup_{\mathcal{P}} |f(a)| + |g(a)| + \sum_{i=0}^{n-1} |f(t_{i+1}) - f(t_i)| + \sum_{i=0}^{n-1} |g(t_{i+1}) - g(t_i)| \\ &\leq \sup_{\mathcal{P}} |f(a)| + \sum_{i=0}^{n-1} |f(t_{i+1}) - f(t_i)| + \sup_{\mathcal{P}} |g(a)| + \sum_{i=0}^{n-1} |g(t_{i+1}) - g(t_i)| = \|f\|_{TV} + \|g\|_{TV}. \end{aligned}$$

Thus  $\|\cdot\|_{TV}$  is indeed a norm.

For the completeness, let  $f_n$  be a Cauchy sequence in  $BV([a, b])$ . For  $\epsilon > 0$ , there exists  $N$  such that  $\|f_m - f_n\|_{TV} < \epsilon$  for all  $m, n \geq N$ . Given any  $x \in [a, b]$ , consider the partition  $\mathcal{P} = \{a < x < b\}$ .

$$\begin{aligned} |f_m(x) - f_n(x)| &= |f_m(x) - f_m(a) + f_m(a) - f_n(a) + f_n(a) - f_n(x)| \\ &\leq |((f_m(x) - f_n(x))) - (f_m(a) - f_n(a))| + |f_m(a) - f_n(a)| \\ &\leq V(f_m - f_n) + |f_m(a) - f_n(a)| = \epsilon. \end{aligned}$$

Thus  $\{f_n(x)\}$  is a Cauchy sequence in  $\mathbb{R}$  and hence converges pointwisely to, say  $f(x)$ . Furthermore, observe that the choice of  $N$  does not depend on  $x$ , and thus the convergence is uniform. We claim that  $f \in BV([a, b])$ . Indeed, for any partition  $\mathcal{P} = \{a = t_0 < \dots < t_n = b\}$ ,

$$\sum_{i=0}^{n-1} |f(t_{i+1}) - f(t_i)| \leq \sum_{i=0}^{n-1} |f(t_{i+1}) - f_N(t_{i+1})| + \sum_{i=0}^{n-1} |f(t_i) - f_N(t_i)| + V(f_N).$$

Since the convergence is uniform, we can choose  $N$  such that  $|f(t) - f_N(t)| \leq \epsilon/(2n)$ . Thus

$$\sum_{i=0}^{n-1} |f(t_{i+1}) - f(t_i)| \leq \epsilon + V(f_N).$$

Since  $f_N$  is of bounded variation, we see that  $f \in BV([a, b])$  as well. Lastly, to show that  $\|f - f_n\|_{TV} \rightarrow 0$ , first note that by definition we have  $|f_n(a) - f(a)| \rightarrow 0$ . It remains to show that  $V(f_n - f) \rightarrow 0$ . For any  $\epsilon > 0$ , there exists  $N$  such that  $V_{\mathcal{P}}(f_m - f_n) < \epsilon$  for all  $m, n \geq N$  and some partition  $\mathcal{P}$ . Taking  $m \rightarrow \infty$ , we obtain  $V_{\mathcal{P}}(f - f_n) < \epsilon$  for all  $n \geq N$ . Since the partition is arbitrary, we have  $V(f - f_n) < \epsilon$  for all  $n \geq N$ . Thus  $f_n \rightarrow f$  in  $BV([a, b])$  and  $BV([a, b])$  is complete. ■

### Theorem 2.36

$M([a, b])$  is isometrically isomorphic to  $BV([a, b])$ .

*Proof.* We define the mapping  $\phi : M([a, b]) \rightarrow BV([a, b])$  by

$$\rho(t) = \phi v(t) = v([a, t]).$$

First, we show that  $\rho \in BV([a, b])$ . For any partition  $\mathcal{P} = \{a = t_0 < \dots < t_n = b\}$ ,

$$\begin{aligned} \sum_{i=0}^{n-1} |\rho(t_{i+1}) - \rho(t_i)| + |\rho(a)| &= \sum_{i=0}^{n-1} |v([a, t_{i+1}]) - v([a, t_i])| + |v(\{a\})| \\ &= \sum_{i=0}^{n-1} |v((t_i, t_{i+1}])| + |v(\{a\})| \\ &= \sum_{i=0}^{n-1} |v|((t_i, t_{i+1}]) + |v|(\{a\}) = |v|([a, b]) = \|v\|. \end{aligned}$$

Since  $\nu$  is a finite signed measure,  $\rho \in BV([a, b])$ . Furthermore, taking supremum over all partitions, we obtain that  $\|\rho\|_{TV} = \|\nu\|$ . It remains to show that  $\phi$  is an isomorphism. Suppose  $\nu, \mu \in M([a, b])$  and  $\phi\nu = \phi\mu$ . Then  $\nu([a, t]) = \mu([a, t])$  for all  $t \in [a, b]$ . Since  $[a, t]$  generates the Borel  $\sigma$ -algebra on  $[a, b]$ , we have  $\nu = \mu$ . Thus  $\phi$  is injective. For surjectivity, let  $\rho \in BV([a, b])$ . Consider the signed measure  $\nu$  defined by  $\nu([a, t]) = \rho(t)$  and  $\nu(\emptyset) = 0$ . Then  $\nu$  is a finite signed measure and  $\phi\nu = \rho$ . The proof is complete. ■

### Lemma 2.37

Let  $X$  be a normed vector space and  $M \subset X$  be a proper subspace. Suppose  $S : M \rightarrow \mathbb{R}$  is a bounded linear functional. Then for every  $x \in X \setminus M$ , there exists a linear  $U : M' \rightarrow \mathbb{R}$  such that  $\|U\|_{M' \rightarrow \mathbb{R}} = \|S\|_{M \rightarrow \mathbb{R}}$ , where  $M' = M + \mathbb{R}x$ .

*Proof.* Clearly  $M'$  is a subspace; furthermore,  $M' = M \oplus \mathbb{R}x$  since if  $v = w + cx = w' + c'x$  for some  $w, w' \in M$  and  $c, c' \in \mathbb{R}$ , then  $(c - c')x = w - w' \in M$ . Since  $x \notin M$ , this implies that  $c = c'$ ,  $w = w'$  and hence the representation is unique.

Now we can define  $U$  on  $M'$  by  $U(w + cx) = Sw + c\lambda$  for any  $w + cx \in M'$  and some  $\lambda \in \mathbb{R}$  to be determined. To make  $U$  have the same norm as  $S$ , we need to find  $\lambda$  such that  $|Sw + c\lambda| \leq \|S\| \|w + cx\|$  holds for all  $w \in M$  and  $c \in \mathbb{R}$ . Clearly if  $c = 0$ , the inequality is already satisfied. For  $c \neq 0$ , by dividing both sides by  $|c|$ , we see that the condition is equivalent to  $|Sw + \lambda| \leq \|S\| \|w + x\|$  for all  $w \in M$ . Now for any  $w, v \in M$ ,

$$Sw - Sv = S(w - v) \leq |S(w - v)| \leq \|S\| \|w - v\| = \|S\| \|w + x - (v + x)\| \leq \|S\| (\|w + x\| + \|v + x\|).$$

Thus

$$Sw - \|S\| \|w + x\| \leq Sv + \|S\| \|v + x\|.$$

Fix  $v$  and taking supremum over all  $w \in M$  on the left,

$$\sup_{w \in M} Sw - \|S\| \|w + x\| \leq Sv + \|S\| \|v + x\|.$$

Taking infimum over all  $v \in M$  on the right,

$$\sup_{w \in M} Sw - \|S\| \|w + x\| \leq \inf_{v \in M} Sv + \|S\| \|v + x\|.$$

Hence there exists  $\lambda \in \mathbb{R}$  such that

$$S(w) - \|S\| \|w + x\| \leq -\lambda \leq S(w) + \|S\| \|w + x\|$$

for all  $w \in M$ . Picking this  $\lambda$ , we see that

$$|Sw + \lambda| \leq \|S\| \|w + x\|$$

as desired. Thus  $U$  is a bounded linear functional on  $M'$  with  $\|U\|_{M' \rightarrow \mathbb{R}} = \|S\|_{M \rightarrow \mathbb{R}}$ . Also, on

$M$ ,  $U = S$  and hence  $U$  is an extension of  $S$ . ■

**Theorem 2.38** (Hahn-Banach)

Let  $X$  be a normed vector space and  $M \subset X$  be a subspace. Suppose  $S : M \rightarrow \mathbb{R}$  is a bounded linear functional on  $M$ . Then there exists a bounded linear functional  $T : X \rightarrow \mathbb{R}$  such that  $T|_M = S$  and  $\|T\|_{X \rightarrow \mathbb{R}} = \|S\|_{M \rightarrow \mathbb{R}}$ .

*Proof.* The proof relies on Zorn's lemma.<sup>1</sup> We start by constructing a partial order space. Let  $(P, \preceq)$  be a partial order space with

$$P = \{(U, Y) \mid M \subset Y \subset X, Y \text{ is a subspace of } X, U \text{ is a bounded extension of } S \text{ on } Y\}$$

and the partial order:  $(U_1, Y_1) \preceq (U_2, Y_2)$  if  $Y_1 \subset Y_2$  and  $U_2$  is a bounded extension of  $U_1$  on  $Y_2$ . Clearly the pair indeed forms a partial order space. We now check the assumptions of Zorn's lemma. Let  $C = \{(U_\alpha, Y_\alpha) \mid \alpha \in A\}$  with an arbitrary index set  $A$  be a chain in  $P$ . Put  $Y = \cup_{\alpha \in A} Y_\alpha$ . We claim that  $Y$  is a subspace of  $X$ . Indeed, for  $y_1, y_2 \in Y$  and  $c_1, c_2 \in \mathbb{R}$ , there exist  $\alpha_1, \alpha_2 \in A$  such that  $y_1 \in Y_{\alpha_1}$  and  $y_2 \in Y_{\alpha_2}$ . Since  $Y$  is a chain, one of them is a subspace of the other, say  $Y_{\alpha_1}$  is a subspace of  $Y_{\alpha_2}$ . Then  $y_1, y_2 \in Y_{\alpha_2}$  and hence  $c_1 y_1 + c_2 y_2 \in Y_{\alpha_2} \subset Y$ . Thus  $Y$  is a subspace.

Next we need to define a bounded linear functional  $U$  on  $Y$  so that  $U$  is a bounded extension of  $S$  on  $Y$ . For  $y \in Y$ , we can find an  $\alpha \in A$  such that  $y \in Y_\alpha$  and set  $U(y) = U_\alpha(y)$ . Such  $U$  is well-defined since if  $\alpha_1$  and  $\alpha_2$  are two indices satisfying  $y \in Y_{\alpha_1} \cap Y_{\alpha_2}$ , then  $U_{\alpha_1}(y) = U_{\alpha_2}(y)$  since one of them is an extension of the other. Also,  $U$  is linear since  $U_\alpha$  is linear for every  $\alpha \in A$ . Lastly,  $U$  is a bounded extension of  $U_\alpha$  on  $Y$  for any  $\alpha \in A$  because every  $U_{\alpha'}$  with  $(U_\alpha, Y_\alpha) \preceq (U_{\alpha'}, Y_{\alpha'})$  is a bounded extension of  $U_\alpha$ . We conclude that  $(U, Y) \in P$  is an upper bound of  $C$ .

By Zorn's lemma, there exists a maximal element  $(T, Z) \in P$ . We claim that  $Z = X$ . Suppose  $Z \subsetneq X$ . Then there exists  $x \in X \setminus Z$  and also a bounded extension  $T'$  of  $T$  on  $Z + \mathbb{R}x \supsetneq Z$  by [lemma 2.37](#). But then  $(T', Z + \mathbb{R}x) \in P$  and  $(T, Z) \preceq (T', Z + \mathbb{R}x)$ , contradicting the maximality of  $(T, Z)$ . Thus  $Z = X$  and  $T$  is a bounded extension of  $S$  on  $X$ . ■

**Theorem 2.39** (Riesz Representation of  $C([a, b])$ )

$C([a, b])' \cong BV([a, b]) \cong M([a, b])$  isometrically.

*Proof.* In [theorem 2.36](#), we have shown that  $M([a, b]) \cong BV([a, b])$ . We are going to show this by constructing an isometric isomorphism between  $C([a, b])'$  and  $BV([a, b])$ .

Let  $X = C([a, b])$  and  $\ell \in X'$ .  $\ell : X \rightarrow \mathbb{R}$  is a bounded linear functional. We need to find a  $\nu \in M([a, b])$  such that

$$\ell(f) = \int_{[a, b]} f d\nu$$

---

<sup>1</sup>Zorn's lemma states that if every chain in a partially ordered set has an upper bound, then the set has a maximal element. It is a direct consequence of the axiom of choice.



for  $f \in C([a, b])$ . Let  $Y = B([a, b]) = \{f : [a, b] \rightarrow \mathbb{R} \mid f \text{ is bounded}\}$ . By Hahn-Banach theorem, there exists a bounded linear extension  $L : Y \rightarrow \mathbb{R}$  of  $\ell$ . Now if  $f = \chi_{[a, t]} \in Y$ , then

$$L(f) = \int_{[a, b]} \chi_{[a, t]} d\nu = \nu([a, t]) = \rho(t).$$

We claim that  $\rho \in BV([a, b])$ . For any partition  $\mathcal{P} = \{a = t_0 < \dots < t_n = b\}$ ,

$$\begin{aligned} V_{\mathcal{P}}(\rho) &= \sum_{i=0}^{n-1} |\rho(t_{i+1}) - \rho(t_i)| = \sum_{i=0}^{n-1} |L(\chi_{[a, t_{i+1}]} - L(\chi_{[a, t_i]})| \\ &= \sum_{i=0}^{n-1} L(\chi_{(t_i, t_{i+1}]} s_i) = L\left(\sum_{i=0}^{n-1} \chi_{(t_i, t_{i+1}]} s_i\right) \leq \|L\| \left\| \sum_{i=0}^{n-1} \chi_{(t_i, t_{i+1}]} s_i \right\|_{\infty} \leq \|L\| \end{aligned}$$

by letting  $s_i = \text{sgn}(\rho(t_{i+1}) - \rho(t_i))$ . Thus  $\rho \in BV([a, b])$  and  $\|\rho\|_{TV} \leq \|L\| = \|\ell\|$ . To extend to  $f \in C([a, b])$  so that

$$\ell(f) = L(f) = \int_{[a, b]} f d\nu,$$

we first note that by our established result,  $f = \chi_{[a, t]} \in Y$  holds. By linearity so does simple functions. For  $f \in C([a, b])$ , consider

$$h_{\mathcal{P}}(t) = f(a) + \sum_{i=0}^{n-1} f(t_i) \chi_{(t_i, t_{i+1}]}(t).$$

Since  $L$  is continuous and  $h_{\mathcal{P}} \rightarrow f$  uniformly as  $\|\mathcal{P}\| \rightarrow 0$ , we have

$$L(f) = \lim_{\|\mathcal{P}\| \rightarrow 0} L(h_{\mathcal{P}}) = \int_a^b f d\rho.$$

$L$  is an extension of  $\ell$  and hence

$$\ell(f) = \int_a^b f d\rho = f(a)\rho(a) + \int_a^b f d\rho.$$

Finally, we claim that  $\|\ell\| \leq \|\rho\|_{TV} \leq \|L\| = \|\ell\|$ . Take  $f \in X$ .

$$|\ell(f)| = \left| \int_a^b f d\rho \right| \leq \|f\|_{\infty} \|\rho\|_{TV} \leq \|f\|_{\infty} \|L\| = \|\ell\| \|f\|_{\infty}.$$

Hence  $\|\ell\| \leq \|\rho\|_{TV} \leq \|L\| = \|\ell\|$ . It follows that the mapping  $\ell \mapsto \rho$  is isometric. Conversely, if  $\rho \in BV([a, b])$ , define

$$\ell_{\rho}(f) = f(a)\rho(a) + \int_a^b f d\rho.$$

We need to check that  $\ell_{\rho}$  is linear and  $\|\rho\|_{TV} \leq \|\ell\| \leq \|\rho\|_{TV}$ .  $\ell_{\rho}$  has an extension  $L_{\rho} : Y \rightarrow \mathbb{R}$ . Define  $\lambda(t) = L_{\rho}(\chi_{[a, t]})$ . Then  $\|\rho\|_{TV} = \|\lambda\| \leq \|L_{\rho}\| = \|\ell_{\rho}\|$ . ■

**Remark**

If  $\ell \in C([a, b])'$ , there exists  $\rho \in BV([a, b])$  such that

$$\ell(f) = \int_a^b f d\rho;$$

if  $\rho \in BV([a, b])$ ,

$$\ell_\rho(f) = f(a)\rho(a) + \int_a^b f d\rho$$

and  $\|\ell_\rho\| = \|\rho\|_{TV}$ .

**Definition 2.40**

Let  $X$  be a Banach space and  $J : X \rightarrow X''$ , the canonical mapping defined by  $J : x \mapsto (T \mapsto Tx)$  for  $T \in X'$ .  $X$  is called **reflexive** if  $J$  is surjective.

**Remark**

Intuitively,  $X$  is reflexive meaning that  $X \cong X''$ . Such canonical mapping is well-defined since  $\hat{x} = T \mapsto T(x)$  is indeed a bounded linear functional on  $X'$ , which we verify here. This is linear because

$$\hat{x}(cT + S) = (cT + S)(x) = cT(x) + S(x) = c\hat{x}(T) + \hat{x}(S)$$

for  $c \in \mathbb{R}$  and  $S \in X'$ . To show that  $\hat{x}$  is bounded, we have

$$|\hat{x}(T)| = |T(x)| \leq \|T\| \|x\| = \|T\|_{X'} \|x\|_X$$

**Definition 2.41**

A Banach space  $X$  is said to be **uniformly convex** if for all  $\epsilon > 0$ ,  $x, y \in X$  with  $\|x - y\| \geq \epsilon > 0$  and  $\|x\|, \|y\| \leq 1$ , we have

$$\left\| \frac{x + y}{2} \right\| \leq 1 - \delta$$

for some  $\delta > 0$  depending on  $\epsilon$ .

**Theorem 2.42** (Clarkson)

$\mathcal{L}^p(\Omega, \mu)$  is uniformly convex for  $1 < p < \infty$ .

*Proof.* Consider the function  $\alpha : \mathbb{R}^2 \rightarrow [0, \infty)$  defined by

$$\alpha(x) = \frac{|x_1|^p + |x_2|^p}{2} - \left| \frac{x_1 + x_2}{2} \right|^p$$

for  $x = (x_1, x_2) \in \mathbb{R}^2$ . Observe that  $\alpha(x) \geq 0$  for all  $x \in \mathbb{R}^2$  and  $\alpha(x) = 0$  if and only if  $x_1 = x_2$  by the strict convexity of  $|\cdot|^p$ . Now given  $\epsilon > 0$ , choose  $\eta < \epsilon^p/2$ . Consider the set  $D = \{x \in \mathbb{R}^2 \mid |x_1|^p + |x_2|^p = 1, |x_1 - x_2|^p \geq \eta\}$ . Then  $D$  is a compact set. By the compactness, we can set  $\theta = \inf_{x \in D} \alpha(x) > 0$ . We now claim that if  $x \in \mathbb{R}^2$  satisfies  $|x_1 - x_2|^p \geq \eta(|x_1|^p + |x_2|^p)$ , then

$$|x_1|^p + |x_2|^p \leq \frac{\alpha(x)}{\theta}.$$

By the assumption, we may assume that  $x \neq (0, 0)$ . Set  $t = (|x_1|^p + |x_2|^p)^{1/p}$ . Then

$$\left| \frac{x_1}{t} \right|^p + \left| \frac{x_2}{t} \right|^p = 1, \quad \text{and} \quad \left| \frac{x_1}{t} - \frac{x_2}{t} \right|^p \geq \eta.$$

Thus

$$\theta \leq \alpha\left(\frac{x}{t}\right) = \frac{\alpha(x)}{t^p} \Rightarrow |x_1|^p + |x_2|^p = t^p \leq \frac{\alpha(x)}{\theta}.$$

The claim follows.

Now let  $f, g \in \mathcal{L}^p(\Omega, \mu)$  with  $\|f\|_p, \|g\|_p \leq 1$  and  $\|f - g\|_p \geq \epsilon$ . Put

$$E = \{x \in \Omega \mid |f(x) - g(x)|^p \geq \eta(|f(x)|^p + |g(x)|^p)\}.$$

Using the claim and  $\alpha(x) \geq 0$ ,

$$\begin{aligned} \epsilon^p &\leq \int_{\Omega} |f - g|^p d\mu = \int_E |f - g|^p d\mu + \int_{E^c} |f - g|^p d\mu \\ &\leq 2^p \int_E \left| \frac{f - g}{2} \right|^p d\mu + \eta \int_{E^c} |f|^p d\mu + \eta \int_{E^c} |g|^p d\mu \\ &\leq 2^p \int_E \frac{|f|^p + |g|^p}{2} d\mu + 2\eta \\ &\leq 2^{p-1} \int_E \frac{\alpha(f, g)}{\theta} d\mu + 2\eta \\ &\leq \frac{2^{p-1}}{\theta} \int_{\Omega} \frac{|f|^p + |g|^p}{2} - \left| \frac{f + g}{2} \right|^p d\mu + 2\eta \leq \frac{2^{p-1}}{\theta} + 2\eta - \frac{2^{p-1}}{\theta} \left\| \frac{f + g}{2} \right\|_p^p. \end{aligned}$$

Hence

$$\left\| \frac{f + g}{2} \right\|_p^p \leq 1 - \theta \frac{\epsilon^p - 2\eta}{2^{p-1}} \Rightarrow \left\| \frac{f + g}{2} \right\|_p \leq 1 - \delta$$

for some  $\delta > 0$ . We conclude that  $\mathcal{L}^p(\Omega, \mu)$  is uniformly convex for  $1 < p < \infty$ . ■

### **Theorem 2.43** (Milman-Pettis)

*Every uniformly convex Banach space is reflexive.*

*Proof.* ■

### **Corollary 2.44**

$\mathcal{L}^p(\Omega, \mu)$  is reflexive for any  $1 < p < \infty$ .

*Proof.* This is a direct consequence of the Clarkson theorem and Milman-Pettis theorem. Since for every  $1 < p < \infty$ ,  $\mathcal{L}^p(\Omega, \mu)$  is uniformly convex, and every uniformly convex Banach space is reflexive, we conclude that  $\mathcal{L}^p(\Omega, \mu)$  is reflexive. ■

### **Remark**

*This corollary can also be inferred from the Riesz representation theorem twice.*

**Example**

Fix  $x \in X$  and define the functional  $L_x : X' \rightarrow \mathbb{R}$  by  $L_x(\ell) = \ell(x)$ . Then  $L_x$  is a bounded linear functional. To see this, let  $c \in \mathbb{R}$  and  $\ell_1, \ell_2 \in X'$ .

$$L_x(c\ell_1 + \ell_2) = (c\ell_1 + \ell_2)(x) = c\ell_1(x) + \ell_2(x) = cL_x(\ell_1) + L_x(\ell_2).$$

And also

$$\|L_x\| = \sup_{\|\ell\|=1} |L_x(\ell)| = \sup_{\|\ell\|=1} |\ell(x)| \leq \sup_{\|\ell\|=1} \|\ell\| \|x\| = \|x\|.$$

In fact, we have  $\|L_x\| = \|x\|$ . To see this, consider the one-dimensional subspace  $Y = \mathbb{R}x$  and the functional  $s : Y \rightarrow \mathbb{R}$  defined by  $s(\lambda x) = \lambda \|x\|$  for some  $\lambda \neq 0$ . Then by the Hahn-Banach theorem, there exists a bounded linear functional  $s' : X \rightarrow \mathbb{R}$  such that  $s'|_Y = s$  with  $\|s'\|_X = \|s\|_Y$ . Then  $s' \in X'$  and

$$\|L_x\| \geq |L_x(s')| = |s'(x)| = \|x\|.$$

Hence  $\|L_x\| = \|x\|$ .

**Proposition 2.45**

If  $X$  is a finite-dimensional Banach space, then  $X$  is reflexive.

*Proof.* Let  $X$  be a finite-dimensional Banach space. Then  $X$  is isomorphic to  $\mathbb{R}^n$  for some  $n \in \mathbb{N}$ . Since the dual of  $\mathbb{R}^n$  is also  $\mathbb{R}^n$ , we have  $X' \cong X$ . Thus  $X'' \cong X' \cong X$ . Hence  $X$  is reflexive. ■

**Example**

Consider the space  $C[-1, 1]$  with  $\|f\| = \sup_{x \in [-1, 1]} |f(x)|$ . Then  $C[-1, 1]$  is a Banach space, but not reflexive. Suppose  $C[-1, 1]$  is reflexive. Write  $(C[-1, 1])'' \cong C[-1, 1]$ . Then the mapping  $L \mapsto L_x$  is an isomorphism, where  $L : (C[-1, 1])' \rightarrow \mathbb{R}$ ,  $L \in (C[-1, 1])''$ . Then, for all  $\ell \in C[-1, 1]'$ , there exists an  $x \in C[-1, 1]$  and  $L_x \in C[-1, 1]''$  such that

$$\begin{cases} \|L_x\| = 1, & \text{and } \|\ell\| = L_x(\ell), \\ \|x\| = 1, & \text{and } \|\ell\| = \ell(x). \end{cases}$$

Now consider the functional

$$\ell(g) = \int_{-1}^0 g(x) dx - \int_0^1 g(x) dx.$$

Clearly  $\ell$  is a bounded linear functional on  $C[-1, 1]$ .

$$|\ell(g)| \leq \int_{-1}^0 |g(x)| dx + \int_0^1 |g(x)| dx \leq 2 \sup_{x \in [-1, 1]} |g(x)| = 2 \|g\|.$$

Thus  $\|\ell\| \leq 2$ . In fact, we have  $\|\ell\| = 2$  by considering the continuous functions

$$g_\epsilon(x) = \begin{cases} 1, & x \in [-1, -\epsilon], \\ -\frac{x}{\epsilon}, & x \in [-\epsilon, \epsilon], \\ -1, & x \in [\epsilon, 1]. \end{cases}$$

Then  $\|g_\epsilon\| = 1$  and  $\ell(g_\epsilon) = 2 - \epsilon \rightarrow 2$  as  $\epsilon \rightarrow 0$ .

However, if  $f \in C[-1, 1]$  has  $\|f\| = 1$ , then there exists an interval  $I \subset [-1, 1]$  with  $\mu(I) > 0$  such that  $\sup_I |f| < 1 - \delta$ , where  $\delta > 0$ . Then

$$\begin{aligned} |\ell(f)| &= \left| \int_{[-1,0]-I} f(x)dx + \int_{[-1,0] \cap I} f(x)dx - \int_{[0,1] \cap I} f(x)dx - \int_{[0,1]-I} f(x)dx \right| \\ &\leq \mu([-1, 0] - I) + \mu([-1, 0] \cap I)(1 - \delta) + \mu([0, 1] \cap I)(1 - \delta) + \mu([0, 1] - I) \\ &= 2 - \delta\mu(I). \end{aligned}$$

This contradicts the fact that there is an  $x \in C[-1, 1]$  such that  $\|x\| = 1$  and  $\|\ell\| = \ell(x)$ . Thus  $C[-1, 1]$  is not reflexive.

### Theorem 2.46

Let  $X$  be a Banach space. If  $X'$  is separable, then  $X$  is also separable.

*Proof.* By the assumption, let  $\{\ell_n\} \subset X'$  be a countable dense subset. By definition, since  $\|\ell_n\| = \sup_{\|z\|=1} |\ell_n(z)|$ , for each  $n \in \mathbb{N}$ , we can find a  $z_n \in X$  such that  $\|z_n\| = 1$  and  $|\ell_n(z_n)| \geq \frac{1}{2} \|\ell_n\|$ .

Now we claim that  $Y = \text{span}\{z_n\}$  is dense in  $X$ . Suppose not. Then there is an  $x \in X \setminus Y$ . Consider the space  $W = \{cx + y \mid c \in \mathbb{R}, y \in Y\}$ . Define a linear functional  $\ell(v) = c\ell(x) \neq 0$  for  $v = cx + y$  with  $c \neq 0$  on  $W$ . By the Hahn-Banach theorem, we can extend  $\ell$  to  $X$ . For such  $\ell$  on  $X$ , we have  $\ell(z_n) = 0$  for all  $n$  since  $z_n \in Y$ . Assume without loss of generality that  $\|\ell\| = 1$  and  $\|\ell_n - \ell\| \leq \epsilon$  by the density of  $\{\ell_n\}$  in  $X'$ . Hence

$$\|\ell_n\| \geq \|\ell\| - \|\ell_n - \ell\| \geq 1 - \epsilon.$$

On the other hand,

$$\|\ell_n\| \leq 2|\ell_n(z_n)| = 2|\ell_n(z_n) - \ell(z_n)| \leq 2\|\ell_n - \ell\| \|z_n\| \leq 2\epsilon.$$

This implies that  $1 \leq 3\epsilon$ . Picking  $\epsilon < 1/3$  leads to a contradiction. Hence  $Y$  is dense in  $X$ .

Finally, write  $X = \overline{Y} = \overline{\{\sum_{k=1}^M \alpha_k z_k \mid M \geq 1, \alpha_k \in \mathbb{R}\}} = \overline{\{\sum_{k=1}^M \alpha_k z_k \mid M \geq 1, \alpha_k \in \mathbb{Q}\}}$ . Thus  $X$  is separable. ■

### Theorem 2.47

Let  $X$  be a reflexive Banach space. If  $Y \subset X$  is a closed subspace, then  $Y$  is reflexive.

*Proof.* Fix a bounded lineal functional  $L : Y' \rightarrow \mathbb{R}$ . We want to show that there exists a unique  $z \in Y$  such that  $L(\ell) = L_z(\ell) = \ell(z)$  for all  $\ell \in Y'$ . Suppose  $\ell : X \rightarrow \mathbb{R}$  is a bounded linear functional on  $X$ . Consider its restriction on  $Y$ ,  $\ell|_Y$ . Note that  $\|\ell|_Y\| \leq \|\ell\|$ .

Now for  $L$ , we can extend by Hahn-Banach theorem to  $L_0 : X' \rightarrow \mathbb{R}$ . For  $m \in X'$ ,  $m : X \rightarrow \mathbb{R}$ , consider its restriction on  $Y$ ,  $m|_Y$ . Then  $L_0(m) = L(m|_Y)$ . We check that  $L_0$  is linear and bounded. For  $c \in \mathbb{R}$  and  $m, \ell \in X'$ ,

$$L_0(cm + \ell) = L((cm + \ell)|_Y) = L(cm|_Y + \ell|_Y) = cL(m|_Y) + L(\ell|_Y) = cL_0(m) + L_0(\ell).$$

And also

$$|L_0(m)| = |L(m|_Y)| \leq \|L\| \|m|_Y\| \leq \|L\| \|m\|. \Rightarrow \|L_0\| \leq \|L\|.$$

Thus  $L_0$  is a bounded linear functional on  $X'$ . We now use the reflexivity of  $X$ . Since  $X'' \cong X$ , there exists a  $z \in X$  such that  $L_0(m) = L_z(m)$  for all  $m \in X'$ .

We claim that  $z \in Y$ . Suppose not. Then there exists a bounded linear functional  $m : \{cz + y \mid c \in \mathbb{R}, y \in Y\} \rightarrow \mathbb{R}$  such that  $m(z) \neq 0$  and  $m(y) = 0$  for all  $y \in Y$ . Extend  $m$  to  $m_0 : X \rightarrow \mathbb{R}$  by Hahn-Banach theorem. Then

$$L_0(m_0) = L(m_0|_Y) = L(0) = 0 \neq m_0(z) = L_0(m_0),$$

which is absurd. Hence  $z \in Y$  and we see that  $L(m) = m(z)$  for all  $m \in Y'$ . Take  $m \in Y'$  and its extension  $m_0 \in X'$ . If  $m_0, m'_0 \in X'$  are two extensions of  $m$ , then  $L_0(m_0) = L_0(m'_0)$  and hence  $L(m) = L_0(m_0) = m_0(z) = m(z)$ . Thus the extension is unique. We conclude that  $Y$  is a reflexive Banach space. ■

## 2.5. Weak Convergence

### Definition 2.48

Let  $(X, \|\cdot\|)$  be a normed space. A sequence  $\{x_n\}$  in  $X$  is said to **converge weakly** to  $x \in X$ , denoted by  $x_n \xrightarrow{w} x$ , if for every  $L \in X'$ ,  $L(x_n) \rightarrow L(x)$  as  $n \rightarrow \infty$ .

### Remark

Strong convergence implies weak convergence. If  $x_n \rightarrow x$ ,

$$|L(x_n) - L(x)| = |L(x_n - x)| \leq \|L\| \|x_n - x\| \rightarrow 0$$

as  $n \rightarrow \infty$ . Thus  $x_n \xrightarrow{w} x$ . However, the converse is not true in general.

### Example

Consider  $\ell^2$ . Note that  $(\ell^2)' \cong \ell^2$ . For all  $L \in (\ell^2)'$ , there exists  $y \in \ell^2$  such that  $L(x) = \sum_{n=1}^{\infty} x_n y_n$ . Let  $x_n = e^n$  be the sequence with 1 at the  $n$ -th position and 0 elsewhere. Then

$x_n \xrightarrow{w} 0$  since for every  $L \in (\ell^2)'$ ,

$$L(x_n) = \sum_i e_i^n y_i = y_n \rightarrow 0$$

for  $y \in \ell^2$ . However,  $\|x_n\|_{\ell^2} = 1$  for every  $n$  and thus  $x_n \not\rightarrow 0$ .

### Example

Consider  $X = C([0, 1])$  with the supremum norm. Let

$$x_n(t) = \begin{cases} nt & \text{if } 0 \leq t \leq 1/n, \\ 2 - nt & \text{if } 1/n \leq t \leq 2/n, \\ 0 & \text{if } 2/n \leq t \leq 1. \end{cases}$$

Then  $\|x_n\|_\infty = 1$  and thus  $x_n \not\rightarrow 0$ . Instead, we have  $x_n \xrightarrow{w} 0$ . Assume not, then we can find  $T \in X'$  and a subsequence  $\{x_{n_k}\}$  such that  $|T(x_{n_k})| \geq \delta > 0$ . For simplicity, we consider the case  $T(x_{n_k}) \geq \delta$ , but the other case is similar. Since  $T \in X'$ ,  $|T(x_{n_k})| \leq \|T\|_{X \rightarrow \mathbb{R}} \|x_{n_k}\|_\infty$ . Let  $y_K = \sum_{k=1}^K x_{n_k}$ . Then  $T(y_K) = \sum_{k=1}^K T(x_{n_k}) \geq K\delta$  and  $T(y_K) \leq \|T\|_{X \rightarrow \mathbb{R}} \|y_K\|_\infty$ . This implies that  $y_K$  cannot be bounded. Now consider  $x_{n_k}$  with  $n_{k+1} \geq 2n_k$ . For  $t \in [0, 1/n_K]$ ,  $x_{n_k}(t) = n_k t$ .

$$y_K(t) = \sum_{k=1}^K n_k t \leq \sum_{k=1}^K n_k / n_K \leq 1 + \sum_{k=1}^K 2^{K-k} \leq 1 + \sum_k 2^{-k} = 2.$$

For  $t \in [1/n_K, 1/n_{K-1}]$ ,

$$y_K(t) = \sum_{k=1}^K x_{n_k}(t) \leq 1 + \sum_{k=1}^{K-1} n_k t \leq 1 + \frac{1}{n_{K-1}} \sum_{k=1}^{K-1} n_k \leq 1 + 1 + \sum_k 2^{-k} = 3.$$

On  $[1/n_K, 1/n_{K-1}]$ , we have  $\|y_K\| \leq 3$ . Thus  $\delta K \leq \|T\|_{X \rightarrow \mathbb{R}} \|y_K\|_\infty \leq 3 \|T\|_{X \rightarrow \mathbb{R}}$ , which is impossible for sufficiently large  $K$ . Hence  $x_n \xrightarrow{w} 0$ .

### Proposition 2.49

$(X, \|\cdot\|_X)$  is a normed space and  $x_n \in X$ . If  $\|x_n\|_X \leq C$  for all  $n \in \mathbb{N}$  and  $L(x_n) \rightarrow L(x)$  for all  $L \in A \subset X'$ , where  $A$  is dense in  $X'$ , then  $x_n \xrightarrow{w} x$  in  $X$ .

*Proof.* Let  $\epsilon > 0$  be given.  $A$  is dense in  $X'$ . For  $T \in X'$ , there is an  $L \in A$  such that  $\|T - L\|_{X' \rightarrow \mathbb{R}} \leq \epsilon$ . Also, there exists  $N$  such that  $|L(x_n) - L(x)| \leq \epsilon$  for all  $n \geq N$ . Then

$$\begin{aligned} |T(x_n) - T(x)| &\leq |T(x_n) - L(x_n)| + |L(x_n) - L(x)| + |L(x) - T(x)| \\ &\leq \|T - L\|_{X' \rightarrow \mathbb{R}} (\|x_n\|_X + \|x\|_X) + |L(x_n) - L(x)| \leq 2C\epsilon + \epsilon \end{aligned}$$

for all  $n \geq N$ . Since  $\epsilon$  is arbitrary,  $x_n \xrightarrow{w} x$ . ■

### Definition 2.50

A space  $X$  is called a **Baire space** if for any sequence of open dense subsets  $\{E_n\}$ ,  $\cap_n E_n$  is dense in  $X$ .

**Theorem 2.51** (Baire Category Theorem)

A complete metric space is a Baire space.

*Proof.* Let  $X$  be a complete metric space and  $\{E_n\}$  be a sequence of open dense subsets in  $X$ . Put  $E = \cap_n E_n$ . We want to show that any nonempty open set  $G \subset X$  intersects  $E$ .

$E_1$  is dense in  $X$  so  $G \cap E_1$  is nonempty. Then there exists  $x_1 \in E_1 \cap G$ . Note that  $E_1 \cap G$  is open; there exists  $1 > \delta_1 > 0$  such that  $B_{\delta_1}(x_1) \subset E_1 \cap G$ . By shrinking  $\delta_1$ , we can have  $\overline{B_{\delta_1}(x_1)} \subset E_1 \cap G$ . Now since  $E_2$  is dense in  $X$ , there exists  $x_2 \in E_2 \cap B_{\delta_1}(x_1)$  and also a  $1/2 > \delta_2 > 0$  such that  $\overline{B_{\delta_2}(x_2)} \subset E_2 \cap B_{\delta_1}(x_1)$ . Continue this process, we obtain a sequence  $\{x_n\}$  and  $\delta_n \leq 1/n$  such that  $\overline{B_{\delta_n}(x_n)} \subset E_n \cap B_{\delta_{n-1}}(x_{n-1})$ .

For every  $m, n \geq N$ , we have  $x_n \in B_{\delta_n}(x_n) \subset \cdots \subset B_{\delta_N}(x_N)$  and  $x_m \in B_{\delta_m}(x_m) \subset \cdots \subset B_{\delta_N}(x_N)$  by construction. Hence  $d(x_n, x_m) \leq 2\delta_N \leq 2/N$  and  $\{x_n\}$  is a Cauchy sequence. Since  $X$  is complete,  $\{x_n\}$  converges to some  $x \in X$ . We claim that  $x \in E \cap G$ . Clearly  $x \in G$ . By construction  $x_m \in \overline{B_{\delta_n}(x_n)}$  for all  $m \geq n$ . Thus  $x \in B_{\delta_n}(x_m) \subset E_n$  for  $m \geq n \geq N$ . We see that  $x \in \cap_n E_n$ . Notice that  $G$  is arbitrary, so  $E$  is dense in  $X$ , proving that  $X$  is a Baire space. ■

**Theorem 2.52** (Uniform Boundedness Principle I)

$X$  is a complete metric space.  $f_\alpha : X \rightarrow \mathbb{R}$  is continuous for every  $\alpha \in A$ , where  $A$  is an index set. If for every  $x \in X$ , there exists  $M(x) < \infty$  such that

$$\sup_{\alpha \in A} |f_\alpha(x)| \leq M(x),$$

then there exists an open  $G$  and a constant  $C < \infty$  such that

$$\sup_{\alpha \in A} |f_\alpha(x)| \leq C$$

for all  $x \in G$ .

*Proof.* By Baire Category Theorem,  $X$  is a Baire space. For each  $n$ , let

$$F_n = \left\{ x \in X \mid \sup_{\alpha \in A} |f_\alpha(x)| \leq n \right\}.$$

We claim that  $F_n$  is closed and  $X = \cup_n F_n$ . Indeed, set  $x_k \rightarrow x \in X$ , where  $x_k \in F_n$  for all  $k$ . For any  $\alpha \in A$ ,  $|f_\alpha(x_k)| \leq n$  for all  $k$  and by continuity of  $f_\alpha$ ,

$$|f_\alpha(x)| = \lim_{k \rightarrow \infty} |f_\alpha(x_k)| \leq n.$$

Hence  $x \in F_n$  and  $F_n$  is closed. Next, for any  $x \in X$ , take  $N \geq M(x)$ . Then  $x \in F_N \subset \cup_n F_n$ . This shows that  $X = \cup_n F_n$ .



Finally, observe that  $F_n$  cannot have empty interiors for all  $n$ . Otherwise,  $\emptyset = X^c = (\cup_n F_n)^c = \cap F_n^c \neq \emptyset$  since  $F_n^c$  are open dense subsets of  $X$ , which is absurd. Hence there is some  $n$  such that  $F_n$  has nonempty interior, say  $G \subset F_n$ . Then  $\sup_{\alpha \in A} |f_\alpha(x)| \leq n$  for all  $x \in G$  as desired. ■

### Definition 2.53

A function  $f : X \rightarrow \mathbb{R}$  is said to be **sub-additive** if  $f(x + y) \leq f(x) + f(y)$  for all  $x, y \in X$ .

### Theorem 2.54 (Uniform Boundedness Principle II)

$X$  is a Banach space.  $\alpha \in A$  is an arbitrary index set.  $f_\alpha : X \rightarrow \mathbb{R}$  are continuous, sub-additive and satisfy  $f_\alpha(cx) = |c| f_\alpha(x)$  for all  $x \in X$  and  $c \in \mathbb{R}$ . If for every  $x \in X$ , there exists  $M(x) < \infty$  such that

$$\sup_{\alpha \in A} |f_\alpha(x)| \leq M(x),$$

then there exists a constant  $C < \infty$  such that

$$\sup_{\alpha \in A} |f_\alpha(x)| \leq C \|x\|_X$$

for all  $x \in X$ .

*Proof.* By theorem 2.52, there exists an open  $G$  and a constant  $C < \infty$  such that

$$\sup_{\alpha \in A} |f_\alpha(x)| \leq C$$

for all  $x \in G$ . The proof will be complete if we can extend  $G$  to  $X$ . Since  $G$  is open, there exists  $r > 0$  such that  $B_r(z) \subset G$  for all  $z \in G$ . For any  $x \in B_r(z)$ ,  $\sup_{\alpha \in A} |f_\alpha(x)| \leq C$  and hence  $\sup_{\alpha \in A} |f_\alpha(z + y)| \leq C$  for all  $y \in B_r(0)$ . Take  $y$  with  $\|y\| \leq r/2$ . Then

$$-2C \leq f_\alpha(y + z) - f_\alpha(z) \leq f_\alpha(y) \leq f_\alpha(y + z) + f_\alpha(-z) = f_\alpha(y + z) + f_\alpha(z) \leq 2C.$$

Hence  $|f_\alpha(y)| \leq 2C$  for all  $y$  with  $\|y\| \leq r/2$ . Take  $x \in X$ .

$$|f_\alpha(x)| = \left| f_\alpha\left(\frac{x}{\|x\|} \frac{r}{2}\right) \right| = \frac{r}{2\|x\|} |f_\alpha(y)| \leq \frac{4C}{r} \|x\|.$$

Thus

$$\sup_{\alpha \in A} |f_\alpha(x)| \leq \frac{4C}{r} \|x\|$$

for all  $x \in X$ . ■

### Corollary 2.55

$X$  is a Banach space.  $L_\alpha \in X'$  and  $\alpha \in A$ . If for every  $x \in X$ , there exists  $M(x) < \infty$  such that  $\sup_{\alpha \in A} |L_\alpha(x)| \leq M(x)$ , then there exists a constant  $C < \infty$  such that  $\sup_{\alpha \in A} \|L_\alpha\| \leq C$ .

*Proof.* Apply **theorem 2.54** to  $f_\alpha(x) = |L_\alpha(x)|$ . First,  $L_\alpha$  is linear and the sub-linearity follows from the triangle inequality. Next,  $|L_\alpha(cx)| = |c| |L_\alpha(x)|$  for all  $c \in \mathbb{R}$ . Also,  $L_\alpha \in X'$  implies that  $f_\alpha$  is continuous. The conclusion follows from **theorem 2.54**. ■

### Corollary 2.56

*X is a normed space.  $x_\alpha \in X$  for all  $\alpha \in A$  with the property that for every  $L \in X'$ , there is  $M(L) < \infty$  such that  $\sup_\alpha |L(x_\alpha)| \leq M(L)$  and  $(X', \|\cdot\|_{X \rightarrow \mathbb{R}})$  is a Banach space. Then there exists  $C < \infty$  such that  $\|x_\alpha\|_X \leq C$  for all  $\alpha \in A$ .*

*Proof.* Apply the **theorem 2.54** to  $f_\alpha(L) = |L(x_\alpha)|$ . First, for  $L, T \in X'$ ,

$$f_\alpha(L + T) = |L(x_\alpha) + T(x_\alpha)| \leq |L(x_\alpha)| + |T(x_\alpha)| = f_\alpha(L) + f_\alpha(T).$$

Next, for  $c \in \mathbb{R}$ ,

$$f_\alpha(cL) = |cL(x_\alpha)| = |c| |L(x_\alpha)| = |c| f_\alpha(L).$$

Finally, to verify that  $f_\alpha$  is continuous, note that for  $L_n \rightarrow L$  in  $X'$ ,

$$|f_\alpha(L_n) - f_\alpha(L)| = |L_n(x_\alpha) - L(x_\alpha)| \leq \|L_n - L\|_{X' \rightarrow \mathbb{R}} \|x_\alpha\|_X \rightarrow 0$$

for each  $\alpha \in A$ . The conclusion follows from **theorem 2.54**. ■

### Corollary 2.57

*X is a normed space and  $x_n \in X$  with  $x_n \xrightarrow{w} x$  in  $X$ . Then there exists  $C < \infty$  such that  $\|x_n\|_X \leq C$  for all  $n$ .*

*Proof.* This is a direct consequence of **corollary 2.56** with  $A = \mathbb{N}$ . ■

### Proposition 2.58

*Let  $f_n \in \mathcal{L}^p(X, \mu)$  and  $1 \leq p < \infty$ . Then  $f_n \xrightarrow{w} f \in \mathcal{L}^p$  if*

$$\lim_{n \rightarrow \infty} \int f_n g d\mu = \int f g d\mu$$

*for all  $g \in \mathcal{L}^{p'}(X, \mu)$  and some  $f$  in  $\mathcal{L}^p$  where  $p'$  is the conjugate exponent of  $p$ .*

*Proof.* By the assumption and Riesz representation theorem, for every  $T \in (\mathcal{L}^p)'$ , there exists a unique  $g \in \mathcal{L}^{p'}$  such that

$$T(f_n) = \int f_n g d\mu \rightarrow \int f g d\mu = T(f).$$

Hence  $f_n \xrightarrow{w} f$ . ■

### Proposition 2.59

*$f_n \in \mathcal{L}^p(X, \mu)$  and  $1 \leq p < \infty$ . If  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$ , then  $f_n$  is bounded and*

$$\|f_n\|_p \leq \liminf_{n \rightarrow \infty} \|f_n\|_p.$$

*Proof.* Consider the function

$$g = \frac{|f|^{p/p'}}{\|f\|_p^{p/p'}}.$$

Note that

$$\|g\|_{p'}^{p'} = \int |g|^{p'} d\mu = \int \frac{|f|^p}{\|f\|_p^p} d\mu = 1.$$

Hence  $g \in \mathcal{L}^{p'}$  with  $\|g\|_{p'} = 1$ . Also notice that  $|g| = |f|^{p/p'} / \|f\|_p^{p/p'} = |f|^{p-1} / \|f\|_p^{p-1}$ . By the weak convergence and Riesz representation theorem,

$$\|f\|_p = \int \frac{|f|^p}{\|f\|_p^{p-1}} d\mu = \int |fg| d\mu = \lim_{n \rightarrow \infty} \int |f_n g| d\mu \leq \liminf_{n \rightarrow \infty} \|f_n\|_p \|g\|_{p'} = \liminf_{n \rightarrow \infty} \|f_n\|_p$$

by the Hölder inequality. Note that by [corollary 2.57](#),  $f_n$  is bounded uniformly in  $n$ . ■

### Proposition 2.60

$1 \leq p < \infty$  and  $1/p + 1/p' = 1$ . Suppose  $f_n \rightarrow f$  in  $\mathcal{L}^p$  and  $g_n \rightarrow g$  in  $\mathcal{L}^{p'}$ . Then

$$\lim_{n \rightarrow \infty} \int f_n g_n d\mu = \int f g d\mu.$$

*Proof.* By the Hölder inequality,

$$\begin{aligned} \left| \int f_n g_n d\mu - \int f g d\mu \right| &\leq \left| \int f_n (g_n - g) d\mu \right| + \left| \int (f_n - f) g d\mu \right| \\ &\leq \|f_n\|_p \|g_n - g\|_{p'} + \|f_n - f\|_p \|g\|_{p'}. \end{aligned}$$

Note that by [proposition 2.59](#),  $f_n$  converges to  $f$  strongly and hence weakly. It follows that  $\|f_n\|$  is bounded by some  $C < \infty$ . Since  $g_n$  converges to  $g$  and  $f_n$  converges to  $f$  in their respective norms, the right hand side of the inequality converges to 0 as  $n \rightarrow \infty$ . ■

### Remark

If we loosen the condition to  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$  and  $g_n \xrightarrow{w} g$  in  $\mathcal{L}^{p'}$ , then the conclusion fails.

### Example

Suppose  $p = p' = 2$  and  $f_n(x) = \sqrt{2/\pi} \sin(nx)$  for  $x \in [0, \pi]$ . Then  $f_n \in \mathcal{L}^2([0, \pi])$  and

$$\int_0^\pi f_n^2 dx = \frac{2}{\pi} \int_0^\pi \sin^2(nx) dx = 1.$$

To see that  $f_n \xrightarrow{w} 0$ , let  $g \in \mathcal{L}^2([0, \pi])$ . For every  $\epsilon > 0$ , there is a step function  $\phi$  such that  $\|g - \phi\|_2 < \epsilon$ . Note that every step function is a finite linear combination of characteristic functions of intervals. Hence it suffices to show that  $f_n \chi_I$  can be arbitrary small for  $n$  sufficiently large. On every interval,

$$\left| \int_I \sin(nx) dx \right| \leq \int_0^{\pi/n} \sin(nx) dx = \frac{2}{n} \rightarrow 0$$

as  $n \rightarrow \infty$ . Thus  $f_n \xrightarrow{w} 0$  in  $\mathcal{L}^2([0, \pi])$ . However,  $f_n$  does not converge to 0 strongly in  $\mathcal{L}^2([0, \pi])$  since  $\|f_n\|_2 = 1 \neq 0$  for all  $n$ .

**Proposition 2.61**

$1 \leq p < \infty$ . Let  $f_n \in \mathcal{L}^p(X, \mu)$  be a bounded sequence of functions. Then  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$  if and only if

$$\lim_{n \rightarrow \infty} \int_A f_n d\mu = \int_A f d\mu$$

for all  $A \in \mathcal{A}$  when  $p = 1$  and for  $A$  with finite measure when  $p > 1$ .

*Proof.*

$$\begin{aligned} f_n \xrightarrow{w} f &\iff \int f_n g d\mu \rightarrow \int f g d\mu \text{ for all } g \in \mathcal{L}^{p'} \\ &\iff \int_A f_n s d\mu \rightarrow \int_A f s d\mu \text{ for all simple } s \in \mathcal{L}^{p'} \\ &\iff \int_A f_n d\mu = \int f_n \chi_A d\mu \rightarrow \int f \chi_A d\mu = \int_A f d\mu \end{aligned}$$

for  $A \in \mathcal{A}$  such that  $\chi_A \in \mathcal{L}^{p'}$ . If  $p = 1$ , then  $A$  can be taken to be any  $A \in \mathcal{A}$ ; if  $p > 1$ , then  $A$  must have finite measure. ■

**Proposition 2.62**

$1 < p < \infty$ . Let  $f_n \in \mathcal{L}^p(X, \mu)$  be a sequence with  $\|f_n\|_p \leq M$  and  $f_n \rightarrow f$  pointwise a.e. Then  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$ .

*Proof.* Since  $\|f_n\|_p \leq M$ ,

$$\int |f|^p d\mu = \int \liminf_{n \rightarrow \infty} |f_n|^p d\mu \leq \liminf_{n \rightarrow \infty} \int |f_n|^p d\mu = M^p$$

by Fatou's lemma. Hence  $f \in \mathcal{L}^p$ . It remains to show that the convergence is weak. By [proposition 2.61](#), it is equivalent to show that

$$\lim_{n \rightarrow \infty} \int_A f_n d\mu = \int_A f d\mu$$

for all  $A \in \mathcal{A}$  with  $\mu(A) < \infty$ . Indeed, by Egorov's theorem, for every  $\epsilon > 0$ , there exists  $F_\epsilon \subset A$  with  $\mu(A - F_\epsilon) \leq \epsilon$  and  $f_n \rightarrow f$  uniformly on  $F_\epsilon$ . Furthermore, by [proposition 1.33](#), we can choose  $F_\epsilon$  so that

$$\int_{A - F_\epsilon} |f_n - f|^p d\mu \leq \epsilon$$

since  $f_n, f \in \mathcal{L}^p$  and so does  $|f_n - f|^p$ . Also, let  $E = \{x \in A - F_\epsilon \mid |f_n - f| > 1\}$ . Then for  $n$

sufficiently large,

$$\begin{aligned}
\int_A |f_n - f| d\mu &\leq \int_{F_\epsilon} |f_n - f| d\mu + \int_{A-F_\epsilon} |f_n - f| d\mu \\
&\leq \int_A \epsilon d\mu + \int_{A-F_\epsilon-E} |f_n - f| d\mu + \int_E |f_n - f| d\mu \\
&\leq \epsilon\mu(A) + \mu(A - F_\epsilon) + \int_{A-F_\epsilon} |f_n - f|^p d\mu \leq \epsilon\mu(A) + \epsilon + \epsilon.
\end{aligned}$$

Hence  $f_n \xrightarrow{w} f$ . ■

### Remark

The proposition fails for  $p = 1$ . Consider  $f_n = n\chi_{[0,1/n]}$ . Then  $\|f_n\|_1 = 1$  and  $f_n \rightarrow 0$  pointwise a.e. However,

$$\int_0^1 f_n(x) dx = 1 \neq 0 = \int_0^1 0 dx.$$

Thus  $f_n$  does not converge weakly to 0 in  $\mathcal{L}^1$ .

### Theorem 2.63 (Radon-Riesz)

$1 < p < \infty$ . Then  $f_n \rightarrow f$  in  $\mathcal{L}^p$  if and only if  $\lim_{n \rightarrow \infty} \|f_n\|_p = \|f\|_p$  and  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$ .

*Proof.* Suppose  $f_n \rightarrow f$  in  $\mathcal{L}^p$ . Then the strong convergence immediately implies the weak convergence. Also, note that  $\|f_n\|_p \leq \|f_n - f\|_p + \|f\|_p$  and thus

$$|\|f_n\|_p - \|f\|_p| \leq \|f_n - f\|_p \rightarrow 0$$

by the strong convergence. Conversely, suppose that  $\|f_n\|_p \rightarrow \|f\|_p$  and  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$ .

Assume  $p \geq 2$ . For any  $y \in \mathbb{R}$ , notice that  $|1 + y|^p \geq 1 + py + c|y|^p$  for some  $c \in (0, 1)$ . Let  $E = \{x \in X \mid f(x) = 0\}$  and apply  $y = (f_n - f)/f$  on  $E^c$ . Then on  $E^c$ ,

$$\left| \frac{f_n}{f} \right|^p \geq 1 + p \left( \frac{f_n - f}{f} \right) + c \left| \frac{f_n - f}{f} \right|^p$$

Thus

$$|f_n|^p \geq |f|^p + p(f_n - f)|f|^{p-1} \operatorname{sgn}(f) + c|f_n - f|^p.$$

Rearranging the inequality and integrating both sides on  $E^c$  gives

$$c \int_{E^c} |f_n - f|^p d\mu \leq \int_{E^c} |f_n|^p - |f|^p d\mu - p \int_{E^c} |f|^{p-1} \operatorname{sgn}(f)(f_n - f) d\mu$$

Note that as shown in the proof of [proposition 2.59](#),  $|f|^{p-1} \operatorname{sgn}(f) \in \mathcal{L}^{p'}$ . By the assumptions we see that

$$\int_{E^c} |f_n - f|^p d\mu \rightarrow 0$$

as  $n \rightarrow \infty$ . On  $E$ , we have  $f = 0$  and

$$\int_E |f_n - f|^p d\mu = \int_E |f_n|^p d\mu \rightarrow 0$$

as  $n \rightarrow \infty$ . Hence  $f_n \rightarrow f$  in  $\mathcal{L}^p$ .

Assume  $1 < p < 2$ . Then we have the same inequality for  $|z| \geq 1$ , i.e.,

$$|1 + z|^p \geq 1 + p|z| + c|z|^p$$

Also, for  $|z| \leq 1$ ,

$$\frac{|1 + z|^p - 1 - pz}{z^2}$$

is strictly positive. Now let  $E_n = \{x \in X \mid |f_n(x) - f(x)| \leq |f(x)|\}$ . Then by applying the same argument above on  $E_n^c$ , we have

$$\int_{E_n^c} |f_n - f|^p d\mu \leq \frac{1}{c} \int_{E_n^c} |f_n|^p - |f|^p d\mu - \frac{p}{c} \int_{E_n^c} |f|^{p-1} \operatorname{sgn}(f)(f_n - f) d\mu$$

as  $n \rightarrow \infty$ . On  $E_n$ , replacing  $z$  by  $(f_n - f)/f$ ,

$$\left| \frac{f_n}{f} \right|^p \geq 1 + p \frac{f_n - f}{f} + c' \left( \frac{f_n - f}{f} \right)^2 \implies |f_n|^p \geq |f|^p + p(f_n - f)|f|^{p-1} \operatorname{sgn}(f) + c'|f_n - f|^2 |f|^{p-2}$$

for some  $c' > 0$ . Thus

$$\int_{E_n} |f_n - f|^2 |f|^{p-2} d\mu \leq \frac{1}{c'} \int_{E_n} |f_n|^p - |f|^p d\mu - \frac{p}{c'} \int_{E_n} |f|^{p-1} \operatorname{sgn}(f)(f_n - f) d\mu.$$

Adding up the two inequalities, we have

$$\int_{E_n^c} |f_n - f|^p d\mu + \int_{E_n} |f_n - f|^2 |f|^{p-2} d\mu \rightarrow 0$$

as  $n \rightarrow \infty$  by the assumptions. Note that on  $E_n$ ,  $|f| \geq |f_n - f|$  and

$$\begin{aligned} \int_{E_n} |f_n - f|^p d\mu &\leq \int_{E_n} |f_n - f| |f|^{p-1} d\mu \leq \left( \int_{E_n} |f_n - f|^2 |f|^{p-2} d\mu \right)^{1/2} \left( \int_{E_n} |f|^p d\mu \right)^{1/2} \\ &\leq \left( \int_{E_n} |f_n - f|^2 |f|^{p-2} d\mu \right)^{1/2} \|f\|_p^{p/2} \rightarrow 0. \end{aligned}$$

Hence  $f_n \rightarrow f$  in  $\mathcal{L}^p$ . We conclude that  $f_n \rightarrow f$  strongly in  $\mathcal{L}^p$  if and only if  $f_n \xrightarrow{w} f$  in  $\mathcal{L}^p$  and  $\|f_n\|_p \rightarrow \|f\|_p$ . ■

### Remark

*Radon-Riesz theorem fails for  $p = 1$ . Consider  $f_n(x) = 1 + \sin(nx)$  on  $X = [-\pi, \pi]$ . Then for*

every  $g \in \mathcal{L}^\infty$ ,

$$\int (f_n - 1)g d\mu \leq \int \sin(nx)g d\mu \rightarrow 0$$

by the step function approximation argument. Also,  $\|f_n\|_1 = 2\pi$  for all  $n$  and hence converges to  $\|1\|_1 = 2\pi$ . However,  $f_n$  does not converge to 1 in  $\mathcal{L}^1$  since

$$\int_{-\pi}^{\pi} |f_n - 1| d\mu = \int_{-\pi}^{\pi} |\sin(nx)| d\mu = 2n \int_0^{\frac{\pi}{2n}} \sin(nx) dx = 2$$

for all  $n$ .

#### Definition 2.64

Let  $X$  be a Banach space. A subset  $K \subset X$  is **weakly sequentially compact** if every sequence  $\{x_n\} \subset K$  has a subsequence  $\{x_{n_k}\}$  such that  $x_{n_k} \xrightarrow{w} x \in K$ .

#### Proposition 2.65

Let  $X$  be a Banach space. If  $K \subset X$  is weakly sequentially compact, then  $K$  is bounded.

*Proof.* Suppose  $K$  is not bounded. Then we can choose an unbounded sequence  $\{x_n\} \subset K$  such that  $\|x_n\| \geq n$  for all  $n$ . By the weakly sequential compactness of  $K$ , there exists a weakly convergent subsequence  $\{x_{n_k}\}$  such that  $x_{n_k} \xrightarrow{w} x \in K$  and also  $\|x_{n_k}\| \geq n_k$ . However, by [corollary 2.57](#),  $\|x_{n_k}\| \leq C$  for some  $C < \infty$ , which is absurd. Hence  $K$  is bounded. ■

#### Theorem 2.66 (Kakutani)

Let  $X$  be a reflexive Banach space. Then the closed unit ball

$$B = \{x \in X \mid \|x\| \leq 1\}$$

is weakly sequentially compact.

*Proof.* We consider first the case when  $X$  is separable. Reflexivity gives that  $X'' \cong X$  and hence  $X''$  is separable. By [theorem 2.46](#),  $X'$  is separable, and there exists a countable dense subset  $\{m_j\} \subset X'$ . Given  $x_n \in X$  with  $\|x_n\| \leq 1$ , we need to show that there exists a subsequence  $x_{n_k} \xrightarrow{w} x \in B$ . Since  $m_j(x_n)$  is a bounded sequence for each  $j$ , we can extract a subsequence  $x_{n_k}$  such that  $m_j(x_{n_{k_j}}) \rightarrow A(m_{k_j})$  as  $j \rightarrow \infty$ , where

$$A(m_j) = \lim_{j \rightarrow \infty} m_j(x_{n_{k_j}}).$$

We claim that for all  $m \in X'$ ,  $m(x_{n_k}) \rightarrow A(m)$  as  $k \rightarrow \infty$ . Indeed, for any  $m \in X'$ , we can find a sequence  $\{m_j\}$  such that  $m_j \rightarrow m$  as  $j \rightarrow \infty$ . Then

$$\begin{aligned} |m(x_{n_k}) - m(x_{n_l})| &\leq |m(x_{n_k}) - m_j(x_{n_k})| + |m_j(x_{n_k}) - m_j(x_{n_l})| + |m_j(x_{n_l}) - m(x_{n_l})| \\ &\leq \|m - m_j\| \|x_{n_k}\| + |m_j(x_{n_k}) - m_j(x_{n_l})| + \|m - m_j\| \|x_{n_l}\| \\ &\leq 2\|m - m_j\| + |m_j(x_{n_k}) - m_j(x_{n_l})| \rightarrow 0 \end{aligned}$$

as  $k, l \rightarrow \infty$ . Hence the sequence  $\{m(x_{n_k})\}$  is Cauchy and  $A$  is well-defined. Notice that  $A$  is also bounded:

$$|A(m)| = \lim_{k \rightarrow \infty} |m(x_{n_k})| \leq \lim_{k \rightarrow \infty} \|m\| \|x_{n_k}\| = \|m\|.$$

We see that  $\|A\| \leq 1$ . Because  $A$  is bounded, it is continuous and thus  $m(x_{n_k}) \rightarrow m(x)$  for some  $x \in X$  by the reflexivity of  $X$ . Such  $x$  belongs to  $B$  since  $\|x\| = \|A\| \leq 1$ . Thus  $B$  is weakly sequentially compact.

For the general case where  $X$  is not separable, consider the sequence  $\{x_n\} \subset B$ . Let  $Y = \{\sum_{n=1}^N \alpha_n x_n \mid N \in \mathbb{N}, \alpha_n \in \mathbb{R}\}$  be the closed subspace of  $X$  spanned by  $\{x_n\}$ . Since  $X$  is reflexive,  $Y$  is also reflexive by [theorem 2.47](#). Note that  $Y$  is also separable. The established results above show that there exists a subsequence  $\{x_{n_k}\} \subset Y$  and  $x \in Y$  such that  $x_{n_k} \xrightarrow{w} x$  in  $Y$ , i.e., for every  $m \in Y'$ ,  $m(x_{n_k}) \rightarrow m(x)$ . Extend the functionals  $m \in Y'$  to  $\ell \in X'$  by Hahn-Banach theorem. Then  $\ell|_Y = m \in Y'$  implies that  $\ell(x_{n_k}) = m(x_{n_k}) \rightarrow m(x) = \ell(x)$ . We conclude that  $x_{n_k} \xrightarrow{w} x \in B$ . Thus  $B$  is weakly sequentially compact. ■

### Example

Let  $p \in (1, \infty)$ . Then  $\mathcal{L}^p(\Omega, \mu)$  is reflexive. Then for all  $\{f_n\}$  with  $\|f_n\|_p \leq 1$ , there exists a subsequence  $f_{n_k} \xrightarrow{w} f$  in  $\mathcal{L}^p$  for some  $f$  with  $\|f\|_p \leq 1$ . By Riesz representation theorem, this is equivalent to saying that for every  $g \in \mathcal{L}^q(\Omega, \mu)$ ,

$$\lim_{k \rightarrow \infty} \int f_{n_k} g d\mu = \int f g d\mu,$$

where  $q$  is the conjugate exponent of  $p$ .

### Definition 2.67

Let  $M$  be a Banach space. A sequence of bounded linear functionals  $\{x_n\} \subset M'$  **converges weakly\*** to  $x$  if for all  $m \in M$ ,  $x_n(m) \rightarrow x(m)$  as  $n \rightarrow \infty$ . We denote the convergence by  $x_n \xrightarrow{w^*} x$ .

### Remark

Since the canonical mapping  $M \rightarrow M''$  is always injective,  $w^*$  convergence is weaker than weak convergence. Allowing for the abuse of notation, we can write  $M \subset M''$ . Consider now a sequence  $x_n \in M'$  with  $x_n \xrightarrow{w} x$  in  $M'$ . Then  $\ell(x_n) \rightarrow \ell(x)$  for any  $\ell \in M''$ . This implies that  $x_n(m) \rightarrow x(m)$  for all  $m \in M$  and hence  $x_n \xrightarrow{w^*} x$  in  $M'$ . Thus weak convergence implies  $w^*$  convergence.

The converse is true if  $M$  is reflexive. However, once we remove the reflexivity condition, the converse fails. Let  $X$  be the space of finite signed measures on  $[-1, 1]$ . We have already seen in [theorem 2.39](#) that  $C([-1, 1])' \cong X$ . Consider the measures  $\nu_n(A) = n\mu(A \cap [-1/n, 1/n])/2$ . We claim that  $\nu_n \xrightarrow{w^*} \delta_0$ , where  $\delta_0$  is the Dirac measure at 0. Indeed, for any  $f \in C([-1, 1])$ , the corresponding functional  $\ell_n$  for  $\nu_n$  is given by

$$\ell_n(f) = \int_{-1}^1 f d\nu_n = \frac{n}{2} \int_{-1/n}^{1/n} f(x) dx \rightarrow f(0) = \ell_0(f),$$



where  $\ell_0$  is the functional defined as  $\ell_0 : f \mapsto f(0)$ . Thus  $\ell_n \xrightarrow{w^*} \ell_0$ .

However,  $\ell_n$  is not weakly convergent to  $\ell_0$ . To see this, consider the evaluation functional  $L_{\{0\}} : \ell \mapsto (\phi\ell)(\{0\})$ , where  $\phi$  is the isometric isomorphism from  $C([-1, 1])'$  to  $X$ . Then  $L_{\{0\}} \in X' = M''$ . However,

$$L_{\{0\}}(\ell_n) = \nu_n(\{0\}) = 0 \not\rightarrow 1 = \delta_0(\{0\}) = L_{\{0\}}(\ell_0).$$

Thus  $\nu_n$  does not converge weakly to  $\delta_0$ .

### Definition 2.68

Let  $M$  be a Banach space. A subset  $K \subset M'$  is **weakly\* sequentially compact** if every sequence  $\{x_n\} \subset K$  has a subsequence  $\{x_{n_k}\}$  such that  $x_{n_k} \xrightarrow{w^*} x \in K$ .

### Theorem 2.69 (Banach-Alaoglu)

Let  $M$  be a separable Banach space. Then the closed unit ball

$$B = \{x \in M' \mid \|x\| \leq 1\}$$

is weakly\* sequentially compact.

*Proof.* This proof is similar to the proof of Kakutani theorem. Let  $\{x_n\} \subset B$  be a sequence. By the separability of  $M$ , for every  $m \in M$ , there is a sequence  $\{m_j\} \subset M$  such that  $m_j \rightarrow m$  as  $j \rightarrow \infty$ . For any fixed  $j$ ,  $|x_n(m_j)| \leq \|m_j\|$  is a bounded sequence. Hence we can extract a subsequence  $x_{n_k}$  such that

$$x_{n_k}(m_j) \rightarrow A(m_j) \text{ as } k \rightarrow \infty$$

for some  $A(m_j) \in \mathbb{R}$ , with  $A(m) = \lim_{k \rightarrow \infty} x_{n_k}(m)$ .  $A$  is a bounded linear functional on  $M$  since it is clearly linear and

$$|A(m)| = \lim_{k \rightarrow \infty} |x_{n_k}(m)| \leq \lim_{k \rightarrow \infty} \|x_{n_k}\| \|m\| = \|m\|,$$

showing that  $\|A\| \leq 1$ . We claim that  $A$  is well-defined, i.e., the limit exists. Indeed, for any  $m \in M$ , we can find a sequence  $\{m_j\}$  such that  $m_j \rightarrow m$  as  $j \rightarrow \infty$ . Then

$$\begin{aligned} |x_{n_k}(m) - x_{n_l}(m)| &\leq |x_{n_k}(m) - x_{n_k}(m_j)| + |x_{n_k}(m_j) - x_{n_l}(m_j)| + |x_{n_l}(m_j) - x_{n_l}(m)| \\ &\leq \|x_{n_k} - x_{n_l}\| \|m\| + |x_{n_k}(m_j) - x_{n_l}(m_j)| + \|x_{n_k} - x_{n_l}\| \|m\| \\ &\leq 2\|x_{n_k} - x_{n_l}\| + |x_{n_k}(m_j) - x_{n_l}(m_j)| \rightarrow 0 \end{aligned}$$

as  $k, l \rightarrow \infty$ . Hence the sequence  $\{x_{n_k}(m)\}$  is Cauchy and  $A$  is well-defined. Because of the boundedness of  $A$ , it is continuous and thus  $x_{n_k}(m) \rightarrow x(m)$  for some  $x \in M'$ . Since  $\|x_{n_k}\| \leq 1$ , we have  $\|x\| \leq 1$ . Thus  $x \in B$  and  $x_{n_k} \xrightarrow{w^*} x$  in  $M'$ . We conclude that  $B$  is weakly\* sequentially compact. ■

**Remark**

Let  $X, Y$  be Banach spaces and  $B(X, Y)$  be the space of bounded linear operators from  $X$  to  $Y$ . Consider the following topologies on  $B(X, Y)$ :

- The uniform topology on  $B(X, Y)$  is the topology induced by the uniform norm:

$$\|m\|_{B(X, Y)} = \sup_{\|x\|_X \leq 1} \|m(x)\|_Y.$$

This is the coarsest among the three topologies.

- The strong topology on  $B(X, Y)$  is the topology generated by the collection of sets:

$$\{B_{x, \epsilon}(T) = \{S \in B(X, Y) \mid \|Sx - Tx\| < \epsilon\} \mid \epsilon > 0, x \in X, T \in B(X, Y)\}.$$

This is the coarsest topology that makes the evaluation map  $m \mapsto m(x)$  continuous for all  $x \in X$ .

- The weak topology on  $B(X, Y)$  is the topology generated by the collection

$$\{B_{y', x, \epsilon}(T) = \{S \in B(X, Y) \mid |y'Sx - y'Tx| < \epsilon\} \mid \epsilon > 0, y' \in Y', x \in X, T \in B(X, Y)\}.$$

**Definition 2.70**

Let  $\{M_n\} \subset B(X, Y)$  be a sequence of bounded linear operators.  $M_n$  **converges strongly** if for any  $x \in X$ ,  $\|M_n x - y\|_Y \rightarrow 0$  for some  $y \in Y$ .

**Proposition 2.71**

If  $\{M_n\} \subset B(X, Y)$  converges strongly, then there is an  $M \in B(X, Y)$  such that  $\|M_n x - Mx\|_Y \rightarrow 0$  for all  $x \in X$ .

*Proof.* Set  $Mx = \lim_{n \rightarrow \infty} M_n x$  for all  $x \in X$ . We check that  $M \in B(X, Y)$ . Linearity is trivial; we check the boundedness. Let  $f_n(x) = \|M_n x\|_Y$ . Then  $f_n$  is sub-additive and  $f_n(\alpha x) = |\alpha| f_n(x)$  for all  $\alpha \in \mathbb{R}$ . If  $x_k \rightarrow x$ ,

$$f_n(x_k) = \|M_n x_k\|_Y \rightarrow \|M x_k\|_Y = f_n(x)$$

for any fixed  $n$  as  $k \rightarrow \infty$ ;  $f_n$  is continuous. Now for any fixed  $x \in X$ ,  $\sup_n f_n(x) = \sup_n \|M_n x\| \leq C(x)$  by the strong convergence. It follows from the uniform boundedness principle that there is  $C_0 < \infty$  such that  $|f_n(x)| \leq C_0 \|x\|_X$  for all  $n$ . Thus

$$\|Mx\|_Y = \lim_{n \rightarrow \infty} \|M_n x\|_Y \leq C_0 \|x\|_X.$$

Hence  $\|M\| \leq C_0$  and  $M \in B(X, Y)$ . ■

**Definition 2.72**

A sequence  $\{M_n\} \subset B(X, Y)$  **converges weakly** if for all  $x \in X$ ,  $M_n x \xrightarrow{w} y \in Y$  for some  $y$ .

**Proposition 2.73**

If  $\{M_n\} \subset B(X, Y)$  converges weakly, then there is an  $M \in B(X, Y)$  such that  $M_n x \xrightarrow{w} Mx$  for all  $x \in X$ .

*Proof.* Set  $Mx = \lim_{n \rightarrow \infty} M_n x$  for all  $x \in X$ . We check that  $M \in B(X, Y)$ . Linearity is trivial; we check the boundedness. Without loss of generality, we can assume that  $\|x\|_X = 1$ . Observe that  $\{M_n x\} \subset Y$  is a weakly convergence sequence and hence weakly sequentially compact; by **proposition 2.65**, it is bounded. Thus there exists  $C < \infty$  such that  $\|M_n x\|_Y \leq C = C \|x\|_X$  for all  $n$ . Taking the limit, we have  $\|Mx\|_Y = \lim_{n \rightarrow \infty} \|M_n x\|_Y \leq C \|x\|_X$ . Hence  $M \in B(X, Y)$ .

Lastly, we check the weak convergence. Indeed, for any  $\ell \in Y'$ ,  $\ell(M_n x) = \ell(Mx)$  for all  $x \in X$  since  $M_n x \rightarrow Mx$  in  $Y$ . Thus  $M_n x \xrightarrow{w} Mx$  in  $Y$ . ■

**2.6. Open Mapping Theorem and Closed Graph Theorem****Proposition 2.74**

If  $X$  is a Baire space and  $F_n$  is a sequence of closed sets in  $X$  such that  $\bigcup_{n=1}^{\infty} F_n = X$ , then there exists some  $n$  and a nonempty open set  $G$  such that  $G \subseteq F_n$ .

*Proof.* Let  $G_n = F_n^c$  be open sets in  $X$ . Then  $\bigcap_{n=1}^{\infty} G_n = \bigcap_{n=1}^{\infty} F_n^c = \left(\bigcup_{n=1}^{\infty} F_n\right)^c = \emptyset$ . By the Baire category theorem, at least one of the  $G_n$  is not dense in  $X$ . Thus there is some  $x \in G^c$  and an open neighborhood  $U$  of  $x$  such that  $U \cap G_n = \emptyset$ . This implies  $U \subseteq F_n$ . ■

**Theorem 2.75 (Open Mapping Theorem)**

Let  $X$  and  $Y$  be Banach spaces and  $T : X \rightarrow Y$  be a bounded surjective linear map. Then for any open set  $U \subset X$ ,  $T(U)$  is open in  $Y$ .

*Proof.* We first claim that for any open ball  $B$  centered at 0 in  $X$ ,  $\overline{T(B)}$  contains an open neighborhood of zero in  $Y$ . By the surjectivity,  $Y \subset T(X) = T(\bigcup_n nB) = \bigcup_n T(nB) \subset \bigcup_n \overline{T(nB)}$ . By **proposition 2.74**, there is some  $n$  such that  $\overline{T(nB)}$  contains an interior point, say  $y$ , and some open ball  $B_r(y) \subset \overline{T(nB)}$ . Then for every  $z \in Y$  with  $\|z\| < r$ ,  $z - y \in B_r(-y) \subset \overline{T(-nB)} = \overline{T(nB)}$  and

$$z = y + (z - y) \in y + B_r(-y) \subset \overline{T(nB)} + \overline{T(nB)} \subset \overline{T(2nB)}.$$

Deviding  $z$  by  $2n$  gives that  $z/2n \in \overline{T(B)}$  and  $B_{r/2n}(0) \subset \overline{T(B)}$ .

Next, let  $B$  be an unit ball. To shorten the notation, denote  $r/2n$  by  $\delta$  and  $B_{r/2n}(0)$  by  $B_\delta$ . Let  $y \in B_\delta$  and  $c_n > 0$  be a sequence. Since  $B_\delta \subset \overline{T(B)}$ ,  $\overline{B_\delta} \subset \overline{T(B)}$ . Thus for every  $z \in Y$  and  $\epsilon > 0$ , we can find some  $x \in X$  such that  $\|x\| < \delta^{-1} \|z\|$  and  $z \in B_\epsilon(T(x))$ . Now taking  $z = y$  and  $\epsilon = c_1$ , we can find an  $x_1$  such that  $\|x_1\| < \delta^{-1} \|y\|$  and  $\|y - Tx_1\| < c_1$ . Similarly, we can take  $z = y - Tx_1$  and  $\epsilon = c_2$  to find an  $x_2$  such that  $\|x_2\| < \delta^{-1} \|y - Tx_1\| < \delta^{-1} c_1$  and  $\|y - Tx_1 - Tx_2\| < c_2$ . Iductively, we find a sequence  $\{x_n\}$  such that  $\|x_n\| < \delta^{-1} c_{n-1}$  and  $\|y - T(\sum_{k=1}^n x_k)\| < c_n$ . Now we choose  $c_n = 2^{-n} c$  for arbitrary  $c > 0$ . Then

$$\left\| \sum_{k=1}^n x_k \right\| \leq \sum_{k=1}^n \|x_k\| \leq \frac{\|y\|}{\delta} + \sum_{k=2}^n \frac{c_{k-1}}{\delta} \leq \frac{\|y\|}{\delta} + \frac{c}{\delta} \sum_{k=1}^{\infty} 2^{-k} = \frac{\|y\|}{\delta} + \frac{c}{\delta}.$$

Hence  $\sum_n x_n$  converges in  $X$  to some  $x$  with  $\|x\| < 1$  by making  $c$  arbitrarily small. Also,

$$\left\| y - T\left(\sum_{k=1}^n x_k\right) \right\| \leq c_n = 2^{-n}c \rightarrow 0.$$

Thus  $Tx = y$  and  $y \in T(B)$ , which implies  $B_\delta \subset T(B)$ .

Finally, let  $U$  be an open set in  $X$ . Then for any  $y \in T(U)$ , there is some  $x \in U$  such that  $y = Tx$ . Since  $U$  is open, there is some  $\epsilon > 0$  such that  $B_\epsilon(x) \subset U$ . By the previous claim, there is some  $s > 0$  such that  $B_s(0) \subset T(B_1(0))$ . Multiplying both sides by  $\epsilon$  gives  $B_{s\epsilon}(0) \subset T(B_\epsilon(0))$ . Then

$$B_{s\epsilon}(y) = y + B_{s\epsilon}(0) \subset y + T(B_\epsilon(0)) = Tx + T(B_\epsilon(0)) = T(x + B_\epsilon(0)) = T(B_\epsilon(x)) \subset T(U).$$

Thus  $T(U)$  is open. This completes the proof. ■

**Theorem 2.76** (Bounded Inverse Theorem)

Let  $X$  and  $Y$  be Banach spaces and  $T : X \rightarrow Y$  be a bounded linear map. If  $T$  is bijective, then  $T^{-1}$  is bounded.

*Proof.* By the open mapping theorem, there is  $r > 0$  such that  $B_r(0) \subset T(B_1(0))$ . For any  $y \in Y$  with  $\|y\| = r/2$ , there exists  $x \in B_1(0)$  such that  $y = Tx$ . For  $z \in Y$ , write

$$z = \frac{rz}{2\|z\|} \frac{2}{r} \|z\|.$$

Then since  $\left\| \frac{rz}{2\|z\|} \right\| = r/2$ , there is some  $x \in B_1(0)$  such that  $\frac{rz}{2\|z\|} = Tx$ . Thus  $z = \frac{2}{r} \|z\| Tx$ ,

$$T^{-1}z = \frac{2}{r} \|z\| x \implies \|T^{-1}z\| \leq \frac{2}{r} \|z\| \|x\|.$$

Note that  $\|x\| \leq 1$ . We see that  $\|T^{-1}\|$  is bounded by  $2/r$ . ■

**Remark**

The completeness in the open mapping theorem is essential. For counterexample, consider  $X$  as the space of all sequences with finitely many nonzero terms equipped with the supremum norm. Define  $T : X \rightarrow X$  by

$$T(x_1, x_2, \dots) = \left(x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots\right).$$

Note that  $X$  is not complete since the sequence  $x^{(n)} = (1, 1/2, \dots, 1/n, 0, 0, \dots)$  converges to  $(1, 1/2, \dots)$ , which does not belong to  $X$ . In this case  $T^{-1}$  exists but is not bounded.

**Definition 2.77**

$X, Y$  are Banach spaces.  $T : X \rightarrow Y$  is a bounded linear map. The set

$$\Gamma(T) = \{(x, Tx) \in X \times Y \mid x \in X\}$$

is called the **graph** of  $T$ . We define the norm of  $x$  on the graph by

$$\|(x, Tx)\|_{\Gamma} = \|x\|_X + \|Tx\|_Y.$$

Note that  $(\Gamma(T), \|\cdot\|_{\Gamma})$  forms a normed space.

**Definition 2.78**

A linear map  $T : X \rightarrow Y$  is called **closed** if its graph is a closed, i.e., for any sequence  $x_n \in X$ , if  $x_n \rightarrow x \in X$  and  $Tx_n \rightarrow y \in Y$ , then  $Tx = y$  and  $(x, y) \in \Gamma(T)$ .

**Remark**

If  $T$  is bounded, it is closed. To see this, note that if  $x_n \rightarrow x \in X$ , by the continuity we have  $Tx_n \rightarrow Tx \in Y$ .

**Theorem 2.79** (Closed Graph Theorem)

Let  $X$  and  $Y$  be Banach spaces and  $T : X \rightarrow Y$  be a linear map. If  $T$  is closed, then  $T$  is bounded.

*Proof.* Observe that  $\Gamma(T)$  is a Banach space with the norm  $\|\cdot\|$  on  $\Gamma(T)$ . This follows from the closedness of  $T$ . Now define  $S : \Gamma(T) \rightarrow X$  by  $S(x, Tx) = x$ . We claim that  $S$  is bounded, linear and bijective. For linearity, let  $(x_1, Tx_1), (x_2, Tx_2) \in \Gamma(T)$  and  $c \in \mathbb{R}$ .

$$S(c(x_1, Tx_1) + (x_2, Tx_2)) = S(cx_1 + x_2, cTx_1 + Tx_2) = cx_1 + x_2 = cS(x_1, Tx_1) + S(x_2, Tx_2).$$

For boundedness,

$$\|S(x, Tx)\|_X = \|x\|_X \leq \|x\|_X + \|Tx\|_Y = \|(x, Tx)\|_{\Gamma}.$$

Thus  $\|S\| \leq 1$ . For bijectivity, notice that

$$S(x_1, Tx_1) = S(x_2, Tx_2) \implies x_1 = S(x_1, Tx_1) = S(x_2, Tx_2) = x_2.$$

and for any  $x \in X$ ,  $(x, Tx) \in \Gamma(T)$  and  $S(x, Tx) = x$ . Thus  $S$  is bounded, linear and bijective.

By the bounded inverse theorem,  $S^{-1} : X \rightarrow \Gamma(T)$  is bounded as well. For any  $x \in X$ ,

$$\|Tx\|_Y = \|(x, Tx)\|_{\Gamma} - \|x\|_X = \|S^{-1}x\|_{\Gamma} - \|x\|_X \leq C \|x\|_X - \|x\|_X = (C - 1) \|x\|_X$$

for some constant  $C < \infty$ . Thus  $T$  is bounded. ■

**Definition 2.80**

Suppose  $X$  is a vector space with norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$ . The norms are said to be **compatible** if  $x_n \rightarrow x$  in  $\|\cdot\|_1$  and  $x_n \rightarrow y$  in  $\|\cdot\|_2$  implies  $x = y$ .

**Definition 2.81**

Let  $X$  be a vector space with norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$ . The norms are said to be **equivalent** if

there are constants  $c_1, c_2 > 0$  such

$$c_1 \|x\|_1 \leq \|x\|_2 \leq c_2 \|x\|_1$$

for all  $x \in X$ .

**Proposition 2.82**

Suppose  $X$  is a vector space with norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$ . If the norms are equivalent, then they are compatible.

*Proof.* Suppose  $\|x_n - x\|_1 \rightarrow 0$  and  $\|x_n - y\|_2 \rightarrow 0$ . By the equivalence,  $\|x - y\|_1 \leq \|x - x_n\|_1 + \|x_n - y\|_1 \leq \|x - x_n\|_1 + c_2 \|x_n - y\|_2 \rightarrow 0$  for some  $c_2 > 0$ . Similarly,  $\|x - y\|_2 \leq c_1 \|x - x_n\|_1 + \|x_n - y\|_2 \rightarrow 0$  for some  $c_1 > 0$ . Thus  $x = y$ . ■

**Proposition 2.83**

If  $(X, \|\cdot\|_1)$  and  $(X, \|\cdot\|_2)$  are Banach spaces. Then the norms are equivalent.

*Proof.* By the closed graph theorem, the identity map  $I : (X, \|\cdot\|_1) \rightarrow (X, \|\cdot\|_2)$  is a closed linear map and thus bounded. Suppose  $x_n \rightarrow x$  in  $\|\cdot\|_1$ . Then  $x_n = Ix_n \rightarrow Ix = x$  in  $\|\cdot\|_2$  by the continuity of  $I$ . Since  $I$  is bounded,  $\|x\|_2 = \|Ix\|_2 \leq c_1 \|x\|_1$  for some  $c_1 > 0$ . Applying the same argument exchanging the roles of  $\|\cdot\|_1$  and  $\|\cdot\|_2$  gives  $\|x\|_1 \leq c_2 \|x\|_2$  for some  $c_2 > 0$ . Hence

$$\frac{1}{c_2} \|x\|_1 \leq \|x\|_2 \leq c_1 \|x\|_1$$

and the norms are equivalent. ■

## 3. Hilbert Space

### 3.1. Cauchy-Schwarz Inequality

#### Definition 3.1

Let  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .  $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{F}$  is an inner product on  $X$  if it satisfies

- (a)  $\langle cx + y, z \rangle = c \langle x, z \rangle + \langle y, z \rangle$  for all  $x, y, z \in X$  and  $c \in \mathbb{F}$ .
- (b)  $\langle x, y \rangle = \overline{\langle y, x \rangle}$  for all  $x, y \in X$ .
- (c)  $\langle x, x \rangle > 0$  for all  $x \neq 0$ .

#### Remark

An inner product automatically induces a norm on  $X$  by  $\|\cdot\| = \sqrt{\langle \cdot, \cdot \rangle}$ .

#### Definition 3.2

A **Hilbert space** is a complete vector space with an inner product inducing a norm that makes it a Banach space. We denote the Hilbert space by  $\mathcal{H}$ .

#### Remark

If  $X$  is a vector space with an inner product but not complete, then  $X$  is called a **pre-Hilbert space**.

#### Proposition 3.3 (Cauchy-Schwarz Inequality)

For all  $x, y \in \mathcal{H}$ ,

$$|\langle x, y \rangle| \leq \|x\| \|y\|.$$

Furthermore, equality holds if and only if  $x$  and  $y$  are linearly dependent.

*Proof.* If  $\langle x, y \rangle = 0$ , then the inequality is trivial. Otherwise, let  $t \in \mathbb{R}$ . Then

$$\begin{aligned} 0 &\leq \left\langle t \frac{|\langle x, y \rangle|}{\langle x, y \rangle} x + y, t \frac{|\langle x, y \rangle|}{\langle x, y \rangle} x + y \right\rangle \\ &= t^2 \|x\|^2 + 2t \Re \left( \frac{|\langle x, y \rangle|}{\langle x, y \rangle} \langle x, y \rangle \right) + \|y\|^2 = t^2 \|x\|^2 + 2t |\langle x, y \rangle| + \|y\|^2. \end{aligned}$$

Hence

$$4 |\langle x, y \rangle|^2 - 4 \|x\|^2 \|y\|^2 \leq 0 \implies |\langle x, y \rangle| \leq \|x\| \|y\|.$$

Note that if the equality holds, then

$$t^2 \|x\|^2 + 2t |\langle x, y \rangle| + \|y\|^2 = t^2 \|x\|^2 + 2t \|x\| \|y\| + \|y\|^2 = (t \|x\| + \|y\|)^2 = 0$$

by taking  $t = -\|y\| / \|x\|$ . But this implies that

$$t \frac{|\langle x, y \rangle|}{\langle x, y \rangle} x + y = 0$$

and so  $x$  and  $y$  are linearly dependent. Conversely, suppose  $cx = y$ . Then  $|\langle x, y \rangle| = |c| \|x\|^2 = \|x\| \|y\|$ . ■

**Proposition 3.4** (Parallelogram Law)

For all  $x, y \in \mathcal{H}$ ,

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

*Proof.* Note that

$$\begin{aligned}\|x + y\|^2 &= \langle x + y, x + y \rangle = \|x\|^2 + 2\Re(\langle x, y \rangle) + \|y\|^2, \\ \|x - y\|^2 &= \langle x - y, x - y \rangle = \|x\|^2 - 2\Re(\langle x, y \rangle) + \|y\|^2.\end{aligned}$$

Adding the two equations gives

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

**Proposition 3.5**

For all  $x \in \mathcal{H}$ ,

$$\|x\| = \sup_{\|y\|=1} |\langle x, y \rangle|.$$

*Proof.* By the Cauchy-Schwarz inequality,

$$|\langle x, y \rangle| \leq \|x\| \|y\| \implies \left| \left\langle x, \frac{y}{\|y\|} \right\rangle \right| \leq \|x\| \implies \sup_{\|y\|=1} |\langle x, y \rangle| \leq \|x\|.$$

Taking  $y = x/\|x\|$  gives the equality and  $\|y\| = 1$ . ■

**Theorem 3.6** (Completion of Pre-Hilbert Space)

Let  $(X, \langle \cdot, \cdot \rangle)$  be a pre-Hilbert space. Then there exists a Hilbert space  $\mathcal{H}$  such that  $X$  is dense in  $\mathcal{H}$  and  $\langle \cdot, \cdot \rangle_*$  on  $\mathcal{H}$  is an extension of  $\langle \cdot, \cdot \rangle$ .

*Proof.* Define  $\langle x, y \rangle_* = \lim_{n \rightarrow \infty} \langle x_n, y_n \rangle$  for Cauchy sequences  $\{x_n\}, \{y_n\} \subset X$  and  $x, y \in \overline{X}$ . We first check that  $\langle \cdot, \cdot \rangle_*$  is well-defined. Note that

$$\begin{aligned}|\langle x_n, y_n \rangle - \langle x_m, y_m \rangle| &\leq |\langle x_n, y_n \rangle - \langle x_n, y_m \rangle| + |\langle x_n, y_m \rangle - \langle x_m, y_m \rangle| \\ &\leq \|x_n\| \|y_n - y_m\| + \|x_n - x_m\| \|y_m\| \rightarrow 0\end{aligned}$$

as  $n, m \rightarrow \infty$  by the Cauchy-Schwarz inequality. Since  $\mathbb{F}$  is complete, the limit exists. To see that  $\langle \cdot, \cdot \rangle_*$  is independent of the choice of sequences, suppose  $\{x_n^1\}, \{y_n^1\}$  and  $\{x_n^2\}, \{y_n^2\}$  are two pairs of Cauchy sequences converging to  $x$  and  $y$  respectively. Then

$$\begin{aligned}|\langle x_n^1, y_n^1 \rangle - \langle x_n^2, y_n^2 \rangle| &\leq |\langle x_n^1, y_n^1 \rangle - \langle x_n^1, y_n^2 \rangle| + |\langle x_n^1, y_n^2 \rangle - \langle x_n^2, y_n^2 \rangle| \\ &\leq \|x_n^1\| \|y_n^1 - y_n^2\| + \|x_n^1 - x_n^2\| \|y_n^2\| \rightarrow 0.\end{aligned}$$



Hence  $\langle x, y \rangle_*$  is well-defined. We now show that  $\langle \cdot, \cdot \rangle_*$  is indeed an inner product on  $\overline{X}$ . For the linearity in the first argument, let  $x, y, z \in \overline{X}$ ,  $\{x_n\}, \{y_n\}, \{z_n\} \subset X$  be Cauchy sequences converging to  $x, y, z$  respectively and  $c \in \mathbb{F}$ . Then

$$\begin{aligned}\langle cx + y, z \rangle_* &= \lim_{n \rightarrow \infty} \langle cx_n + y_n, z_n \rangle = \lim_{n \rightarrow \infty} c \langle x_n, z_n \rangle + \langle y_n, z_n \rangle \\ &= c \lim_{n \rightarrow \infty} \langle x_n, z_n \rangle + \lim_{n \rightarrow \infty} \langle y_n, z_n \rangle = c \langle x, z \rangle_* + \langle y, z \rangle_*.\end{aligned}$$

For the conjugate symmetry, let  $x, y \in \overline{X}$  and  $\{x_n\}, \{y_n\} \subset X$  be Cauchy sequences converging to  $x$  and  $y$  respectively. Then

$$\overline{\langle x, y \rangle_*} = \overline{\lim_{n \rightarrow \infty} \langle x_n, y_n \rangle} = \lim_{n \rightarrow \infty} \overline{\langle x_n, y_n \rangle} = \lim_{n \rightarrow \infty} \langle y_n, x_n \rangle = \langle y, x \rangle_*.$$

For the positive definiteness, let  $x \in \overline{X}$ ,  $x \neq 0$  and  $\{x_n\} \subset X$  be a Cauchy sequence converging to  $x$ . Then

$$\langle x, x \rangle_* = \lim_{n \rightarrow \infty} \langle x_n, x_n \rangle = \lim_{n \rightarrow \infty} \|x_n\|^2 > 0.$$

Hence  $\langle \cdot, \cdot \rangle_*$  is an inner product on  $\overline{X}$  and induces a norm on  $\overline{X}$ . Lastly, for every  $x, y \in \overline{X}$ , pick  $x_n = x$  and  $y_n = y$  to see that

$$\langle x, y \rangle_* = \lim_{n \rightarrow \infty} \langle x, y \rangle = \langle x, y \rangle,$$

which shows that  $\langle \cdot, \cdot \rangle_*$  is an extension of  $\langle \cdot, \cdot \rangle$ . We conclude that  $\mathcal{H} = \overline{X}$  forms a Hilbert space. ■

### Example

Let  $X = C([0, 1])$  with the inner product

$$\langle f, g \rangle = \int_0^1 f(x)g(x)dx.$$

Then  $X$  is a pre-Hilbert space. To see this, set  $f_n(x) = x^n$ .

$$\|f_m - f_n\|^2 = \int_0^1 (x^m - x^n)^2 dx = \frac{1}{2m+1} + \frac{2}{m+n+1} + \frac{1}{2n+1} \rightarrow 0$$

as  $m, n \rightarrow \infty$ . Hence  $\{f_n\}$  is Cauchy in  $X$ . However,  $f_n$  converges to

$$f(x) = \begin{cases} 0 & x \in [0, 1) \\ 1 & x = 1, \end{cases}$$

which is not in  $X$ . Hence  $X$  is not complete. But by the [proposition 1.35](#),  $X$  is dense in  $\mathcal{L}^2([0, 1])$  and so  $X$  can be completed to a Hilbert space  $\mathcal{H} = \mathcal{L}^2([0, 1])$ , which is complete by Riesz-

*Fischer theorem.*

**Definition 3.7**

A set  $X$  is called **convex** if for all  $x, y \in X$  and  $t \in [0, 1]$ ,  $tx + (1 - t)y \in X$ .

**Theorem 3.8**

Let  $K \subset \mathcal{H}$  be a closed convex set. For  $x \in \mathcal{H}$ , define the distance from  $x$  to  $K$  as

$$d(x, K) = \inf_{y \in K} \|x - y\|.$$

Then there exists a unique  $z \in K$  such that  $d(x, K) = \|x - z\|$ .

*Proof.* Let  $\{y_n\} \subset K$  be a sequence such that  $\|y_n - x\| \rightarrow d(x, K)$ . We claim that  $\{y_n\}$  is Cauchy. Let  $\epsilon > 0$  be given. By the parallelogram law,

$$2(\|x - y_n\|^2 + \|x - y_m\|^2) = \|2x - y_n - y_m\|^2 + \|y_n - y_m\|^2$$

Rearranging gives

$$\begin{aligned} \frac{1}{4} \|y_n - y_m\|^2 &= \frac{1}{2} \|x - y_n\|^2 + \frac{1}{2} \|x - y_m\|^2 - \left\|x - \frac{y_n + y_m}{2}\right\|^2 \\ &\leq \frac{1}{2} (d(x, K) + \epsilon)^2 + \frac{1}{2} (d(x, K) + \epsilon)^2 - d(x, K)^2 \\ &= \epsilon^2 + 2\epsilon d(x, K) \end{aligned}$$

for all  $m, n \geq N$  for some  $N \in \mathbb{N}$ . The inequality follows from the fact that  $(y_n + y_m)/2 \in K$  by the convexity of  $K$ . Since  $\epsilon > 0$  is arbitrary, we conclude that  $\{y_n\}$  is Cauchy. By the completeness of  $\mathcal{H}$ ,  $\{y_n\}$  converges to some  $z \in \mathcal{H}$ . Since  $K$  is closed,  $z \in K$ . To see the uniqueness, suppose  $z_1, z_2 \in K$  are such that  $\|x - z_1\| = \|x - z_2\| = d(x, K)$ . Then by the parallelogram law,

$$\begin{aligned} 4d(x, K)^2 &= 2\|x - z_1\|^2 + 2\|x - z_2\|^2 = \|z_1 - z_2\|^2 + \|2x - z_1 - z_2\|^2 \\ &= \|z_1 - z_2\|^2 + 4\left\|x - \frac{z_1 + z_2}{2}\right\|^2. \end{aligned}$$

Hence

$$\|z_1 - z_2\|^2 = 4d(x, K)^2 - 4\left\|x - \frac{z_1 + z_2}{2}\right\|^2 \leq 4d(x, K)^2 - 4d(x, K)^2 = 0$$

and so  $z_1 = z_2$ . ■

**Definition 3.9**

$Y \subset \mathcal{H}$  is a closed subspace. The **orthogonal complement** of  $Y$ , denoted by  $Y^\perp$ , is defined as

$$Y^\perp = \{x \in \mathcal{H} \mid \langle x, y \rangle = 0 \text{ for all } y \in Y\}.$$

**Proposition 3.10**

$Y \subset \mathcal{H}$  is a closed subspace. Then

(a)  $Y^\perp$  is a closed subspace.

(b)  $\mathcal{H} = Y \oplus Y^\perp$ .

(c)  $(Y^\perp)^\perp = Y$ .

*Proof.* For (a), we first check that  $Y^\perp$  is a subspace. First note that  $0 \in Y^\perp$ . Also, if  $x, z \in Y^\perp$  and  $c \in \mathbb{F}$ , then

$$\langle cx + z, y \rangle = c \langle x, y \rangle + \langle z, y \rangle = 0$$

for all  $y \in Y$ . Hence  $cx + z \in Y^\perp$ . This shows that  $Y^\perp$  is a subspace. To see that  $Y^\perp$  is closed, let  $\{x_n\} \subset Y^\perp$  be a sequence converging to  $x$ . Then for all  $y \in Y$ ,

$$|\langle x, y \rangle - \langle x_n, y \rangle| = |\langle x - x_n, y \rangle| \leq \|x - x_n\| \|y\| \rightarrow 0$$

by the Cauchy-Schwarz inequality. Thus  $\langle \cdot, y \rangle$  is a continuous functional on  $\mathcal{H}$ . Therefore,

$$\langle x, y \rangle = \lim_{n \rightarrow \infty} \langle x_n, y \rangle = 0$$

for all  $y \in Y$  and so  $x \in Y^\perp$ . This shows that  $Y^\perp$  is closed.

For (b), notice that  $Y$  as a closed subspace is convex. By [theorem 3.8](#), for any  $u \in \mathcal{H}$ , there exists a unique  $y \in Y$  such that  $\|u - y\| \leq \|u - y'\|$  for all  $y' \in Y$ . Let  $z = u - y$ . We claim that  $z \in Y^\perp$ . To see this, let  $y' \in Y$  and  $t \in \mathbb{R}$ . Then

$$\begin{aligned} \|z\|^2 &= \|u - y\|^2 \leq \|u - y - ty'\|^2 \\ &= \|u - y\|^2 - 2t\Re(\langle u - y, y' \rangle) + t^2 \|y'\|^2 \\ &= \|z\|^2 - 2t\Re(\langle z, y' \rangle) + t^2 \|y'\|^2. \end{aligned}$$

Rearranging gives

$$2t\Re(\langle z, y' \rangle) - t^2 \|y'\|^2 \leq 0.$$

If  $y' = 0$ , we have  $\langle z, y' \rangle = 0$ ; if  $y' \neq 0$ , then take  $t = \Re(\langle z, y' \rangle / \|y'\|^2)$ . Substituting this back gives

$$0 \geq 2 \frac{(\Re(\langle z, y' \rangle))^2}{\|y'\|^2} - \frac{(\Re(\langle z, y' \rangle))^2}{\|y'\|^2} = \frac{(\Re(\langle z, y' \rangle))^2}{\|y'\|^2}.$$

Hence  $\Re(\langle z, y' \rangle) = 0$  for all  $y' \in Y$ . Similarly, replacing  $t$  with  $it$  gives  $\Im(\langle z, y' \rangle) = 0$  for all  $y' \in Y$ . Therefore,  $\langle z, y' \rangle = 0$  for all  $y' \in Y$  and so  $z \in Y^\perp$ . Since our choice of  $y$  is unique, we can write  $u = y + z$  uniquely for  $y \in Y$  and  $z \in Y^\perp$ . This shows that  $\mathcal{H} = Y \oplus Y^\perp$ .

For (c), note that we can apply (a) and (b) to  $Y^\perp$  and obtain that  $(Y^\perp)^\perp$  is a closed subspace and  $\mathcal{H} = Y \oplus Y^\perp = (Y^\perp)^\perp \oplus Y^\perp$ . It follows that for every  $u \in \mathcal{H}$ , we can write  $u = y + z = x + z$  for  $x \in (Y^\perp)^\perp$ ,  $y \in Y$  and  $z \in Y^\perp$  by the uniqueness of decomposition. This implies that  $y = x$  and hence  $(Y^\perp)^\perp = Y$ . ■

### 3.2. Separability and Orthonormal Basis

#### Definition 3.11

A Hilbert space  $\mathcal{H}$  is said to be **separable** if there exists a countable dense subset in  $\mathcal{H}$ .

#### Definition 3.12

$\{x_\alpha \mid \alpha \in A\} \subset \mathcal{H}$ , where  $A$  is an arbitrary index set. The **linear span** of  $\{x_\alpha \mid \alpha \in A\}$  is defined as

$$\text{span} \{x_\alpha\} = \left\{ \sum_{\alpha \in A} c_\alpha x_\alpha \mid c_\alpha \in \mathbb{F} = \mathbb{R} \text{ or } \mathbb{C} \right\},$$

where the sum is a finite sum.

#### Definition 3.13

$\{x_\alpha \mid \alpha \in A\} \subset \mathcal{H}$ , where  $A$  is an arbitrary index set. The **closed linear span** of  $\{x_\alpha \mid \alpha \in A\}$  is defined as the smallest closed subspace of  $\mathcal{H}$  containing  $\{x_\alpha \mid \alpha \in A\}$ .

#### Proposition 3.14

Let  $Y = \overline{\text{span}(\{x_\alpha\})} \subset \mathcal{H}$  be a closed linear span of  $\{x_\alpha\}$ . Then for any  $x \in \mathcal{H}$ ,  $\langle x, x_\alpha \rangle = 0$  for all  $\alpha \in A$  if and only if  $\langle x, z \rangle = 0$  for all  $z \in Y$ .

*Proof.* Assume first that for  $x \in \mathcal{H}$ ,  $\langle x, x_\alpha \rangle = 0$  for all  $\alpha \in A$ . For each  $z \in Y$ , write  $z = \sum_{\alpha_j \in A} c_{\alpha_j} x_{\alpha_j}$ . Then

$$\langle x, z \rangle = \lim_{M \rightarrow \infty} \left\langle x, \sum_{j=1}^M c_{\alpha_j} x_{\alpha_j} \right\rangle = \lim_{M \rightarrow \infty} \sum_{j=1}^M \overline{c_{\alpha_j}} \langle x, x_{\alpha_j} \rangle = 0.$$

The converse is trivial since  $x_\alpha \in Y$  for all  $\alpha \in A$ . ■

#### Definition 3.15

$\{x_\alpha \mid \alpha \in A\}$  is said to be **orthonormal** if  $\langle x_\alpha, x_\beta \rangle = \delta_{\alpha\beta}$  for all  $\alpha, \beta \in A$ .

#### Definition 3.16

$\{x_\alpha \mid \alpha \in A\}$  forms a **orthonormal basis** of  $\mathcal{H}$  if it is orthonormal and  $\overline{\text{span}(\{x_\alpha\})} = \mathcal{H}$ .

#### Remark

This definition of basis is different from the definition of basis in linear algebra. In linear algebra, one can only express a vector as a finite linear combination of basis vectors; however, in Hilbert space, one can express a vector as a countable linear combination of basis vectors.

#### Lemma 3.17 (Bessel's Inequality)

Let  $\{x_\alpha \mid \alpha \in A\}$  be an orthonormal set in  $\mathcal{H}$ . For any  $x \in \mathcal{H}$ , let  $c_\alpha = \langle x, x_\alpha \rangle$ . Then

- (a) The set  $\{\alpha \mid c_\alpha \neq 0\}$  is at most countable.
- (b)  $\sum_\alpha |c_\alpha|^2 \leq \|x\|^2$ .

*Proof.* We assume that (a) is established and prove (b) first. Let  $J \subset A$  be a countable subset with  $J = \{\alpha_k \mid k \in \mathbb{N}\}$ . For each  $M \in \mathbb{N}$ ,

$$\begin{aligned} 0 &\leq \left\| \sum_{k=1}^M c_{\alpha_k} x_{\alpha_k} - x \right\|^2 = \sum_{k=1}^M |c_{\alpha_k}|^2 - 2\Re \left( \left\langle x, \sum_{k=1}^M c_{\alpha_k} x_{\alpha_k} \right\rangle \right) + \|x\|^2 \\ &= \sum_{k=1}^M |c_{\alpha_k}|^2 - 2\Re \left( \sum_{k=1}^M \overline{c_{\alpha_k}} \langle x, x_{\alpha_k} \rangle \right) + \|x\|^2 = \sum_{k=1}^M |c_{\alpha_k}|^2 - 2 \sum_{k=1}^M |c_{\alpha_k}|^2 + \|x\|^2 \\ &= \|x\|^2 - \sum_{k=1}^M |c_{\alpha_k}|^2 \implies \sum_{k=1}^M |c_{\alpha_k}|^2 \leq \|x\|^2. \end{aligned}$$

Taking  $M \rightarrow \infty$ , we have  $\sum_{k=1}^{\infty} |c_{\alpha_k}|^2 \leq \|x\|^2$ .

Now we turn back to establish (a). For  $m \in \mathbb{N}$ , let  $J_m = \{\alpha \in A \mid |c_{\alpha}| \geq 1/m\}$ . Then  $J_m$  is finite or we can find infinitely many  $\alpha \in J_m$  such that  $|c_{\alpha}| \geq 1/m$ . This implies that  $x \in \mathcal{H}$  and

$$\|x\|^2 = \sum_{\alpha \in A} |c_{\alpha}|^2 \geq \sum_{\alpha \in J_m} |c_{\alpha}|^2 \geq \sum_{\alpha \in J_m} \frac{1}{m^2} = \infty,$$

which is absurd. Thus  $J_m$  is finite for all  $m \in \mathbb{N}$ . Observe that  $\bigcup_{m \in \mathbb{N}} J_m = \{\alpha \in A \mid c_{\alpha} \neq 0\}$ . It follows that as a countable union of finite sets,  $\{\alpha \in A \mid c_{\alpha} \neq 0\}$  is at most countable. (b) follows from (a) and the previous argument. ■

### Remark

*There is a non-separable Hilbert space. Consider an uncountable set  $S$ . Let*

$$\mathcal{H} = \left\{ f : S \rightarrow \mathbb{R} \mid \sum_{s \in S} f(s)^2 < \infty, f(S) \setminus \{0\} \text{ is at most countable} \right\}.$$

*Then  $\mathcal{H}$  is a Hilbert space with the inner product*

$$\langle f, g \rangle = \sum_{s \in S} f(s)g(s).$$

*To see that  $\mathcal{H}$  is a Hilbert space, note that the countable union of countably many non-zero points is countable. Also, it is not separable since the set*

$$\{e_s : S \rightarrow \mathbb{R} \mid e_s(t) = \delta_{st}\}$$

*forms an orthonormal set in  $\mathcal{H}$  and for each  $s \neq r$ ,  $\|e_s - e_r\| = \sqrt{2}$ . This shows that it is nowhere dense in  $\mathcal{H}$ . Thus  $\mathcal{H}$  is not separable.*

### Proposition 3.18

Let  $\{x_\alpha\}$  be an orthonormal set and  $Y$  be the closed linear span of  $\{x_\alpha\}$ . Then

$$Y = \left\{ \sum_j c_j x_{\alpha_j} \mid \sum_j |c_j|^2 < \infty, \alpha_j \in A \right\}.$$

*Proof.* Let  $S = \left\{ \sum_j c_j x_{\alpha_j} \mid \sum_j |c_j|^2 < \infty, \alpha_j \in A \right\}$ . For  $x \in S$ ,  $x = \sum_j c_j x_{\alpha_j}$  with  $\sum_j |c_j|^2 < \infty$ . Then  $z_n = \sum_{j=1}^n c_j x_{\alpha_j} \rightarrow x$  as  $n \rightarrow \infty$ . Each  $z_n \in Y$  and thus  $x \in Y$ . Hence  $S \subset Y$ .

Conversely, we claim that  $S$  is a closed subspace of  $\mathcal{H}$ . Clearly  $0 \in S$ . For  $c \in \mathbb{F}$ ,  $x = \sum_j c_j x_{\alpha_j} \in S$  and  $y = \sum_j d_j x_{\alpha_j} \in S$ , we have

$$cx + y = c \sum_j c_j x_{\alpha_j} + \sum_j d_j x_{\alpha_j} = \sum_j (cc_j + d_j) x_{\alpha_j} \in S,$$

where the summation is over all  $j$  such that either  $c_j \neq 0$  or  $d_j \neq 0$ . To see that  $S$  is closed, let  $z_n \in S$  where  $z_n = \sum_j c_j^n x_{\alpha_j}$  with  $\sum_j |c_j^n|^2 < \infty$  for all  $n \in \mathbb{N}$ . Let  $J_n = \{ \alpha_j^n \mid j \in \mathbb{N} \}$  and  $J = \bigcup_n J_n \subset A$  is at most countable. Consider the transformation  $T : S \rightarrow \ell^2$  defined by  $\sum_j c_j x_{\alpha_j} \mapsto \{c_j\}$ . Such definition is well-defined since if  $\sum_j c_j x_{\alpha_j} = \sum_j d_j x_{\alpha_j}$ , then  $\sum_j (c_j - d_j) x_{\alpha_j} = 0$  and thus  $c_j = d_j$  for all  $j$  since every  $x_{\alpha_j}$  is orthogonal and thus linearly independent. Furthermore,  $T$  is clearly linear. Also, it is isometric since

$$\left\| \sum_j c_j x_{\alpha_j} \right\|^2 = \sum_j |c_j|^2 = \|\{c_j\}\|^2.$$

For  $z_n \in S$ ,  $z_n \rightarrow z$ . Since  $z_n$  is Cauchy and  $T$  is isometric,  $\{c_j^n\}$  is Cauchy in  $\ell^2$  and thus converges to some  $\{c_j\} \in \ell^2$ . Define  $w = \sum_j c_j x_{\alpha_j} \in S$ . It follows that  $Tz_n \rightarrow Tw$ . Hence  $z = w \in S$  by the isometry of  $T$ . Thus  $S$  is closed. It follows that by the definition of  $Y$ ,  $Y \subset S$ . We conclude that  $Y = S$ .  $\blacksquare$

**Lemma 3.19** (Gram-Schmidt)

Suppose  $\{x_\alpha\}$  is an orthonormal set in  $\mathcal{H}$  with  $\overline{\text{span}(\{x_\alpha\})} \neq \mathcal{H}$ . Then there exists  $y \in \mathcal{H}$  such that  $\{x_\alpha\} \cup \{y\}$  is orthonormal.

*Proof.* Pick  $z \in \mathcal{H}$  such that  $z \notin \overline{\text{span}(\{x_\alpha\})}$ . By lemma 3.17, there are at most countably many  $\alpha$  such that  $\langle z, x_\alpha \rangle \neq 0$ . Let  $\alpha_j$  denumerate all  $\alpha$  such that  $\langle z, x_\alpha \rangle \neq 0$ . Set  $\hat{z} = \sum_j \langle z, x_{\alpha_j} \rangle x_{\alpha_j}$ . For each  $x_{\alpha_k}$ ,

$$\begin{aligned} \langle z - \hat{z}, x_{\alpha_k} \rangle &= \lim_{m \rightarrow \infty} \left\langle z - \sum_{j=1}^m \langle z, x_{\alpha_j} \rangle x_{\alpha_j}, x_{\alpha_k} \right\rangle \\ &= \lim_{m \rightarrow \infty} \langle z, x_{\alpha_k} \rangle - \sum_{j=1}^m \langle z, x_{\alpha_j} \rangle \delta_{jk} \\ &= \langle z, x_{\alpha_k} \rangle - \langle z, x_{\alpha_k} \rangle = 0. \end{aligned}$$

And for those  $x_\alpha$  such that  $\langle z, x_\alpha \rangle = 0$ ,  $\langle z - \hat{z}, x_\alpha \rangle = \langle z, x_\alpha \rangle - \langle \hat{z}, x_\alpha \rangle = 0$  since  $x_{\alpha_j}$  and  $x_\alpha$  are orthogonal. Now set  $y = (z - \hat{z})/\|z - \hat{z}\|$ . Then  $\{x_\alpha\} \cup \{y\}$  forms an orthonormal set. ■

### Theorem 3.20

*Every Hilbert space has an orthonormal basis.*

*Proof.* We plan to use Zorn's lemma. Denote the space consisting of all orthonormal sets in  $\mathcal{H}$  by  $\mathcal{O}$ . Define a partial order as the inclusion of sets. Let  $C \subset \mathcal{O}$  be a chain. We claim that  $B = \bigcup_{\{x_\alpha\} \in C} \{x_\alpha\}$  is an upper bound of  $C$ . By construction we have  $\{x_\alpha\} \subset B$  for all  $\{x_\alpha\} \in C$ . We need to show that  $B \in \mathcal{O}$ . For distinct  $x_\alpha, x_\beta \in B$ , they belong to a common set  $C \in C \subset \mathcal{O}$ . Hence  $C$  is orthonormal and  $\langle x_\alpha, x_\beta \rangle = 0$ . Also, it is clear that for every  $x_\alpha \in B$ ,  $x_\alpha$  belongs to some  $C \in C$  and thus  $\|x_\alpha\| = 1$ . It follows that  $B$  is also orthonormal. By Zorn's lemma, there exists a maximal element in  $\mathcal{O}$ , say  $B$ , such that if  $C \in \mathcal{O}$  and  $B \subset C$ , then  $B = C$ . We claim that  $B$  is an orthonormal basis. It suffices to check that  $\overline{\text{span}(B)} = \mathcal{H}$ . Suppose not, then by lemma 3.19, there exists  $y \in \mathcal{H}$  such that  $\{x_\alpha\} \cup \{y\}$  forms an orthonormal set. This contradicts the maximality of  $B$ . We conclude that  $B$  is an orthonormal basis. ■

### Theorem 3.21

*Let  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ . Suppose that  $\mathcal{H}$  is a Hilbert space. Then  $\mathcal{H}$  is separable if and only if  $\mathcal{H}$  has a countable orthonormal basis.*

*Proof.* Suppose that  $\mathcal{H}$  has a countable orthonormal basis  $\{x_n\}$ . Then consider the sets

$$A_n = \left\{ \sum_{j=1}^n c_j x_j \mid c_j \in S \right\},$$

where  $S = \mathbb{Q}$  if  $\mathbb{F} = \mathbb{R}$  and  $S = \mathbb{Q} + \mathbb{Q}i$  if  $\mathbb{F} = \mathbb{C}$ . Since  $S$  is countable, each  $A_n$  being a finite union of countable sets is countable. Put  $A = \bigcup_n A_n$  and let  $\epsilon > 0$  be given. Since  $A$  is a countable union of countable sets, it is also countable. For every  $x \in \mathcal{H}$ , we can write  $x = \sum_j \langle x, x_j \rangle x_j$  with

$$\left\| \sum_{j=N+1}^{\infty} \langle x, x_j \rangle x_j \right\| < \frac{\epsilon}{2}$$

for some  $N \in \mathbb{N}$ . Since  $S$  is dense in  $\mathbb{F}$ , we can pick some  $c_j \in S$  with  $|c_j - \langle x, x_j \rangle| < \epsilon/2^{j+1}$ . Then

$$\begin{aligned} \left\| x - \sum_{j=1}^N c_j x_j \right\| &= \left\| \sum_{j=1}^{\infty} \langle x, x_j \rangle x_j - \sum_{j=1}^N \langle x, x_j \rangle x_j + \sum_{j=1}^N \langle x, x_j \rangle x_j - \sum_{j=1}^N c_j x_j \right\| \\ &\leq \left\| \sum_{j=1}^{\infty} \langle x, x_j \rangle x_j - \sum_{j=1}^N \langle x, x_j \rangle x_j \right\| + \left\| \sum_{j=1}^N \langle x, x_j \rangle x_j - \sum_{j=1}^N c_j x_j \right\| \\ &\leq \left\| \sum_{j=N+1}^{\infty} \langle x, x_j \rangle x_j \right\| + \sum_{j=1}^N |\langle x, x_j \rangle - c_j| \|x_j\| \leq \frac{\epsilon}{2} + \sum_{j=1}^N \frac{\epsilon}{2^{j+1}} \leq \epsilon. \end{aligned}$$

It follows that  $A$  is dense in  $\mathcal{H}$  and hence  $\mathcal{H}$  is separable.

Conversely, suppose that  $\mathcal{H}$  is separable. Let  $S \subset \mathcal{H}$  be a countable subset. Assume that every orthonormal basis of  $\mathcal{H}$  is uncountable. Denote an orthonormal basis of  $\mathcal{H}$  by  $\{x_\alpha\}$ . For each distinct  $x_\alpha, x_\beta \in S$ ,  $\|x_\alpha - x_\beta\| = \sqrt{2}$ . Consider the open balls  $B_{1/2}(x_\alpha)$ . They are clearly disjoint since if  $y$  lies in two such balls, then  $\sqrt{2} = \|x_\alpha - x_\beta\| \leq \|x_\alpha - y\| + \|y - x_\beta\| < 1$ , which is absurd. Now since  $S$  is dense in  $\mathcal{H}$ , for each  $\alpha$  we can find some  $s_\alpha \in S$  such that  $s_\alpha \in B_{1/2}(x_\alpha)$ . It follows that each  $s_\alpha$  is distinct and thus  $S$  is uncountable. This contradicts to our assumption that  $S$  is countable. Thus  $\mathcal{H}$  must have a countable orthonormal basis. ■

**Proposition 3.22**

Let  $\mathcal{H}$  be a Hilbert space and  $\{x_\alpha \mid \alpha \in A\}, \{y_\beta \mid \beta \in B\}$  be two orthonormal bases in  $\mathcal{H}$ . Then  $\text{card}(A) = \text{card}(B)$ .

*Proof.* Fixed an  $\alpha \in A$ ,  $B_\alpha = \{\beta \in B \mid \langle y_\beta, x_\alpha \rangle \neq 0\}$  is at most countable by [lemma 3.17](#) and  $B_\alpha \subset B$ . We claim that  $B \subset \bigcup_{\alpha \in A} B_\alpha$ . Take  $\beta \in B$ , we can write  $y_\beta = \sum_k \langle y_\beta, x_{\alpha_k} \rangle x_{\alpha_k}$  with at least one  $\langle y_\beta, x_{\alpha_k} \rangle \neq 0$ . Hence  $\beta \in B_{\alpha_k}$  for some  $\alpha_k \in A$ . It follows that  $B \subset \bigcup_{\alpha \in A} B_\alpha$  and hence  $\text{card}(B) \leq \text{card}(A)$ . By symmetry, we have  $\text{card}(A) \leq \text{card}(B)$  and thus  $\text{card}(A) = \text{card}(B)$ . ■

**Remark**

If  $\mathcal{H}$  is separable, then  $\mathcal{H}$  has a countable orthonormal basis and hence every orthonormal basis of  $\mathcal{H}$  is countable.

**Proposition 3.23** (Parseval's Identity)

Let  $\{x_\alpha\}$  be an orthonormal basis of  $\mathcal{H}$ . Then

$$\|x\|^2 = \sum_j |\langle x, x_{\alpha_j} \rangle|^2.$$

*Proof.* Let  $x \in \mathcal{H}$ . Write  $x = \sum_j c_j x_{\alpha_j}$  with  $\sum_j |c_j|^2 < \infty$ . Then

$$\langle x, x_{\alpha_k} \rangle = \lim_{M \rightarrow \infty} \left\langle \sum_{j=1}^M c_j x_{\alpha_j}, x_{\alpha_k} \right\rangle = \lim_{M \rightarrow \infty} \sum_{j=1}^M c_j \langle x_{\alpha_j}, x_{\alpha_k} \rangle = c_k.$$

It follows that

$$\|x\|^2 = \lim_{M \rightarrow \infty} \left\langle \sum_{j=1}^M c_j x_{\alpha_j}, \sum_{j=1}^M c_j x_{\alpha_j} \right\rangle = \lim_{M \rightarrow \infty} \sum_{j=1}^M |c_j|^2 = \sum_j |c_j|^2 = \sum_j |\langle x, x_{\alpha_j} \rangle|^2.$$

■

### 3.3. Riesz Representation and Bilinear Form

**Proposition 3.24**

$\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .  $T : \mathcal{H} \rightarrow \mathbb{F}$  is a nonzero bounded linear functional.



- (a)  $\mathcal{H} = \text{span}(\{w\}) \oplus \ker(T)$  for  $w \notin \ker(T)$ .
- (b) If  $S, T$  are bounded linear functionals and  $\ker(S) = \ker(T)$ , then there exists  $c \in \mathbb{F}$  such that  $S = cT$ .
- (c)  $\ker(T)$  is closed.

*Proof.* For (a), since  $T$  is nonzero, there is some  $w$  such that  $Tw \neq 0$ . For  $x \in \mathcal{H}$ , set  $\alpha = Tx/Tw$  and  $u = x - \alpha w$ . Then  $x = \alpha w + u$  and

$$Tu = Tx - \frac{Tx}{Tw}Tw = 0.$$

Hence  $u \in \ker(T)$ . Also, if  $v \in \text{span}(\{w\}) \cap \ker(T)$ ,  $v = cw$  and  $Tv = 0$ . Then  $cTw = Tv = 0$ ;  $c = 0$  and thus  $v = 0$ . Therefore,  $\text{span}(\{w\}) \cap \ker(T) = \{0\}$  and  $\mathcal{H} = \text{span}(\{w\}) \oplus \ker(T)$ .

To see (b), note that if  $S = 0$ ,  $\mathcal{H} = \ker(S) = \ker(T)$ . Thus  $T = 0$ . If  $S \neq 0$ , by (a) we can write  $\mathcal{H} = \text{span}(\{w\}) \oplus \ker(S) = \text{span}(\{w\}) \oplus \ker(T)$ . Then for every  $x \in \mathcal{H}$ ,  $x = \alpha w + u$  for some  $\alpha \in \mathbb{F}$  and  $u \in \ker(T) = \ker(S)$ . Then  $Tw \neq 0$  and

$$Sx = S(\alpha w + u) = \alpha Sw = \alpha Tw \frac{Sw}{Tw} = \frac{Sw}{Tw} T(\alpha w + u) = \frac{Sw}{Tw} Tx.$$

Taking  $c = Sw/Tw$  gives  $S = cT$ .

For (c), let  $x_n \in \ker(T)$  be a sequence such that  $x_n \rightarrow x \in \mathcal{H}$ . Since  $T$  is continuous,

$$Tx = \lim_{n \rightarrow \infty} Tx_n = 0.$$

Hence  $x \in \ker(T)$  and  $\ker(T)$  is closed. ■

### Theorem 3.25 (Riesz Representation on $\mathcal{H}$ )

$\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .  $T : \mathcal{H} \rightarrow \mathbb{F}$  is a bounded linear functional. Then there exists a unique  $x^* \in \mathcal{H}$  such that  $Ty = \langle y, x^* \rangle$  for all  $y \in \mathcal{H}$ .

*Proof.* If  $T = 0$ , pick  $x^* = 0$  then  $Ty = 0 = \langle y, 0 \rangle$ . If  $T \neq 0$ , there is some  $w \in \mathcal{H}$  such that  $Tw \neq 0$ . By [proposition 3.24](#), we can write  $\mathcal{H} = \text{span}(\{w\}) \oplus \ker(T)$  with  $\ker(T)$  closed. Also,  $\mathcal{H} = \ker(T) \oplus \ker(T)^\perp$  by [proposition 3.10](#). We claim that  $\ker(T)^\perp = \text{span}(\{w\})$ . First note that  $\ker(T)^\perp \neq \{0\}$  or we would have  $\mathcal{H} = \ker(T)$  and  $T = 0$ , contradicting to our assumption. Now if  $z_1, z_2 \in \ker(T)^\perp$ , write  $z_1 = \alpha_1 w + u_1$  and  $z_2 = \alpha_2 w + u_2$  for some  $\alpha_1, \alpha_2 \in \mathbb{F}$  and  $u_1, u_2 \in \ker(T)$ . Then  $\alpha_2 z_1 - \alpha_1 z_2 = \alpha_2 u_1 - \alpha_1 u_2 \in \ker(T)$  and  $\alpha_2 z_1 - \alpha_1 z_2 \in \ker(T)^\perp$ . Hence  $\alpha_2 z_1 - \alpha_1 z_2 = 0$  and  $z_1, z_2$  are linearly dependent. Now define  $S : \mathcal{H} \rightarrow \mathbb{F}$  by  $Sx = \langle x, w \rangle$ . Then  $S$  is a bounded linear functional and  $\ker(S) = \{x \in \mathcal{H} \mid \langle x, w \rangle = 0\} = (\ker(T)^\perp)^\perp = \ker(T)$  by [proposition 3.10](#). Applying (b) of [proposition 3.24](#) gives  $cS = T$  for some  $c \in \mathbb{F}$ . Then  $Tx = cSx = c \langle x, w \rangle = \langle x, \bar{c}w \rangle$ . Set  $x^* = \bar{c}w$  proves the existence of  $x^*$ .

To see uniqueness, suppose  $x_1^*, x_2^* \in \mathcal{H}$  are such that  $Ty = \langle y, x_1^* \rangle = \langle y, x_2^* \rangle$  for all  $y \in \mathcal{H}$ . Then  $\langle y, x_1^* - x_2^* \rangle = 0$  for all  $y \in \mathcal{H}$ . Hence  $x_1^* - x_2^* = 0$  and  $x_1^* = x_2^*$ . Such  $x^*$  is unique. ■

**Definition 3.26**

Let  $X, Y$  be vector spaces.  $T : X \rightarrow Y$  is called **skew-linear** if  $T(cx + y) = \bar{c}Tx + Ty$  for all  $x, y \in X$  and  $c \in \mathbb{F}$ .

**Definition 3.27**

$\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is called a **bilinear form** if

- (a)  $B(\cdot, x)$  is linear for all  $x \in \mathcal{H}$ .
- (b)  $B(x, \cdot)$  is skew-linear for all  $x \in \mathcal{H}$ .

**Definition 3.28**

A bilinear form  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is called **bounded** if there exists  $C < \infty$  such that  $|B(x, y)| \leq C \|x\| \|y\|$  for all  $x, y \in \mathcal{H}$ .

**Definition 3.29**

A bilinear form  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is called **coercive** if there exists  $\delta > 0$  such that  $B(x, x) \geq \delta \|x\|^2$  for all  $x \in \mathcal{H}$ .

**Theorem 3.30 (Lax-Milgram I)**

Let  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ .  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is a bounded coercive bilinear form. Then for every  $L \in \mathcal{H}'$ , there exists  $x \in \mathcal{H}$  such that  $Ly = B(y, x)$  for all  $y \in \mathcal{H}$ .

*Proof.* Fixed  $x \in \mathcal{H}$ . Then  $B(\cdot, x)$  is a bounded linear functional defined on  $\mathcal{H}$ . By Riesz representation, there exists a unique  $x^* \in \mathcal{H}$  such that  $B(y, x) = \langle y, x^* \rangle$  for all  $y \in \mathcal{H}$ . Define  $T : \mathcal{H} \rightarrow \mathcal{H}$  by  $Tx = x^*$ . Such definition is well-defined because  $x^*$  is unique. We claim that  $T$  is bounded and linear. For linearity, let  $x, y, z \in \mathcal{H}$  and  $c \in \mathbb{F}$ . Then

$$\langle y, T(cx + z) \rangle = B(y, cx + z) = \bar{c}B(y, x) + B(y, z) = \bar{c} \langle y, Tx \rangle + \langle y, Tz \rangle = \langle y, cTx + Tz \rangle.$$

Hence  $T(cx + z) = cTx + Tz$ . For boundedness, by [proposition 3.5](#),

$$\|Tx\| = \sup_{\|y\|=1} |\langle y, Tx \rangle| = \sup_{\|y\|=1} |B(y, x)| \leq \sup_{\|y\|=1} C \|x\| \|y\| = C \|x\|.$$

Hence  $T$  is bounded.

Next, let  $A = T(\mathcal{H})$ . We claim that  $A$  is closed. Let  $y_n \rightarrow y$  and  $y_n \in A$ . By the boundedness of  $T$  we have  $\|Tx\| \leq C \|x\|$ . Also, by the coerciveness and [proposition 3.5](#),

$$\delta \|x\|^2 \leq B(x, x) \leq |B(x, x)| = |\langle x, Tx \rangle| = \|x\| \left| \left\langle \frac{x}{\|x\|}, Tx \right\rangle \right| \leq \|x\| \sup_{\|y\|=1} |\langle y, Tx \rangle| = \|x\| \|Tx\|.$$

So  $\|x\| \leq \frac{1}{\delta} \|Tx\|$ . Then we see that the norms  $\|\cdot\|$  and  $\|T(\cdot)\|$  are equivalent. For  $y_n \in A$ , we can find  $x_n \in \mathcal{H}$  such that  $Tx_n = y_n$ . Since  $y_n$  is Cauchy,  $x_n$  is also Cauchy by the equivalence of norms. By the completeness of  $\mathcal{H}$ ,  $x_n \rightarrow x \in \mathcal{H}$ . Then the boundedness of  $T$  implies the continuity and  $Tx_n \rightarrow Tx$ . It follows that  $y = Tx$  by the uniqueness of the limit. Hence  $y \in A$  and  $A$  is closed.

Finally, we claim that  $A = \mathcal{H}$ . Assume not. Then because  $A$  is closed,  $\mathcal{H} = A \oplus A^\perp$  with  $A^\perp \neq \{0\}$  by **proposition 3.10**. There is some  $z \in A$  such that  $\langle z, y \rangle = 0$  for all  $y \in A^\perp$ . This implies  $B(z, x) = \langle z, Tx \rangle = 0$  for all  $x \in \mathcal{H}$ . Taking  $x = z$  gives  $0 = B(z, z) \geq \delta \|z\|^2$  by the coerciveness of  $B$ . Hence  $z = 0$ ,  $A^\perp = \{0\}$ , and  $A = \mathcal{H}$ , a contradiction. We conclude that  $A = \mathcal{H}$ .

For any bounded linear functional  $L$ , there is  $x^* \in \mathcal{H}$  such that  $Ly = \langle y, x^* \rangle$  for all  $y \in \mathcal{H}$  by Riesz representation. Then there is  $x \in \mathcal{H}$  such that  $Tx = x^*$ . Then for all  $y \in \mathcal{H}$ ,

$$Ly = \langle y, x^* \rangle = \langle y, Tx \rangle = B(y, x).$$

This completes the proof. ■

### Remark

*Lax-Milgram theorem ensures the existence of weak solutions to linear PDEs. For example, consider the Poisson equation*

$$-\Delta u = f \quad \text{on } \Omega \subset \mathbb{R}^d, \quad u|_{\partial\Omega} = 0$$

for  $f \in \mathcal{L}^2(\Omega)$ .  $\Delta = \sum_{i=1}^d \partial_i^2$  is the Laplacian. Then for all  $\varphi \in C_c^\infty$ ,

$$L(\varphi) = B(u, \varphi) = \int_{\Omega} \nabla u \cdot \nabla \varphi = \int_{\Omega} f \varphi.$$

### Definition 3.31

A bilinear form  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is called **symmetric** if  $B(x, y) = B(y, x)$  for all  $x, y \in \mathcal{H}$ .

### Theorem 3.32 (Lax-Milgram II)

Let  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ . Suppose that  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is a bounded symmetric coercive bilinear form and  $L \in \mathcal{H}'$  is a bounded linear functional. Then

$$\inf_{x \in \mathcal{H}} \frac{1}{2} B(x, x) - Lx$$

is attained at a unique  $x \in \mathcal{H}$ .

*Proof.* Set  $F(x) = \frac{1}{2} B(x, x) - Lx$  and  $\alpha = \inf_{x \in \mathcal{H}} F(x)$ . We check that  $\alpha$  is finite. Notice that for all  $x \in \mathcal{H}$ ,

$$-M < \frac{1}{2} \delta \|x\|^2 - \|L\| \|x\| \leq \frac{1}{2} B(x, x) - |Lx| \leq F(x) \leq \left| \frac{1}{2} B(x, x) - Lx \right| \leq \frac{1}{2} C \|x\|^2 + \|L\| \|x\| < \infty$$

for some finite  $M, C > 0$  and  $\delta > 0$ . The first inequality is due to  $\delta > 0$  and the quadratic function is bounded below. Taking infimum of  $F(x)$  gives that  $\alpha$  is finite.

Now by definition we can find  $u_n \in \mathcal{H}$  such that  $F(u_n) \rightarrow \alpha$ . We claim that  $u_n$  is Cauchy.

For all  $n, m \in \mathbb{N}$ , we have

$$\begin{aligned}
F(u_m) + F(u_n) &= \frac{1}{2}B(u_m, u_m) - Lu_m + \frac{1}{2}B(u_n, u_n) - Lu_n \\
&= 2\left(B\left(\frac{u_m}{2}, \frac{u_m}{2}\right) + B\left(\frac{u_n}{2}, \frac{u_n}{2}\right)\right) - 2L\left(\frac{u_m + u_n}{2}\right) \\
&= 2\left[\frac{1}{2}\left(B\left(\frac{u_m + u_n}{2}, \frac{u_m + u_n}{2}\right) + B\left(\frac{u_m - u_n}{2}, \frac{u_m - u_n}{2}\right)\right) - L\left(\frac{u_m + u_n}{2}\right)\right] \\
&= 2F\left(\frac{u_m + u_n}{2}\right) + B\left(\frac{u_m - u_n}{2}, \frac{u_m - u_n}{2}\right) \geq 2\alpha + \frac{\delta}{4} \|u_m - u_n\|^2.
\end{aligned}$$

For arbitrary  $\epsilon > 0$ , there is  $N \in \mathbb{N}$  such that for all  $m, n \geq N$ ,  $F(u_m), F(u_n) \leq \alpha + \epsilon$ . Then

$$2\alpha + \frac{\delta}{4} \|u_m - u_n\|^2 \leq F(u_m) + F(u_n) \leq 2\alpha + 2\epsilon \implies \|u_m - u_n\|^2 \leq \frac{8\epsilon}{\delta}.$$

Since  $\epsilon$  is arbitrary, we obtain that  $u_n$  is Cauchy. By the completeness,  $u_n \rightarrow u$  for some  $u \in \mathcal{H}$ . We check that  $u$  is the minimizer of  $F$ , i.e.  $F(u) = \alpha$ . Observe that if for  $u_n \rightarrow u$  and  $v_n \rightarrow v$ , we have  $B(u_n, v_n) \rightarrow B(u, v)$  and  $Lu_n \rightarrow Lu$ . Indeed,  $Lu_n \rightarrow u$  by the boundedness and hence the continuity of  $L$ . Also,

$$|B(u_n, v_n) - B(u, v)| \leq |B(u_n - u, v_n)| + |B(u, v_n - v)| \leq C \|u_n - u\| \|v_n\| + C \|u\| \|v_n - v\| \rightarrow 0$$

since  $u_n \rightarrow u$  as a Cauchy sequence must be bounded. This implies

$$F(u_n) = \frac{1}{2}B(u_n, u_n) - Lu_n \rightarrow \frac{1}{2}B(u, u) - Lu = F(u).$$

The uniqueness of the limit of  $u_n$  ensures that the minimizer is unique. ■

### Theorem 3.33

Let  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ . Suppose that  $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{F}$  is a bounded symmetric coercive bilinear form and  $L \in \mathcal{H}'$  is a bounded linear functional.  $x_0 \in \mathcal{H}$  is a minimizer of  $F(x) = \frac{1}{2}B(x, x) - Lx$  if and only if  $B(x_0, y) = L(y)$  for all  $y \in \mathcal{H}$ .

*Proof.* Suppose that  $x_0$  is a solution to  $B(x_0, y) = L(y)$  for all  $y \in \mathcal{H}$ . Then for all  $x \in \mathcal{H}$ ,

$$\begin{aligned}
F(x_0 + x) - F(x_0) &= \frac{1}{2}B(x_0 + x, x_0 + x) - L(x_0 + x) - \frac{1}{2}B(x_0, x_0) + L(x_0) \\
&= B(x_0, x) - L(x) + \frac{1}{2}B(x, x) \geq \frac{\delta}{2} \|x\|^2 \geq 0
\end{aligned}$$

by the coerciveness of  $B$ . Hence  $F(x_0) \leq F(x_0 + x)$  for all  $x \in \mathcal{H}$  and  $x_0$  is a minimizer.

Conversely, suppose that  $x_0$  minimizes  $F$ . For all  $t \in \mathbb{R}$  and  $y \in \mathcal{H}$ , consider the function

$\phi(t) = F(x_0 + ty)$ . Then since  $x_0$  minimizes  $F$ ,  $\phi'(t)|_{t=0} = 0$ . We compute that

$$\begin{aligned}\phi(t) &= F(x_0 + ty) = \frac{1}{2}B(x_0 + ty, x_0 + ty) - L(x_0 + ty) \\ &= \frac{1}{2}B(x_0, x_0) + tB(x_0, y) + \frac{1}{2}t^2B(y, y) - L(x_0) - tL(y).\end{aligned}$$

Differentiating gives

$$0 = \phi'(t)|_{t=0} = B(x_0, y) - L(y) + tB(y, y)|_{t=0} = B(x_0, y) - L(y)$$

for each given  $y \in \mathcal{H}$ . Hence  $x_0$  satisfies  $B(x_0, y) = L(y)$  for all  $y \in \mathcal{H}$ . ■

### 3.4. Symmetric and Compact Operators

#### Definition 3.34

$\mathcal{D}(A) \subset \mathcal{H}$  is dense in  $\mathcal{H}$ . A linear operator  $A : \mathcal{D}(A) \rightarrow \mathcal{H}$  is said to be **symmetric** if  $\langle Ax, y \rangle = \langle x, Ay \rangle$  for all  $x, y \in \mathcal{D}(A)$ .

#### Remark

Note that the domain  $\mathcal{D}(A)$  is dense in  $\mathcal{H}$ . It follows that by density, the domain can often be extended to  $\mathcal{H}$ . For simplicity, we consider the domain to be  $\mathcal{H}$ , but the domain can be any dense subset of  $\mathcal{H}$ .

#### Definition 3.35

$\lambda \in \mathbb{F}$  is an **eigenvalue** of a linear operator  $A : \mathcal{H} \rightarrow \mathcal{H}$  if there exists a non-zero vector  $x \in \mathcal{H}$  such that  $Ax = \lambda x$ . The vector  $x$  is called the **eigenvector** corresponding to the eigenvalue  $\lambda$ .

#### Proposition 3.36

Let  $A : \mathcal{H} \rightarrow \mathcal{H}$  be a symmetric operator. The followings are true.

- (a)  $\langle Ax, x \rangle \in \mathbb{R}$  for all  $x \in \mathcal{H}$ .
- (b) If  $\lambda \in \mathbb{F}$  is an eigenvalue of  $A$ , then  $\lambda \in \mathbb{R}$ .
- (c) If  $\lambda_1, \lambda_2 \in \mathbb{F}$  are two distinct eigenvalues with respect to eigenvectors  $x_1, x_2 \in \mathcal{H}$ , then  $\langle x_1, x_2 \rangle = 0$ .
- (d) Suppose  $\{x_\alpha\}$  is an orthonormal basis of  $\mathcal{H}$  with the property that each  $x_\alpha$  is an eigenvector of  $A$  corresponding to the eigenvalue  $\lambda_\alpha$ . Then if  $\mu \in \mathbb{F}$  is also an eigenvalue of  $A$ , then  $\mu = \lambda_\alpha$  for some  $\alpha$ .

*Proof.* For (a),  $\langle Ax, x \rangle = \langle x, Ax \rangle = \overline{\langle Ax, x \rangle}$ . Then  $\Im(\langle Ax, x \rangle) = 0$  and  $\langle Ax, x \rangle \in \mathbb{R}$ .

For (b), let  $x \in \mathcal{H}$  be the corresponding eigenvector to  $\lambda$ . Then

$$\langle Ax, x \rangle = \langle \lambda x, x \rangle = \lambda \langle x, x \rangle = \lambda \|x\|^2 \implies \lambda = \frac{\langle Ax, x \rangle}{\|x\|^2} \in \mathbb{R}.$$

For (c), by symmetry and (b), we have

$$\lambda_1 \langle x_1, x_2 \rangle = \langle Ax_1, x_2 \rangle = \langle x_1, Ax_2 \rangle = \overline{\lambda_2} \langle x_1, x_2 \rangle = \lambda_2 \langle x_1, x_2 \rangle.$$

Since  $\lambda_1 \neq \lambda_2$ ,  $\langle x_1, x_2 \rangle = 0$ .

For (d), let  $\mu \in \mathbb{F}$  be an eigenvalue of  $A$  with eigenvector  $y \in \mathcal{H}$ ,  $y \neq 0$ . We claim that  $\mu = \lambda_\alpha$  for some  $\alpha$ . Suppose not. Then write  $y = \sum_j c_j x_{\alpha_j}$ , where  $c_j \in \mathbb{F}$ . We see that

$$\|y\|^2 = \lim_{M \rightarrow \infty} \left\langle y, \sum_{j=1}^M c_j x_{\alpha_j} \right\rangle = \lim_{M \rightarrow \infty} \overline{c_j} \langle y, x_{\alpha_j} \rangle = 0$$

by (c), but this is a contradiction since  $y \neq 0$ . Thus  $\mu = \lambda_\alpha$  for some  $\alpha$ . ■

### Definition 3.37

A linear operator  $A : \mathcal{H} \rightarrow \mathcal{H}$  is called **bounded** if

$$\|A\| = \sup_{\|x\|=1} \|Ax\| < \infty.$$

### Proposition 3.38

$A : \mathcal{H} \rightarrow \mathcal{H}$  is a symmetric bounded linear operator. Then

$$\|A\| = \sup_{\|x\|=1} |\langle Ax, x \rangle|.$$

*Proof.* Assume  $\|x\| = 1$ . By Cauchy-Schwarz inequality,  $|\langle Ax, x \rangle| \leq \|Ax\| \|x\| = \|Ax\|$ . Taking supremum,

$$\sup_{\|x\|=1} |\langle Ax, x \rangle| \leq \sup_{\|x\|=1} \|Ax\| = \|A\|.$$

To see the reverse inequality, note that  $\|Ax\|^2 = \langle Ax, Ax \rangle = \langle A^2x, x \rangle$ . For any nonzero  $\lambda \in \mathbb{R}$ , define  $x^+ = \lambda x + \frac{1}{\lambda} Ax$  and  $x^- = \lambda x - \frac{1}{\lambda} Ax$ . Then  $x = \frac{1}{2\lambda}(x^+ + x^-)$  and  $Ax = \frac{\lambda}{2}(x^+ - x^-)$ . Now,

$$\begin{aligned} \langle A^2x, x \rangle &= \left\langle A \left( \frac{\lambda}{2}(x^+ - x^-) \right), \frac{1}{2\lambda}(x^+ + x^-) \right\rangle \\ &= \frac{1}{4} \langle Ax^+ - Ax^-, x^+ + x^- \rangle \\ &= \frac{1}{4} (\langle Ax^+, x^+ \rangle + \langle Ax^+, x^- \rangle - \langle Ax^-, x^+ \rangle - \langle Ax^-, x^- \rangle) \end{aligned}$$

Notice that  $\langle A^2x, x \rangle$ ,  $\langle Ax^+, x^+ \rangle$  and  $\langle Ax^-, x^- \rangle$  are real numbers by [proposition 3.36](#); hence  $\Im(\langle Ax^+, x^- \rangle - \langle Ax^-, x^+ \rangle) = 0$ . Also,  $\langle Ax^-, x^+ \rangle = \overline{\langle x^+, Ax^- \rangle} = \overline{\langle Ax^+, x^- \rangle}$ . We have  $\Im(\langle Ax^+, x^- \rangle) =$

0 and  $\langle Ax^+, x^- \rangle - \langle Ax^-, x^+ \rangle = 0$ . Thus, letting  $C = \sup_{\|x\|=1} |\langle Ax, x \rangle|$ ,

$$\begin{aligned} \langle A^2x, x \rangle &= \frac{1}{4} (\langle Ax^+, x^+ \rangle + \langle Ax^-, x^- \rangle) \\ &\leq \frac{1}{4} C (\|x^+\|^2 + \|x^-\|^2) \\ &= \frac{1}{4} C (\langle x^+, x^+ \rangle + \langle x^-, x^- \rangle) \\ &= \frac{1}{4} C \left( 2\lambda^2 \|x\|^2 + \frac{2}{\lambda^2} \|Ax\|^2 \right) = \frac{1}{2} C \left( \lambda^2 \|x\|^2 + \frac{1}{\lambda^2} \|Ax\|^2 \right). \end{aligned}$$

Notice that for  $a, b \in \mathbb{R}$ ,  $(a - b)^2 \geq 0$  and thus  $a^2 + b^2 \geq 2ab$ . Hence

$$\lambda^2 \|x\|^2 + \frac{1}{\lambda^2} \|Ax\|^2 \geq 2\lambda \|x\| \frac{1}{\lambda} \|Ax\| = 2 \|Ax\| \|x\|.$$

We see that

$$\|Ax\|^2 = \langle A^2x, x \rangle \leq \frac{1}{2} C \inf_{\lambda \neq 0} \lambda^2 \|x\|^2 + \frac{1}{\lambda^2} \|Ax\|^2 \leq C \|Ax\| \|x\|.$$

Clearly if  $Ax = 0$  the inequality holds. Suppose  $\|Ax\| \neq 0$ . Then deviding both sides by  $\|Ax\|$  and taking supremum gives

$$\|A\| = \sup_{\|x\|=1} \|Ax\| \leq C \sup_{\|x\|=1} \|x\| = C = \sup_{\|x\|=1} |\langle Ax, x \rangle|.$$

We conclude that  $\|A\| = \sup_{\|x\|=1} |\langle Ax, x \rangle|$ . ■

### Definition 3.39

$X$  and  $Y$  are normed spaces.  $M \subset X$ .  $A : M \rightarrow Y$  is an operator. We say that  $A$  is **compact** if  $A$  is continuous and for every bounded sequence  $x_n \in M$ , the sequence  $Ax_n \in Y$  has a convergent subsequence.

### Remark

A compact operator  $A$  transfers bounded sets in  $X$  to relatively compact sets in  $Y$ .

### Example

Consider the integral operator  $A : C([0, 1]) \rightarrow C([0, 1])$  equipped with the supremum norms. Define

$$Au(x) = \int_0^1 K(x, y) f(y) dy,$$

where  $K \in C([0, 1]^2)$ . We verify that  $A$  is well-defined, i.e.,  $Au \in C([0, 1])$ . Let  $x_n \rightarrow x \in [0, 1]$ .

$$\begin{aligned} |Au(x_n) - Au(x)| &= \left| \int_0^1 K(x_n, y) u(y) dy - \int_0^1 K(x, y) u(y) dy \right| \\ &\leq \int_0^1 |K(x_n, y) - K(x, y)| |u(y)| dy \leq \|K(x_n, \cdot) - K(x, \cdot)\|_\infty \|u\|_\infty. \end{aligned}$$

Since  $K$  is continuous,  $(x_n, y) \rightarrow (x, y)$  implies  $K(x_n, y) \rightarrow K(x, y)$ . It follows that  $Au(x_n) \rightarrow Au(x)$ . Hence  $Au \in C([0, 1])$ .

We claim that  $A$  is compact. Let  $\{u_n\}$  be a bounded sequence in  $C([0, 1])$ . By Arzelà-Ascoli theorem, it suffices to show that  $\{Au_n\}$  is bounded and uniformly equicontinuous. To see the boundedness, note that by assumption we have  $\|u_n\|_\infty \leq M$  for all  $n$ . Also, since  $K$  is continuous on a compact set, we have  $K(x, y) \leq C$  for all  $x, y \in [0, 1]$ . Then

$$\begin{aligned}\|Au_n\|_\infty &= \sup_{x \in [0, 1]} \left| \int_0^1 K(x, y) u_n(y) dy \right| \\ &\leq \sup_{x \in [0, 1]} \int_0^1 |K(x, y)| |u_n(y)| dy \\ &\leq \sup_{x \in [0, 1]} \int_0^1 C \|u_n\|_\infty dy = C \|u_n\|_\infty \leq CM.\end{aligned}$$

Thus  $\{Au_n\}$  is bounded. To see the uniform equicontinuity, let  $\epsilon > 0$  be given. By continuity of  $K$ , we can find  $\delta > 0$  such that whenever  $|x - z| < \delta$ ,  $|K(x, y) - K(z, y)| < \epsilon/M$  for all  $y \in [0, 1]$ . Then for each  $n \in \mathbb{N}$ ,

$$\begin{aligned}|Au_n(x) - Au_n(z)| &= \left| \int_0^1 K(x, y) u_n(y) dy - \int_0^1 K(z, y) u_n(y) dy \right| \\ &\leq \int_0^1 |K(x, y) - K(z, y)| |u_n(y)| dy \leq \frac{\epsilon}{M} \|u_n\|_\infty \leq \epsilon.\end{aligned}$$

Thus  $\{Au_n\}$  is uniformly equicontinuous. By Arzelà-Ascoli theorem,  $\{Au_n\}$  has a convergent subsequence, i.e.,  $A$  is compact.

### Definition 3.40

Let  $A : \mathcal{H} \rightarrow \mathcal{H}$  be a linear operator. If  $\lambda \in \mathbb{F}$  is an eigenvalue of  $A$ , then the corresponding **eigenspace** is defined as

$$E_\lambda = \{x \in \mathcal{H} : Ax = \lambda x\}.$$

### Remark

Clearly  $E_\lambda$  is a subspace of  $\mathcal{H}$ .

### Theorem 3.41 (Spectral Theorem for Compact Symmetric Operators)

Let  $\mathcal{H}$  be a separable Hilbert space. Suppose that  $A : \mathcal{H} \rightarrow \mathcal{H}$  is a symmetric compact linear operator. Then the followings are true.

- (a) There exists an at most countable orthonormal basis  $\{x_j\}$ , in which each  $x_j$  is an eigenvector of  $A$  corresponding to an eigenvalue  $\lambda_j$ .
- (b) If  $\lambda_i \neq \lambda_j$ , then  $\langle x_i, x_j \rangle = 0$ .
- (c) For any  $\lambda \neq 0$ ,  $\dim(E_\lambda) < \infty$ .
- (d) If  $\dim(\mathcal{H}) = \infty$ , then either  $\lambda_j \rightarrow 0$  or there is only finitely many  $\lambda_j \neq 0$ .



*Proof.* First note that if  $\mathcal{H} = \{0\}$ , then the statements are vacuously true. We assume that  $\mathcal{H} \neq \{0\}$ . Assume first that  $\dim(\mathcal{H}) = \infty$  and  $\ker(A) = \{0\}$ . By the assumptions we can find  $x \in \mathcal{H}$  such that  $\|Ax\| > 0$  and thus  $\|A\| > 0$ . By **proposition 3.38**,  $\|A\| = \sup_{\|x\|=1} |\langle Ax, x \rangle|$ . Hence there exists a sequence  $z_n \in \mathcal{H}$  such that  $|\langle Az_n, z_n \rangle| \rightarrow \|A\|$  with  $\|z_n\| = 1$ . Let  $\lambda_1 \in \mathbb{R}$  satisfying that  $\lambda_1 = \operatorname{sgn}(\langle Az_n, z_n \rangle) \|A\|$  for  $n$  greater than some  $N$  so that the sign of  $\langle Az_n, z_n \rangle$  does not alternate. Now notice that

$$\begin{aligned} 0 &\leq \|\lambda_1 z_n - Az_n\|^2 \\ &= |\lambda_1|^2 \|z_n\|^2 + \|Az_n\|^2 - 2|\lambda_1| \langle Az_n, z_n \rangle \\ &\leq 2|\lambda_1|^2 - 2\lambda_1 \langle Az_n, z_n \rangle \leq 2|\lambda_1|^2 - 2|\lambda_1| |\langle Az_n, z_n \rangle| \rightarrow 0. \end{aligned}$$

Hence  $\lambda_1 z_n - Az_n \rightarrow 0$ . Since  $A$  is compact,  $\{Az_n\}$  has a convergent subsequence, say  $A(z_{n_k}) \rightarrow y$ . Then  $\lambda_1 z_{n_k} \rightarrow \lambda_1 x_1$  for some  $x_1 \in \mathcal{H}$  with  $Ax_1 = y$  and then  $z_{n_k} \rightarrow x_1$ . Since  $A$  is continuous,  $A(z_{n_k}) \rightarrow Ax_1$  implies that

$$Ax_1 = \lim_{k \rightarrow \infty} A(z_{n_k}) = \lim_{k \rightarrow \infty} \lambda_1 z_{n_k} = \lambda_1 x_1.$$

Note that  $\|z_{n_k}\| = 1$  and thus  $\|x_1\| = 1$ . We have shown that there exists an eigenvector  $x_1$  corresponding to an eigenvalue  $\lambda_1$ , with  $\|x_1\| = 1$  and  $|\lambda_1| = \|A\|$ .

Next, define  $W_1 = \operatorname{span}(\{x_1\})$  and  $W_1^\perp = \{y \in \mathcal{H} \mid \langle y, x_1 \rangle = 0\}$ . Consider  $A_1 = A|_{W_1^\perp}$ . We verify that  $A_1 : W_1^\perp \rightarrow W_1^\perp$  is well-defined. For any  $y \in W_1^\perp$ ,

$$\langle A_1 y, x_1 \rangle = \langle Ay, x_1 \rangle = \langle y, Ax_1 \rangle = \langle y, \lambda_1 x_1 \rangle = \lambda_1 \langle y, x_1 \rangle = 0.$$

Hence  $A_1 y \in W_1^\perp$ . Observe that  $A_1$  is also symmetric since for every  $y_1, y_2 \in W_1^\perp$ ,

$$\langle A_1 y_1, y_2 \rangle = \langle Ay_1, y_2 \rangle = \langle y_1, Ay_2 \rangle = \langle y_1, A_1 y_2 \rangle.$$

We show that  $A_1$  is compact. Suppose  $y_n \in W_1^\perp$  and  $y_n \rightarrow y \in W_1^\perp$ . Then  $A_1 y_n = Ay_n \rightarrow Ay = A_1 y$  by the continuity of  $A$ . Also, if  $\{y_n\}$  is a bounded sequence in  $W_1^\perp$ , then  $\{A_1 y_n\} = \{Ay_n\} \subset W_1^\perp$  has a convergent subsequence. Since  $W_1$  is finite-dimensional,  $W_1$  is itself closed and thus so does  $W_1^\perp$  by **proposition 3.10**. It follows that the subsequence converges in  $W_1^\perp$ . Hence  $A_1$  is compact.

Now by similar argument as above, we can find  $x_2 \in W_1^\perp$  such that  $\|x_2\| = 1$ ,  $|\lambda_2| = \|A_1\| \leq \|A\| = |\lambda_1|$  and  $Ax_2 = A_1 x_2 = \lambda_2 x_2$  for some  $\lambda_2 \in \mathbb{R}$ . Continue the process. We obtain a sequence  $\{x_j\}$  such that each  $x_j$  is an eigenvector of  $A$  corresponding to an eigenvalue  $\lambda_j$  with  $|\lambda_j| \leq |\lambda_{j-1}|$  and  $\|x_j\| = 1$ . Furthermore, observe that  $\langle x_i, x_j \rangle = 0$  for all  $i \neq j$  and  $|\lambda_{i+1}| = \|A_i\|$ .

We verify (d) first. Notice that  $|\lambda_j|$  decreases and bounded below by 0. Hence  $|\lambda_j| \rightarrow \alpha$ . By the compactness of  $A$ , there exists a subsequence  $x_{n_i}$  such that  $Ax_{n_i}$  converges. Thus  $Ax_{n_i}$

is Cauchy and

$$\|Ax_{n_i} - Ax_{n_j}\|^2 = \|\lambda_{n_i}x_{n_i} - \lambda_{n_j}x_{n_j}\|^2 = |\lambda_{n_i}|^2 + |\lambda_{n_j}|^2.$$

Note that the left hand side converges to 0 and the right hand side converges to  $2\alpha^2$ . Hence  $\alpha = 0$ .

Next, we show that  $\dim(E_\lambda) < \infty$  for  $\lambda \neq 0$ . Suppose not. Then we can find a countable orthonormal basis  $\{x_i\}$  of  $E_\lambda$  with each  $x_i$  corresponding to  $\lambda$ . By the compactness of  $A$ , there exists a subsequence  $x_{n_i}$  such that  $Ax_{n_i}$  converges and thus Cauchy.

$$2|\lambda|^2 = |\lambda|^2 \|x_{n_i}\|^2 + |\lambda|^2 \|x_{n_j}\|^2 = \|\lambda x_{n_i} - \lambda x_{n_j}\|^2 = \|Ax_{n_i} - Ax_{n_j}\|^2 \rightarrow 0.$$

Hence  $|\lambda| = 0$ , a contradiction. Thus  $\dim(E_\lambda) < \infty$ .

Lastly, we show that  $\text{span}(\{x_j\}) = \mathcal{H}$ . For every  $x \in \mathcal{H}$ , consider the partial sum  $z_n = \sum_{i=1}^n \langle x, x_i \rangle x_i$ . We want to show that  $z_n \rightarrow x$ .  $Az_n = \sum_{i=1}^n \lambda_i \langle x, x_i \rangle x_i$ . Notice that

$$\langle x - z_n, x_j \rangle = \left\langle x - \sum_{i=1}^n \langle x, x_i \rangle x_i, x_j \right\rangle = \langle x, x_j \rangle - \langle x, x_j \rangle = 0.$$

Hence  $x - z_n \in W_n^\perp$ . Thus since  $\|x - z_n\|$  is bounded,

$$\|Ax - Az_n\| = \|A(x - z_n)\| = \|A_n(x - z_n)\| \leq \|A_n\| \|x - z_n\| = |\lambda_{n+1}| \|x - z_n\| \rightarrow 0$$

Hence  $Az_n \rightarrow Ax$  and thus we can write  $Ax = \sum_j \lambda_j \langle x, x_j \rangle x_j$ . If  $y = \sum_j \langle x, x_j \rangle x_j$ , then  $Ay = \sum_j \lambda_j \langle x, x_j \rangle x_j = Ax$ . Because  $A$  has zero kernel,  $x = y = \sum_j \langle x, x_j \rangle x_j$ . Thus we have  $\text{span}(\{x_j\}) = \mathcal{H}$ .

Now we drop the assumption that  $\ker(A) = \{0\}$ . Note that  $\ker(A)$  is a closed subspace of  $\mathcal{H}$  since if  $x_n \in \ker(A)$  and  $x_n \rightarrow x \in \mathcal{H}$ , by continuity of  $A$ ,

$$Ax = \lim_{n \rightarrow \infty} Ax_n = 0 \implies x \in \ker(A).$$

It follows that by **proposition 3.10**,  $\mathcal{H} = \ker(A) \oplus \ker(A)^\perp$ . For  $\ker(A)$ , we apply **theorem 3.20** to find an orthonormal basis  $\{w_k\}$  of  $\ker(A)$ . Note that since  $\{w_k\} \subset \ker(A)$ , each  $w_k$  is an eigenvector of  $A$  corresponding to the eigenvalue 0. Also, define  $A^\perp : \ker(A)^\perp \rightarrow \ker(A)^\perp$  by  $A^\perp y = Ay$ . We verify that such definition is well-defined, i.e.,  $A^\perp y \in \ker(A)^\perp$ . For each  $y \in \ker(A)^\perp$ ,

$$\langle A^\perp y, w \rangle = \langle Ay, w \rangle = \langle y, Aw \rangle = 0$$

for all  $w \in \ker(A)$ . Hence  $A^\perp y \in \ker(A)^\perp$ . Also,  $A^\perp$  inherits the compactness and symmetry of  $A$ . By the previous argument, we can find an orthonormal basis  $\{x_j\}$  of  $\ker(A)^\perp$  such that each  $x_j$  is an eigenvector of  $A^\perp$  and thus  $A$ , corresponding to an eigenvalue  $\lambda_j$ . Notice that  $\langle w_k, x_j \rangle = 0$ . Thus  $\{w_k\} \cup \{x_j\}$  forms the desired orthonormal basis of  $\mathcal{H}$ .

Finally, if  $\dim(\mathcal{H}) = \infty$ , then (c) and (d) are vacuously true. (a) follows by applying the

above construction with the process being terminated in finite steps. Once (a) is established, (b) follows by observing that  $x_i$  and  $x_j$  are distinct eigenvectors in the orthonormal basis and thus must be orthogonal. ■

### Definition 3.42

Let  $\mathcal{H}$  be a separable Hilbert space and  $A : \mathcal{H} \rightarrow \mathcal{H}$  be a symmetric compact linear operator. Then for each  $x \in \mathcal{H}$ , we can find  $\{x_j\}$  and  $\{w_k\}$  are orthonormal bases for  $\ker(A)^\perp$  and  $\ker(A)$  respectively such that

$$x = \sum_j \langle x, x_j \rangle x_j + \sum_k \langle x, w_k \rangle w_k.$$

Such decomposition is called the **spectral decomposition** of  $A$ .

### Theorem 3.43 (Fredholm Alternative)

Let  $\mathcal{H}$  be separable. Suppose  $A : \mathcal{H} \rightarrow \mathcal{H}$  is a symmetric compact linear operator,  $\lambda \neq 0$ . Let  $N_\lambda = \{x \in \mathcal{H} \mid Ax = \lambda x\}$ . Then the equation

$$\lambda x - Ax = z$$

has a solution if and only if  $z \in N_\lambda^\perp$ . Furthermore, if  $\lambda$  is not an eigenvalue of  $A$ , then the solution is unique.

*Proof.* Consider the orthonormal eigenbasis  $\{x_j\} \cup \{w_k\}$  of  $A$  with nonzero eigenvalues  $\lambda_j$  for  $x_j$  and zeros for  $w_k$ . Suppose first that  $\lambda \neq \lambda_j$  for all  $j$ . This is equivalent to that  $N_\lambda = \{0\}$  and  $N_\lambda^\perp = \mathcal{H}$ . For every  $z \in \mathcal{H}$ , by setting

$$x = \sum_j \frac{1}{\lambda - \lambda_j} \langle z, x_j \rangle x_j + \sum_k \frac{1}{\lambda} \langle z, w_k \rangle w_k,$$

we see that

$$\begin{aligned} \lambda x - Ax &= \lambda \sum_j \frac{1}{\lambda - \lambda_j} \langle z, x_j \rangle x_j + \lambda \sum_k \frac{1}{\lambda} \langle z, w_k \rangle w_k - \sum_j \frac{\lambda_j}{\lambda - \lambda_j} \langle z, x_j \rangle x_j \\ &= \sum_j \langle z, x_j \rangle x_j + \sum_k \langle z, w_k \rangle w_k = z. \end{aligned}$$

We verify that such  $x$  indeed belongs to  $\mathcal{H}$ .

$$\|x\|^2 = \sum_j \left| \frac{1}{\lambda - \lambda_j} \langle z, x_j \rangle \right|^2 + \sum_k \left| \frac{1}{\lambda} \langle z, w_k \rangle \right|^2 \leq \sup_j \frac{1}{|\lambda - \lambda_j|^2} \|z\|^2 + \sup_k \frac{1}{|\lambda|^2} \|z\|^2 < C_\lambda \|z\|^2$$

for some  $C_\lambda$  by the Parseval's identity. Since  $\lambda \neq 0$ ,  $C_\lambda$  is finite and thus  $x \in \mathcal{H}$ . To check the uniqueness, it suffices to show that the homogeneous equation  $\lambda x - Ax = 0$  implies  $x = 0$ . Indeed, if  $x \neq 0$  satisfies  $\lambda x - Ax = 0$ , then  $\lambda$  becomes an eigenvalue of  $A$  with eigenvector

$x$ , which contradicts to our assumption. Hence the solution is unique. The converse is trivial since  $N_\lambda^\perp = \mathcal{H}$ .

Now suppose that  $\lambda = \lambda_j$  for some  $j$ , say  $j = 1$ . Then  $N_\lambda = E_{\lambda_1}$ . If  $z \in E_{\lambda_1}^\perp$ , then since  $\dim(E_{\lambda_1}) < \infty$  by the spectral theorem,  $E_{\lambda_1}$  is a closed subspace of  $\mathcal{H}$  and hence  $E_{\lambda_1}^\perp$  by [proposition 3.10](#). Thus for such  $z$ , we can write  $z = \sum_{j:\lambda_j \neq \lambda_1} \langle z, x_j \rangle x_j + \sum_k \langle z, w_k \rangle w_k$ . Set

$$x = \sum_{j:\lambda_j \neq \lambda_1} \frac{1}{\lambda - \lambda_j} \langle z, x_j \rangle x_j + \sum_k \frac{1}{\lambda} \langle z, w_k \rangle w_k.$$

Then

$$\begin{aligned} \lambda x - Ax &= \lambda \sum_{j:\lambda_j \neq \lambda_1} \frac{1}{\lambda - \lambda_j} \langle z, x_j \rangle x_j + \lambda \sum_k \frac{1}{\lambda} \langle z, w_k \rangle w_k - \sum_{j:\lambda_j \neq \lambda_1} \frac{\lambda_j}{\lambda - \lambda_j} \langle z, x_j \rangle x_j \\ &= \sum_{j:\lambda_j \neq \lambda_1} \langle z, x_j \rangle x_j + \sum_k \langle z, w_k \rangle w_k = z. \end{aligned}$$

We verify that such  $x$  indeed belongs to  $\mathcal{H}$ . By exactly the same argument as above,

$$\|x\|^2 \leq \sup_{j:\lambda_j \neq \lambda} \frac{1}{|\lambda - \lambda_j|^2} \|z\|^2 + \sup_k \frac{1}{|\lambda|^2} \|z\|^2 < C_\lambda \|z\|^2.$$

Since by the spectral theorem we have  $\lambda_j \rightarrow 0$  as  $j \rightarrow \infty$ ,  $C_\lambda < \infty$ . We conclude that the equation  $\lambda x - Ax = z$  has a solution if  $z \in N_\lambda^\perp$ . Conversely, if  $x$  is a solution, then for every  $x_{j_i}$  with  $\lambda_{j_i} = \lambda_1 = \lambda$ ,

$$\begin{aligned} \langle z, x_{j_i} \rangle &= \langle \lambda x - Ax, x_{j_i} \rangle = \langle \lambda x, x_{j_i} \rangle - \langle Ax, x_{j_i} \rangle \\ &= \lambda \langle x, x_{j_i} \rangle - \langle x, Ax_{j_i} \rangle = 0. = \lambda \langle x, x_{j_i} \rangle - \lambda_{j_i} \langle x, x_{j_i} \rangle = 0. \end{aligned}$$

We see that  $z \in N_\lambda^\perp$ . This completes the proof. ■

### Remark

*If  $\lambda$  is an eigenvalue of  $A$ , we can actually find infinitely many solutions. Since every eigenvector  $x$  corresponding to  $\lambda$  would become a homogeneous solution, for any solution  $x_0$  such that  $\lambda x_0 - Ax_0 = z$ ,  $x_0 + x$  forms another solution for any  $x \in E_\lambda$ . In other words, the set of solutions is  $x_0 + E_\lambda$ .*

## 4. Approximation Theory and Fourier Theory

### 4.1. Approximation by Polynomials

#### Proposition 4.1

Let  $X$  be a finite-dimensional vector space. Then every norm on  $X$  is equivalent.

*Proof.* This can be seen as a special case of [proposition 2.83](#), as any finite-dimensional vector space is a Banach space. However, we also have a simple proof here.

Let  $\{e_1, \dots, e_n\}$  be a basis for  $X$ . For any  $x \in X$ , we can write  $x = \sum_{i=1}^n x_i e_i$ . For any norm  $\|\cdot\|$  on  $X$ ,

$$\|x\| = \left\| \sum_{i=1}^n x_i e_i \right\| \leq \sum_{i=1}^n |x_i| \|e_i\| \leq \left( \max_{1 \leq i \leq n} \|e_i\| \right) \sum_{i=1}^n |x_i| = C_1 \|x\|_1.$$

where  $\|\cdot\|_1$  is the  $\ell^1$  norm. Also, this implies that  $\|\cdot\| : X \rightarrow \mathbb{R}$  is continuous with respect to the  $\ell^1$  norm since

$$|\|x\| - \|y\|| \leq \|x - y\| \leq C_1 \|x - y\|_1.$$

Now for any  $x \neq 0$ , the function  $f(x) = \left\| \frac{x}{\|x\|_1} \right\|$  is continuous on  $S = \{x \in X \mid \|x\|_1 = 1\}$ , which is compact. By extreme value theorem,  $f$  attains its minimum on  $S$ , which leads to

$$\frac{\|x\|}{\|x\|_1} = \left\| \frac{x}{\|x\|_1} \right\| \geq C_2 > 0$$

for some  $C_2 > 0$  since  $x \neq 0$ . Thus  $\|x\| \geq C_2 \|x\|_1$  and the norms are equivalent. ■

#### Remark

Every finite-dimensional normed vector space is complete.

#### Remark

Every closed ball in a finite-dimensional normed vector space is compact.

#### Theorem 4.2

Let  $X$  be a Banach space and  $Y$  be a finite-dimensional subspace. For any  $x \in X$ , there exists a  $y^* \in Y$  such that

$$\|x - y^*\| = \inf_{y \in Y} \|x - y\|.$$

*Proof.* Since  $Y$  is a subspace,  $0 \in Y \subset X$ . Then  $\|x - y^*\| \leq \|x\|$ . Consider the closed ball  $B = \{y \in Y \mid \|x - y\| \leq \|x\|\}$ . Let  $f(y) = \|x - y\|$ . Observe that

$$|f(y) - f(z)| = |\|x - y\| - \|x - z\|| \leq \|y - z\|,$$

and  $f$  is continuous. Since  $B$  is compact,  $f$  attains its minimum at some point  $y^* \in B \subset Y$ . Thus  $\|x - y^*\| = \inf_{y \in Y} \|x - y\|$ . ■

**Proposition 4.3**

Let  $X$  be a Banach space and  $Y \subset X$  be a finite-dimensional subspace. Suppose that for each  $x \in X$ , there corresponds a unique  $y_x \in Y$  such that  $\|x - y_x\| = \inf_{y \in Y} \|x - y\|$ . Then the map  $P : X \rightarrow Y$  defined by  $P : x \mapsto y_x$  is continuous.

*Proof.* Let  $x_n \rightarrow x$  in  $X$ . Since

$$\|P(x_n)\| = \|P(x_n) - x_n + x_n\| \leq \|P(x_n) - x_n\| + \|x_n\| \leq 2\|x_n\|,$$

$P(x_n)$  is bounded. By Bolzano-Weierstrass theorem, there is a subsequence  $x_{n_k}$  such that  $P(x_{n_k}) \rightarrow P(\hat{x})$  for some  $\hat{x} \in X$ . It remains to show that  $P(x) = P(\hat{x})$ . Indeed,

$$\|P(x_{n_k}) - x_{n_k}\| \leq \|P(x) - x_{n_k}\|$$

for all  $k$ . Letting  $k \rightarrow \infty$  gives  $\|P(\hat{x}) - x\| \leq \|P(x) - x\|$ . Since the minimizer is unique,  $P(x) = P(\hat{x})$  and  $x = \hat{x}$ . Hence  $P$  is continuous. ■

**Theorem 4.4**

Let  $X$  be a Banach space,  $Y \subset X$  be a subspace, and  $x \in X$ . Then the set

$$Y_x = \{y \in Y \mid y = \operatorname{argmin}_{y \in Y} \|x - y\|\}$$

is a bounded convex set.

*Proof.* Since  $0 \in Y$ , for all  $y \in Y_x$ ,  $\|y\| \leq \|x - y\| + \|x\| \leq 2\|x\|$ . Thus  $Y_x$  is bounded. To see the convexity, let  $y_1, y_2 \in Y_x$  and  $t \in [0, 1]$ . Then  $ty_1 + (1 - t)y_2 \in Y$  and

$$\begin{aligned} \|x - (ty_1 + (1 - t)y_2)\| &= \|t(x - y_1) + (1 - t)(x - y_2)\| \\ &\leq t\|x - y_1\| + (1 - t)\|x - y_2\| \\ &= \|x - y_1\| = \|x - y_2\| \end{aligned}$$

since  $y_1, y_2 \in Y_x$ . Thus  $ty_1 + (1 - t)y_2 \in Y_x$  and  $Y_x$  is convex. ■

**Definition 4.5**

A Banach space  $(X, \|\cdot\|)$  has a strictly convex norm if

$$\|x + y\| < \|x\| + \|y\|$$

for all  $x, y \in X$  such that  $\alpha x \neq \beta y$  for all  $\alpha, \beta \in \mathbb{R}$ .

**Remark**

$\mathcal{L}^p$  spaces are strictly convex for  $1 < p < \infty$ .

**Definition 4.6**

For any bounded function  $f : [0, 1] \rightarrow \mathbb{R}$ , the Bernstein polynomial of degree  $n$  for  $f$  is defined

as

$$B_n(f)(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}.$$

**Theorem 4.7 (Weierstrass)**

Let  $f \in C([a, b])$ . Then for any  $\epsilon > 0$ , there exists a polynomial  $p$  such that  $\|f - p\|_\infty < \epsilon$ .

*Proof.* First consider the mapping  $\sigma : x \mapsto a + (b - a)x$  for  $x \in [a, b]$ . Then by replacing  $f$  with  $f \circ \sigma$ , we can assume that  $a = 0$  and  $b = 1$ .

Now consider the Bernstein polynomial  $B_n(f)$ . Since  $f$  is continuous on  $[0, 1]$ , which is compact,  $f$  is uniformly continuous. Thus for any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \epsilon$ . Let  $F = \{k \in \{0, \dots, n\} \mid |x - \frac{k}{n}| < \delta\}$ . We can compute that

$$\begin{aligned} |B_n(f)(x) - f(x)| &= \left| f(x) - \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} \right| \\ &\leq \sum_{k \in F} \left| f(x) - f\left(\frac{k}{n}\right) \right| \binom{n}{k} x^k (1-x)^{n-k} + \sum_{k \notin F} \left| f(x) - f\left(\frac{k}{n}\right) \right| \binom{n}{k} x^k (1-x)^{n-k} \\ &\leq \sum_{k=0}^n \epsilon \binom{n}{k} x^k (1-x)^{n-k} + 2 \|f\|_\infty \sum_{k \notin F} \binom{n}{k} x^k (1-x)^{n-k} \\ &\leq \epsilon + 2 \|f\|_\infty \sum_{k=0}^n \mathbb{1}_{\left\{ \left| x - \frac{k}{n} \right| \geq \delta \right\}} \binom{n}{k} x^k (1-x)^{n-k} \\ &\leq \epsilon + 2 \|f\|_\infty \sum_{k=0}^n \frac{1}{\delta^2} \left( x - \frac{k}{n} \right)^2 \binom{n}{k} x^k (1-x)^{n-k} \\ &= \epsilon + 2 \frac{\|f\|_\infty}{\delta^2} \sum_{k=0}^n \left( x^2 - \frac{2k}{n} x + \frac{k^2}{n^2} \right) \binom{n}{k} x^k (1-x)^{n-k}. \end{aligned}$$

Now let

$$S(x, y) = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} = (x + y)^n.$$

Then

$$\begin{aligned} nx(x + y)^{n-1} &= x \frac{\partial S}{\partial x} = \sum_{k=0}^n k \binom{n}{k} x^k y^{n-k}, \\ n(n-1)x^2(x + y)^{n-2} &= x^2 \frac{\partial^2 S}{\partial x^2} = \sum_{k=0}^n k(k-1) \binom{n}{k} x^k y^{n-k}. \end{aligned}$$

Taking  $y = 1 - x$  gives

$$\sum_{k=0}^n \left( x^2 - \frac{2k}{n} x + \frac{k^2}{n^2} \right) \binom{n}{k} x^k (1-x)^{n-k} = x^2 - \frac{2}{n} x \cdot nx + \frac{1}{n^2} (n(n-1)x^2 + nx) = \frac{x(1-x)}{n} \leq \frac{1}{4n}.$$

Hence we obtain the estimate

$$|B_n(f)(x) - f(x)| \leq \epsilon + \frac{\|f\|_\infty}{2n\delta^2}.$$

Letting  $n \rightarrow \infty$  and by the arbitrariness of  $\epsilon$ , we see that  $B_n(f) \rightarrow f$  uniformly. ■

**Remark**

*An alternative expression for the Weierstrass theorem is that for such  $f$ , there exists a sequence of polynomials  $p_n$  such that  $p_n \rightarrow f$  uniformly.*

**Remark**

*As a direct consequence of the Weierstrass theorem, the polynomial space is dense in  $C([0, 1])$ .*

**Definition 4.8**

A map  $T : C([a, b]) \rightarrow C([a, b])$  is said to be **positive** if  $T(f) \geq 0$  for all  $f \geq 0$ .

**Proposition 4.9**

The map  $U : C([0, 1]) \rightarrow C([0, 1])$  defined by  $f \mapsto B_n(f)$  is linear, positive, and continuous.

*Proof.* To show the linearity, let  $c \in \mathbb{R}$  and  $f, g \in C([0, 1])$ . Then

$$\begin{aligned} U(cf + g)(x) &= \sum_{k=0}^n \left( cf\left(\frac{k}{n}\right) + g\left(\frac{k}{n}\right) \right) \binom{n}{k} x^k (1-x)^{n-k} \\ &= c \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} + \sum_{k=0}^n g\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} = cU(f)(x) + U(g)(x). \end{aligned}$$

To show the positivity, let  $f \geq 0$ . Then  $f\left(\frac{k}{n}\right) x^k (1-x)^{n-k} \geq 0$  for all  $k$  and  $x \in [0, 1]$ . Then the sum is nonnegative and  $U(f) \geq 0$ .

To show the continuity, it is enough to show the boundedness of  $U$ .

$$\begin{aligned} \|U(f)\|_\infty &= \sup_{x \in [0, 1]} \left| \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} \right| \\ &\leq \sup_{x \in [0, 1]} \sum_{k=0}^n \left| f\left(\frac{k}{n}\right) \right| \binom{n}{k} x^k (1-x)^{n-k} \\ &\leq \sup_{x \in [0, 1]} \sum_{k=0}^n \|f\|_\infty \binom{n}{k} x^k (1-x)^{n-k} = \|f\|_\infty. \end{aligned}$$

Hence  $U$  is a bounded linear operator and thus continuous. ■

**Theorem 4.10 (Korovkin)**

Let  $T_n : C([0, 1]) \rightarrow C([0, 1])$  be positive linear maps. Suppose that  $T_n(f_i) \rightarrow f_i$  uniformly for  $i = 0, 1, 2$  with  $f_i(x) = x^i$ . Then  $T_n(f) \rightarrow f$  uniformly for all  $f \in C([0, 1])$ .



*Proof.* Let  $\epsilon > 0$ . Since  $f$  is continuous on a compact set, we can assume that it is Lipschitz with constant  $L$ . Now observe that

$$|f(x) - f(a)| \leq L|x - a| \leq L\epsilon + L\frac{(x-a)^2}{\epsilon}.$$

This can be verify as follows,

$$\begin{cases} L|x - a| \leq L\epsilon \leq L\epsilon + L\frac{(x-a)^2}{\epsilon} & \text{if } |x - a| \leq \epsilon, \\ L|x - a| \leq L\frac{(x-a)^2}{\epsilon} \leq L\epsilon + L\frac{(x-a)^2}{\epsilon} & \text{if } |x - a| > \epsilon. \end{cases}$$

Next, we apply  $T_n$  and note that we have  $|T_n(f)| = T_n(|f|)$ . Then,

$$\begin{aligned} |T_n(f)(x) - f(a)| &\leq |T_n(f)(x) - f(a)T_n(f_0)(x)| + |f(a)T_n(f_0)(x) - f(a)| \\ &= T_n(|f - f(a)|)(x) + |f(a)| |T_n(f_0)(x) - f_0| \\ &\leq L \left( \epsilon T_n(f_0)(x) + \frac{1}{\epsilon} T_n(f_2 - 2af_1 + a^2f_0)(x) \right) + \|f\|_\infty \|T_n(f_0) - f_0\|_\infty \\ &\leq L\epsilon(T_n(f_0)(x) - f_0(x)) + L\epsilon f_0(x) + \frac{L}{\epsilon}(T_n(f_2)(x) - f_2(x)) + \frac{L}{\epsilon}f_2(x) \\ &\quad - \frac{2aL}{\epsilon}(T_n(f_1)(x) - f_1(x)) - \frac{2aL}{\epsilon}f_1(x) + \frac{a^2L}{\epsilon}(T_n(f_0)(x) - f_0(x)) \\ &\quad + \frac{a^2L}{\epsilon}f_0(x) + \|f\|_\infty \|T_n(f_0) - f_0\|_\infty \\ &\leq L\epsilon \|T_n(f_0) - f_0\|_\infty + \frac{L}{\epsilon} \|T_n(f_2) - f_2\|_\infty + \frac{2aL}{\epsilon} \|T_n(f_1) - f_1\|_\infty \\ &\quad + \frac{a^2L}{\epsilon} \|T_n(f_0) - f_0\|_\infty + \|f\|_\infty \|T_n(f_0) - f_0\|_\infty + L\epsilon + \frac{L}{\epsilon}(x-a)^2. \end{aligned}$$

Now taking  $a = x$  and then taking supremum over  $x \in [0, 1]$  gives

$$\begin{aligned} \|T_n(f) - f\|_\infty &\leq L\epsilon \|T_n(f_0) - f_0\|_\infty + \frac{L}{\epsilon} \|T_n(f_2) - f_2\|_\infty \\ &\quad + \frac{2L}{\epsilon} \|T_n(f_1) - f_1\|_\infty + \frac{L}{\epsilon} \|T_n(f_0) - f_0\|_\infty + \|f\|_\infty \|T_n(f_0) - f_0\|_\infty + L\epsilon. \end{aligned}$$

By the assumptions, there is  $N$  such that  $n > N$  implies that

$$\|T_n(f_i) - f_i\|_\infty < \epsilon^2, \quad i = 0, 1, 2.$$

Thus,

$$\|T_n(f) - f\|_\infty < L\epsilon^3 + 5L\epsilon + \|f\|_\infty \epsilon^2.$$

Since  $\epsilon$  is arbitrary, we obtain that  $T_n(f) \rightarrow f$  uniformly. ■

### Example

Let  $f \in C([0, 1])$  and  $L_n(f)$  be the polygonal approximation of  $f$  with nodes at  $k/n$  for  $k =$

$0, \dots, n$ , i.e.,

$$L_n(f)(x) = f\left(\frac{k}{n}\right) + n\left(f\left(\frac{k+1}{n}\right) - f\left(\frac{k}{n}\right)\right)\left(x - \frac{k}{n}\right), \quad x \in \left[\frac{k}{n}, \frac{k+1}{n}\right].$$

Now  $L_n(1) = 1$ ,  $L_n(x) = x$ , and  $\|L_n(x^2) - x^2\|_\infty \leq \max_{0 \leq k \leq n-1} \frac{(k+1)^2}{n^2} - \frac{k^2}{n^2} \leq \frac{1}{n} \rightarrow 0$ . By the Korovkin theorem, we can conclude that  $L_n(f) \rightarrow f$  uniformly for all  $f \in C([0, 1])$ .

#### Definition 4.11

Let  $f$  be a bounded function on  $[a, b]$ . The **modulus of continuity** of  $f$  is defined as

$$\omega_f(\delta) = \sup_{\substack{|x-y| \leq \delta \\ x, y \in [a, b]}} |f(x) - f(y)|.$$

#### Definition 4.12

A function  $f$  is said to be **Lipschitz of order  $\alpha$**  if

$$|f(x) - f(y)| \leq M |x - y|^\alpha$$

for some  $M > 0$  and all  $x, y \in [a, b]$ .

#### Proposition 4.13

Let  $f$  be a bounded function on  $[a, b]$ . Then

- (a)  $\omega_f(\delta_1) \leq \omega_f(\delta_2)$  for all  $\delta_1 \leq \delta_2$ .
- (b) If  $f'$  exists and is bounded, then  $\omega_f(\delta) \leq M\delta$  for some  $M$ .
- (c) If  $f$  is Lipschitz of order  $\alpha$ , then  $\omega_f(\delta) \leq M\delta^\alpha$  for some  $M$  and all  $\delta > 0$ .

*Proof.* For (a), note that we have  $|x - y| \leq \delta_1 \leq \delta_2$  for all  $x, y \in [a, b]$ .

For (b), from the mean value theorem, we have that if  $|x - y| \leq \delta$ , then

$$|f(x) - f(y)| = |f'(c)(x - y)| \leq M |x - y| \leq M\delta$$

for some  $c \in [a, b]$  and some  $M > 0$ .

For (c), we have that

$$|f(x) - f(y)| \leq M |x - y|^\alpha \leq M\delta^\alpha$$

for  $|x - y| \leq \delta$ . ■

#### Lemma 4.14

Let  $f$  be a bounded function on  $[a, b]$  and  $\delta > 0$ . Then

- (a)  $\omega_f(n\delta) \leq n\omega_f(\delta)$  for all  $n \in \mathbb{N}$ .
- (b)  $\omega_f(\lambda\delta) \leq (1 + \lambda)\omega_f(\delta)$  for all  $\lambda > 0$ .

*Proof.* For (a), let  $x < y$  be such that  $|x - y| \leq n\delta$ . We can split  $[x, y]$  into  $n$  intervals of length at most  $\delta$ , say  $[z_0, z_1], \dots, [z_{n-1}, z_n]$ . Then  $|z_i - z_{i-1}| \leq \delta$  for all  $i$  and

$$|f(x) - f(y)| \leq \sum_{i=1}^n |f(z_i) - f(z_{i-1})| \leq n\omega_f(\delta).$$

For (b), let  $n \in \mathbb{N}$  be such that  $n - 1 \leq \lambda \leq n$ . Then

$$\omega_f(\lambda\delta) \leq \omega_f(n\delta) \leq n\omega_f(\delta) \leq (1 + \lambda)\omega_f(\delta).$$

■

### Theorem 4.15

For any  $f \in C([0, 1])$ , the Bernstein polynomial  $B_n(f)$  satisfies

$$\|B_n(f) - f\|_\infty \leq \frac{3}{2}\omega_f\left(\frac{1}{\sqrt{n}}\right).$$

*Proof.* By lemma 4.14, setting  $\delta = 1/\sqrt{n}$  and  $\lambda = \sqrt{n}|x - \frac{k}{n}|$ , then

$$\begin{aligned} |f(x) - B_n f(x)| &\leq \sum_{k=0}^n \left| f(x) - f\left(\frac{k}{n}\right) \right| \binom{n}{k} x^k (1-x)^{n-k} \leq \sum_{k=0}^n \omega_f\left(\left|x - \frac{k}{n}\right|\right) \binom{n}{k} x^k (1-x)^{n-k} \\ &\leq \sum_{k=0}^n \omega_f\left(\frac{1}{\sqrt{n}}\right) \left(1 + \sqrt{n}\left|x - \frac{k}{n}\right|\right) \binom{n}{k} x^k (1-x)^{n-k} \\ &\leq \omega_f\left(\frac{1}{\sqrt{n}}\right) \left\{ 1 + \sqrt{n} \sum_{k=0}^n \left|x - \frac{k}{n}\right| \binom{n}{k} x^k (1-x)^{n-k} \right\} \\ &\leq \omega_f\left(\frac{1}{\sqrt{n}}\right) \left\{ 1 + \sqrt{n} \left( \sum_{k=0}^n \left|x - \frac{k}{n}\right|^2 \binom{n}{k} x^k (1-x)^{n-k} \right)^{1/2} \left( \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} \right)^{1/2} \right\} \\ &\leq \omega_f\left(\frac{1}{\sqrt{n}}\right) \left\{ 1 + \sqrt{n} \frac{1}{2\sqrt{n}} \right\} = \frac{3}{2}\omega_f\left(\frac{1}{\sqrt{n}}\right). \end{aligned}$$

The fourth inequality follows from the Cauchy-Schwarz inequality. ■

### Theorem 4.16

Let  $X$  be a metric space and  $Y \subset X$  be a compact subset. Then for any  $f \in X$ , there exists a  $p^* \in Y$  such that  $d(f, p^*) \leq d(f, q)$  for all  $q \in Y$ .

*Proof.* Let  $d^*$  be the shortest distance from  $f$  to  $Y$ , i.e.,  $d^* = \inf_{q \in Y} d(f, q)$ . Then there exists a sequence  $q_n \in Y$  such that  $d(f, q_n) \rightarrow d^*$ . From the compactness of  $Y$ , there exists a subsequence  $q_{n_k}$  such that  $q_{n_k} \rightarrow p^* \in Y$ . We claim that  $p^*$  is the desired point. Indeed, for any  $\epsilon > 0$ , there is an  $N$  such that  $d(f, q_{n_k}) \leq d^* + \epsilon$  and  $d(q_{n_k}, p^*) \leq \epsilon$  for all  $k \geq N$ . Then

$$d(f, p^*) \leq d(f, q_{n_k}) + d(q_{n_k}, p^*) \leq d^* + 2\epsilon.$$

Since  $\epsilon$  is arbitrary, we have  $d(f, p^*) \leq d^*$ , which completes the proof.  $\blacksquare$

#### Theorem 4.17

If  $X$  is a strictly convex Banach space, and  $Y$  is a convex compact subset of  $X$ , then for any  $f \in X$ , there exists a unique  $p^* \in Y$  such that  $\|f - p^*\| = \inf_{q \in Y} \|f - q\|$ .

*Proof.* The existence of such  $p^*$  follows from [theorem 4.16](#). Denote the shortest distance from  $f$  to  $Y$  by  $d^*$ . To show that  $p^*$  is unique, suppose that there are two such points  $p_1^*$  and  $p_2^*$ . Then by the convexity of  $Y$ ,  $\frac{1}{2}p_1^* + \frac{1}{2}p_2^* \in Y$  and from the strict convexity of  $X$ ,

$$\left\| f - \frac{1}{2}p_1^* - \frac{1}{2}p_2^* \right\| < \frac{1}{2}\|f - p_1^*\| + \frac{1}{2}\|f - p_2^*\| = d^*.$$

This contradicts the minimality of  $d^*$  and thus  $p^*$  is unique.  $\blacksquare$

#### Definition 4.18

A function  $g$  on  $[a, b]$  satisfies the **equioscillation condition** of degree  $n$  if there are  $n + 2$  points  $a \leq x_0 < x_1 < \dots < x_{n+1} \leq b$  such that  $g(x_i) = (-1)^i \|g\|$  for  $i = 0, \dots, n + 1$ .

#### Theorem 4.19 (Chebyshev Equioscillation theorem)

Let  $f \in C[a, b]$  and  $p \in P_n$  be the polynomial of degree  $n$ . Let  $r = f - p$ . Then  $r$  satisfies the equioscillation condition of degree  $n$  if and only if  $\|f - p\|_\infty \leq \|f - q\|_\infty$  for all  $q \in P_n$ .

*Proof.* First assume that  $r$  satisfies the equioscillation condition of degree  $n$ . If  $p$  is not the best approximation to  $f$  in  $P_n$ , then there is  $q \in P_n$  such that  $\|f - (p + q)\|_\infty < \|f - p\|_\infty$ . This implies that  $\|r - q\|_\infty < \|r\|_\infty$ . By the equioscillation condition,  $|r(x_i) - q(x_i)| < |r(x_i)|$  for all  $i = 0, \dots, n + 1$ . This means that  $q$  has the same sign with  $r$  at each  $x_i$ , so  $q$  must change sign  $n + 1$  times. This contradicts the fact that  $q \in P_n$ .

Conversely, suppose that  $p \in P_n$  is the best approximation to  $f$  in  $P_n$  in uniform norm. Let  $R = \|r\|_\infty$ . Since  $r$  is uniform continuous on  $[a, b]$ , we can split  $[a, b]$  into subintervals  $[t_i, t_{i+1}]$  such that  $|r(x) - r(y)| < R/2$  for all  $x, y \in [t_i, t_{i+1}]$ . Now observe that if  $[t_i, t_{i+1}]$  contains a local extremum of  $r$ , then  $r$  must have same sign in  $[t_i, t_{i+1}]$ . Denote the intervals by  $I_k$  and rearrange them so that  $r$  has maximum in  $I_1, \dots, I_{k_1}$  and minimum in  $I_{k_1+1}, \dots, I_{k_1+k_2}$ . The rest intervals are denumerated by  $I_{k_1+k_2+1}, \dots, I_{k_1+k_2+k_3}$ . By construction we see that the intervals with extremum points are disjoint.

We claim that  $k_1 + k_2 \geq n + 2$ . Assume that  $k_1 + k_2 \leq n + 1$ . Consider the polynomial

$$q(x) = \pm \prod_{i=1}^{k_1+k_2-1} (x - z_i),$$

where  $z_i$  are the points chosen with  $\max I_i < z_i < \min I_{i+1}$  for  $i = 1, \dots, k_1 + k_2 - 1$ . Notice that  $q(x) \neq 0$  for all  $x$  lying in  $I_i$  for  $i = 1, \dots, k_1 + k_2$ . We select the sign of  $q$  such that  $q$  has the same sign as  $r$  in  $I_i$  for  $i = 1, \dots, k_1 + k_2$ . We show that  $p + \lambda q$  gives a better approximation

than  $p$  for some  $\lambda > 0$ . Let  $S = \cup_{i=1}^{k_1+k_2} I_i$  and  $N = \cup_{i=k_1+k_2+1}^{k_1+k_2+k_3} I_i$ . Then for  $x \in S$ ,

$$|f(x) - (p(x) + \lambda q(x))| = |r(x) - \lambda q(x)| \leq R - \lambda \min |q(x)| < R.$$

And for  $x \in N$ ,

$$|f(x) - (p(x) + \lambda q(x))| = |r(x) - \lambda q(x)| \leq R + \lambda \max |q(x)| < \frac{R}{2} + \lambda \|q\|_\infty < R$$

by taking  $\lambda = \frac{R}{2\|q\|_\infty}$ . This contradicts the assumption that  $p$  is the best approximation in  $P_n$ . Hence  $k_1 + k_2 \geq n + 2$ . Since  $p$  lies in  $P_n$ ,  $p$  has at most  $n$  interior extremum plus the two endpoints; we have  $n + 2$  extremum points, yielding the equioscillation condition. ■

### Remark

The Chebyshev equioscillation theorem gives us a way to compute the best polynomial approximation to a function in uniform norm. The approach is as follows. Consider the approximation polynomial  $p(x) = \sum_{k=0}^n a_k x^k$  and the error  $h = \|f - p\|_\infty$ . Our goal is to find the coefficients  $a_k$  and the error  $h$  as well. The Chebyshev equioscillation theorem gives the extremum points  $x_0, \dots, x_{n+1}$  such that  $f(x_i) = (-1)^i h$ . Then we have the system of equations

$$\begin{pmatrix} 1 & x_0 & \cdots & x_0^n & -1 \\ 1 & x_1 & \cdots & x_1^n & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & x_{n+1} & \cdots & x_{n+1}^n & (-1)^n \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \\ h \end{pmatrix} = \begin{pmatrix} f(x_0) \\ f(x_1) \\ \vdots \\ f(x_{n+1}) \end{pmatrix}.$$

Since  $x_i$  are unknown, we need to guess a set of  $x_i$  and solve the system of equations. The iteration continues until  $\|f - p\|_\infty = h$ .

## 4.2. Fourier Series

### Definition 4.20

The **Fourier series** of a function  $f$  is given by

$$Sf(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx),$$

where the **Fourier coefficients**  $a_k$  and  $b_k$  are given by

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(kx) dx, \quad b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(kx) dx.$$

Or, alternatively,

$$Sf(x) = \sum_{k=-\infty}^{\infty} c_k e^{ikx},$$

where the **Fourier coefficients**  $c_k$  are given by

$$c_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx.$$

**Definition 4.21**

The **Truncated Fourier series** of a function  $f$  is denoted by

$$S_N f(x) = \frac{a_0}{2} + \sum_{k=1}^N a_k \cos(kx) + b_k \sin(kx) = \sum_{k=-N}^N c_k e^{ikx}.$$

**Proposition 4.22**

Let  $a_k$  and  $b_k$  be the Fourier coefficients of a function  $f$ . Then

- (a) If  $f \in \mathcal{L}^1$ ,  $|a_k|, |b_k| \leq C \|f\|_1$  for some constant  $C > 0$ .
- (b) If  $f \in \mathcal{L}^\infty$ ,  $|a_k|, |b_k| \leq C \|f\|_\infty$  for some constant  $C > 0$ .

*Proof.* To see (a), compute that

$$|a_k| \leq \frac{1}{\pi} \int_{-\pi}^{\pi} |f(x)| |\cos(kx)| dx \leq \frac{1}{\pi} \int_{-\pi}^{\pi} |f(x)| dx = C \|f\|_1$$

and similarly for  $b_k$ . For (b),

$$|a_k| \leq \frac{1}{\pi} \int_{-\pi}^{\pi} |f(x)| |\cos(kx)| dx \leq \frac{1}{\pi} \int_{-\pi}^{\pi} \|f\|_\infty dx = 2 \|f\|_\infty = C \|f\|_\infty.$$

The proof for  $b_k$  is analogous. ■

**Lemma 4.23** (Riemann-Lebesgue I)

Let  $f \in \mathcal{L}^1[a, b]$ . Then

$$\lim_{n \rightarrow \infty} \int_a^b f(x) e^{-inx} dx = 0.$$

*Proof.* Let  $\epsilon > 0$ . Since  $f \in \mathcal{L}^1[a, b]$ , there is a step function  $g$  such that  $\|f - g\|_1 < \epsilon$ . For any interval  $E$ ,

$$\left| \int_a^b \chi_E(x) e^{-inx} dx \right| \leq \left| \int_E \cos(nx) dx \right| + \left| \int_E \sin(nx) dx \right| \leq \frac{2\pi}{n} \rightarrow 0$$

as  $n \rightarrow \infty$ . A step function is a linear combination of characteristic functions of intervals, and thus  $\left| \int_a^b g(x) e^{-inx} dx \right| \rightarrow 0$  as  $n \rightarrow \infty$ . Therefore,

$$\begin{aligned} \left| \int_a^b f(x) e^{-inx} dx \right| &\leq \left| \int_a^b (f(x) - g(x)) e^{-inx} dx \right| + \left| \int_a^b g(x) e^{-inx} dx \right| \\ &\leq \|f - g\|_1 + \left| \int_a^b g(x) e^{-inx} dx \right| \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ . ■

**Definition 4.24**

The space of piecewise continuous functions on  $[a, b]$  is denoted by  $PC[a, b]$ . The symbol  $PC^n[a, b]$  denotes the space of functions having continuous derivatives up to order  $n - 1$ , with the  $n$ -th derivative being piecewise continuous.

**Proposition 4.25**

Let  $f \in PC^1[-\pi, \pi]$  and

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx).$$

Then

$$f'(x) = \sum_{k=1}^{\infty} -ka_k \sin(kx) + kb_k \cos(kx).$$

*Proof.* Differentiation term by term gives the desired result. ■

**Remark**

If  $f \in PC^n[-\pi, \pi]$ , then

$$|a_k|, |b_k| \leq \frac{\|f\|_{PC^n}}{k^n},$$

where  $\|f\|_{PC^n} = \sum_{j=0}^n \|f^{(j)}\|_{\infty}$ .

**Definition 4.26**

Let  $f$  be a function on  $\mathbb{R}$ . The **right-limit** and the **left-limit** of  $f$  at  $x$  are defined by

$$f(x^+) = \lim_{h \rightarrow 0^+} f(x + h), \quad f(x^-) = \lim_{h \rightarrow 0^+} f(x - h).$$

**Definition 4.27**

A **kernel** is a function  $k : X \times X \rightarrow \mathbb{R}$  such that

- (a)  $k(x, y) = k(y, x)$  for all  $x, y \in X$ ,
- (b) For finitely many points  $x_1, \dots, x_n \in X$  and scalars  $a_1, \dots, a_n \in \mathbb{R}$ ,

$$\sum_{i=1}^n \sum_{j=1}^n a_i a_j k(x_i, x_j) \geq 0.$$

**Definition 4.28**

The **Dirichlet kernel** is defined by

$$D_N(x) = \frac{1}{2\pi} \sum_{k=-N}^N e^{ikx}.$$

**Remark**

The Dirichlet kernel can be simplified to

$$D_N(x) = \frac{\sin((N + 1/2)x)}{2\pi \sin(x/2)}.$$

To see this, note that

$$2\pi D_N(x)(e^{ix} - 1) = e^{i(N+1)x} - e^{-iNx} = \frac{e^{i(N+1/2)x} - e^{-i(N+1/2)x}}{e^{ix/2} - e^{-ix/2}}.$$

And thus,

$$D_N(x) = \frac{\sin((N + 1/2)x)}{2\pi \sin(x/2)}.$$

Some other properties of the Dirichlet kernel include  $D_N(-x) = D_N(x)$  and  $\int_{-\pi}^{\pi} D_N(x)dx = 1$ .

**Definition 4.29**

Let  $f, g : X \rightarrow \mathbb{R}$ . The **convolution** of  $f$  and  $g$  is defined by

$$(f * g)(x) = \int_X f(x - y)g(y)dy.$$

**Proposition 4.30**

For any  $2\pi$ -periodic function  $f \in PC$ ,

$$S_N f = D_N * f.$$

*Proof.* Compute that

$$\begin{aligned} S_N f(x) &= \sum_{k=-N}^N c_k e^{ikx} = \sum_{k=-N}^N \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) e^{-iky} dy e^{ikx} \\ &= \int_{-\pi}^{\pi} f(y) \frac{1}{2\pi} \sum_{k=-N}^N e^{ik(x-y)} dy = \int_{-\pi}^{\pi} f(y) D_N(x - y) dy = (D_N * f)(x). \end{aligned}$$

■

**Theorem 4.31** (Dirichlet-Jordan)

Let  $f$  be a  $2\pi$ -periodic function and piecewise Lipschitz. Then

$$\lim_{N \rightarrow \infty} S_N f(x) = \frac{f(x^+) + f(x^-)}{2}.$$

In particular, if  $f$  is continuous at  $x$ , then

$$\lim_{N \rightarrow \infty} S_N f(x) = f(x).$$



*Proof.* Since  $f$  is  $2\pi$ -periodic,

$$\begin{aligned}
S_N f(x) &= \int_{-\pi}^{\pi} D_N(x-y) f(y) dy = \int_{-\pi}^{\pi} D_N(y) f(x-y) dy \\
&= \int_0^{\pi} D_N(y) f(x-y) dy + \int_{-\pi}^0 D_N(y) f(x-y) dy \\
&= \int_0^{\pi} D_N(y) f(x-y) dy + \int_0^{\pi} D_N(-y) f(x+y) dy \\
&= \int_0^{\pi} D_N(y) (f(x-y) + f(x+y)) dy.
\end{aligned}$$

Notice that

$$\frac{1}{2}(f(x^+) + f(x^-)) = \int_0^{\pi} D_N(y) (f(x^+) + f(x^-)) dy.$$

Thus for given  $x$ , we have

$$\begin{aligned}
\left| S_N f(x) - \frac{f(x^+) + f(x^-)}{2} \right| &\leq \left| \int_0^{\pi} D_N(y) (f(x+y) - f(x^+)) dy \right| \\
&\quad + \left| \int_0^{\pi} D_N(y) (f(x-y) - f(x^-)) dy \right|.
\end{aligned}$$

We claim that

$$\int_0^{\pi} D_N(y) |f(x+y) - f(x^+)| dy \rightarrow 0$$

as  $N \rightarrow \infty$  and the other integral is similar. There is a  $\delta > 0$  such that  $f$  is continuous on  $[x, x + \delta]$ . Thus  $f$  is uniformly continuous on  $[x, x + \delta]$ , and  $|f(x+y) - f(x^+)| < Cy$  for some constant  $C > 0$  and  $y \in [0, \delta]$ . Then

$$\int_0^{\delta} |D_N(y)| |f(x+y) - f(x^+)| dy \leq C \int_0^{\delta} y |D_N(y)| dy \leq C \int_0^{\delta} dy = C\delta,$$

because  $|D_N(t)| \leq 1/|t|$ . On the other hand,

$$\begin{aligned}
\int_{\delta}^{\pi} |D_N(y)| |f(x+y) - f(x^+)| dy &\leq \frac{1}{2\pi} \int_{\delta}^{\pi} \frac{\sin((N+1/2)y)}{\sin(y/2)} |f(x+y) - f(x^+)| dy \\
&\leq \frac{1}{2\pi \sin(\delta/2)} \int_{\delta}^{\pi} \sin((N+1/2)y) g(y) dy \rightarrow 0,
\end{aligned}$$

as  $N \rightarrow \infty$  by the Riemann-Lebesgue lemma, where  $g(y) = |f(x+y) - f(x^+)|$  is a continuous function on  $[\delta, \pi]$ . Hence we have

$$\left| \int_0^{\pi} D_N(y) (f(x+y) - f(x^+)) dy \right| \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

We now see that

$$\left| S_N f(x) - \frac{f(x^+) + f(x^-)}{2} \right| \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

The pointwise convergence is achieved whenever  $f$  is continuous since  $f(x^+) = f(x^-) = f(x)$ . ■

**Definition 4.32**

The series  $\sigma_N f$  is defined by

$$\sigma_N f(x) = \frac{1}{N+1} \sum_{n=0}^N S_n f(x).$$

**Remark**

The series  $\sigma_N f$  is the Cesaro's mean of the Fourier series of  $f$ .

**Definition 4.33**

The **Fejer kernel** is defined by

$$F_N(x) = \frac{1}{N+1} \sum_{k=0}^N D_k(x).$$

**Remark**

The Fejer kernel can be simplified to

$$F_N(x) = \frac{\sin^2\left(\frac{N+1}{2}x\right)}{2\pi(N+1)\sin^2(x/2)}.$$

To see this, note that

$$\begin{aligned} F_N(t) &= \frac{1}{N+1} \sum_{k=0}^N D_k(t) = \frac{1}{2\pi(N+1)} \sum_{k=0}^N \frac{\sin((k+1/2)t)}{\sin(t/2)} \\ &= \frac{1}{2\pi(N+1)\sin^2(t/2)} \sum_{k=0}^N \sin((k+1/2)t) \sin(t/2) \\ &= \frac{1}{2\pi(N+1)\sin^2(t/2)} \sum_{k=0}^N \cos(kt) - \cos((k+1)t) \\ &= \frac{1}{4\pi(N+1)\sin^2(t/2)} (1 - \cos((N+1)t)) = \frac{\sin^2\left(\frac{N+1}{2}t\right)}{2\pi(N+1)\sin^2(t/2)}. \end{aligned}$$

Some other properties of the Fejer kernel include that if  $f = 1$ , then

$$\sigma_N f = \int_{-\pi}^{\pi} F_N(x) dx = 1,$$

and that  $F_N(-x) = F_N(x)$ ,  $F_N \geq 0$ . Also,

$$\sigma_N f(x) = \frac{1}{N+1} \sum_{k=0}^N \int_{-\pi}^{\pi} D_k(x-y) f(y) dy = \int_{-\pi}^{\pi} \left( \frac{1}{N+1} \sum_{k=0}^N D_k(x-y) \right) f(y) dy = F_N * f(x).$$

**Definition 4.34**

$C_{2\pi}$  denote the space of  $2\pi$ -periodic continuous functions.  $C_{2\pi}^k$  denotes the space of  $2\pi$ -periodic functions having continuous derivatives up to order  $k$ .

**Theorem 4.35** (Fejer)

Let  $f \in C_{2\pi}$ . Then  $\sigma_N f \rightarrow f$  uniformly.

*Proof.* Let  $\epsilon > 0$ . There is a  $\delta > 0$  such that  $|f(x) - f(y)| < \epsilon$  whenever  $|x - y| < \delta$ . Observe that if  $|t| \geq \delta$ , then

$$F_N(t) \leq \frac{1}{2\pi(N+1)\sin^2(\delta/2)} \rightarrow 0$$

as  $N \rightarrow \infty$ . Hence

$$\begin{aligned} |\sigma_N f(x) - f(x)| &= \left| \int_{-\pi}^{\pi} F_N(x-y) f(y) dy - f(x) \right| \leq \int_{-\pi}^{\pi} F_N(x-y) |f(y) - f(x)| dy \\ &\leq \int_{|x-y| \leq \delta} F_N(x-y) |f(y) - f(x)| dy + \int_{\delta \leq |x-y| \leq \pi} F_N(x-y) |f(y) - f(x)| dy \\ &\leq \epsilon \int_{-\pi}^{\pi} F_N(x-y) dy + 2 \|f\|_{\infty} \int_{\delta \leq |x-y| \leq \pi} F_N(x-y) dy \\ &= \epsilon + 2 \|f\|_{\infty} \frac{\pi - \delta}{2\pi(N+1)\sin^2(\delta/2)} \rightarrow 0 \end{aligned}$$

as  $N \rightarrow \infty$ . ■

**Definition 4.36**

The **trigonometric polynomial** of degree  $N$  is a function of the form

$$TP_N(x) = \sum_{k=0}^N a_k \cos(kx) + b_k \sin(kx).$$

The **trigonometric polynomial space** is denoted by  $TP = \cup_N TP_N$ .

**Theorem 4.37**

Under  $\mathcal{L}^2[-\pi, \pi]$ ,  $PC_{2\pi} \subset \overline{TP}$ .

*Proof.* Since continuous functions are dense in  $PC_{2\pi}$ , it suffices to show that continuous functions can be approximated by trigonometric polynomials. Let  $f \in C_{2\pi}$ . By the Fejer theorem,  $\sigma_N f \rightarrow f$  uniformly. Since  $\sigma_N f$  is a trigonometric polynomial,  $f$  can be approximated by trigonometric polynomials. ■

**Definition 4.38**

The **best approximation error** of a function  $f$  by a trigonometric polynomial is defined by  $\tilde{E}_N(f) = \inf_{p \in TP_N} \|p - f\|_{\infty}$ .

**Definition 4.39**

For  $f, g > 0$ ,  $f \lesssim g$  if there is some constant  $c > 0$  such that  $f \leq cg$ .

**Theorem 4.40**

For  $f \in C_{2\pi}$ ,

$$\|S_N f - f\|_\infty \lesssim (1 + \log N) \tilde{E}_N(f).$$

*Proof.* Recall that  $S_N f = D_N * f$ . Then

$$|S_N f(x)| = \left| \int_{-\pi}^{\pi} D_N(x-y) f(y) dy \right| \leq \int_{-\pi}^{\pi} |D_N(x-y)| |f(y)| dy \leq \|f\|_\infty \int_{-\pi}^{\pi} |D_N(t)| dt.$$

Observe that

$$|D_N(t)| = \frac{1}{2\pi} \left| \frac{\sin((N+1/2)t)}{\sin(t/2)} \right| \leq \min \left\{ \frac{2N+1}{2\pi}, \frac{1}{2|t|} \right\}.$$

Thus

$$\begin{aligned} \int_{-\pi}^{\pi} |D_N(t)| dt &\leq \int_{|t| \leq \pi/(2N+1)} \frac{2N+1}{2\pi} dt + \int_{\pi/(2N+1) \leq |t| \leq \pi} \frac{1}{2|t|} dt \\ &\leq \frac{2N+1}{2\pi} \frac{2\pi}{2N+1} + 2 \cdot \frac{1}{2} \log(2N+1) \lesssim (1 + \log N). \end{aligned}$$

Now let  $q^*$  be the best approximation to  $f$  in  $TP_N$ . Notice that  $S_N q^* = q^*$  and  $S_N$  is a linear operator. Then

$$\|S_N f - q^*\|_\infty = \|S_N f - S_N q^*\|_\infty \leq \|S_N\| \|f - q^*\|_\infty \lesssim (1 + \log N) \tilde{E}_N(f)$$

as desired. ■

**Theorem 4.41**

If  $f \in C_{2\pi}$  is  $L$ -Lipschitz, then

- (a)  $\|\sigma_N f - f\|_\infty \lesssim \frac{1+\log N}{N} L$ ,
- (b)  $\|S_N f - f\|_\infty \lesssim \frac{(1+\log N)^2}{N} L$ .

*Proof.* For (a) we have

$$\sigma_N f(x) = (F_N * f)(x) = \int_{-\pi}^{\pi} F_N(x-y) f(y) dy.$$

And thus

$$\begin{aligned} |\sigma_N f(x) - f(x)| &\leq \int_{-\pi}^{\pi} F_N(x-y) |f(y) - f(x)| dy \\ &= \int_{-\pi}^{\pi} F_N(u) |f(x-u) - f(x)| du \leq L \int_{-\pi}^{\pi} F_N(u) |u| du. \end{aligned}$$

Observe that

$$|F_N(u)| = \frac{1}{2\pi(N+1)} \left| \frac{\sin^2\left(\frac{N+1}{2}u\right)}{\sin^2(u/2)} \right| \leq \min \left\{ \frac{N+1}{2\pi}, \frac{\pi}{2(N+1)|u|^2} \right\}.$$

Then

$$\begin{aligned}
L \int_{-\pi}^{\pi} F_N(u) |u| du &\leq L \int_{|u| \leq \pi/(N+1)} \frac{N+1}{2\pi} |u| du + L \int_{\pi/(N+1) \leq |u| \leq \pi} \frac{\pi}{2(N+1) |u|^2} |u| du \\
&\leq L \int_{|u| \leq \pi/(N+1)} \frac{N+1}{2\pi} \frac{\pi}{N+1} du + \frac{L\pi}{2(N+1)} \cdot 2 \log(N+1) \\
&= \frac{L\pi}{N+1} + L\pi \frac{\log(N+1)}{N+1} \lesssim \frac{1 + \log N}{N} L,
\end{aligned}$$

proving (a).

For (b), from **theorem 4.40** we have

$$\|S_N f - f\|_{\infty} \lesssim (1 + \log N) \tilde{E}_N(f) \lesssim \|\sigma_N f - f\|_{\infty} \lesssim \frac{(1 + \log N)^2}{N} L,$$

proving (b). ■

#### Definition 4.42

The **Chebyshev polynomials** are defined by  $T_n(x) = \cos(n \cos^{-1}(x))$  for  $n = 0, 1, \dots$  on  $[-1, 1]$ .

#### Remark

The Chebyshev polynomials have the following recurrence property,

$$\begin{aligned}
T_{n+1}(x) &= \cos((n+1) \cos^{-1}(x)) = \cos(n \cos^{-1}(x)) \cos(\cos^{-1}(x)) - \sin(n \cos^{-1}(x)) \sin(\cos^{-1}(x)) \\
&= x T_n(x) - \frac{1}{2} \cos((n-1) \cos^{-1}(x)) + \frac{1}{2} \cos((n+1) \cos^{-1}(x)) \\
&= x T_n(x) - \frac{1}{2} T_{n-1}(x) + \frac{1}{2} T_{n+1}(x).
\end{aligned}$$

Then

$$T_{n+1}(x) = 2x T_n(x) - T_{n-1}(x).$$

Immediately we see that  $T_n(x) \in P_n[-1, 1]$ .

#### Proposition 4.43

$\{T_n\}_{n=0}^{\infty}$  forms an orthogonal set with respect to the inner product

$$\langle f, g \rangle_T = \frac{1}{\pi} \int_{-1}^1 f(x) g(x) \frac{dx}{\sqrt{1-x^2}}.$$

*Proof.* Using the change of variable  $x = \cos(\theta)$ , a direct computation gives

$$\begin{aligned}
\langle T_m, T_n \rangle_T &= \frac{1}{\pi} \int_{-1}^1 \cos(m \cos^{-1}(x)) \cos(n \cos^{-1}(x)) \frac{dx}{\sqrt{1-x^2}} = \frac{1}{\pi} \int_0^{\pi} \cos(m\theta) \cos(n\theta) d\theta \\
&= \frac{1}{2\pi} \int_0^{\pi} \cos((m+n)\theta) + \cos((m-n)\theta) d\theta = 0
\end{aligned}$$

for  $m \neq n$ . ■

**Proposition 4.44**

Let  $E[-\pi, \pi]$  be the subspace of  $C[-1, 1]$  consisting of all even continuous function on  $[-\pi, \pi]$ . Consider the mapping  $\Phi : C[-1, 1] \rightarrow E[-\pi, \pi]$  defined by  $\Phi : f \rightarrow f \circ \cos$ . Then the followings are true.

- (a)  $\Phi$  is well-defined and is an isomorphism.
- (b)  $(\Phi T_n)(\theta) = \cos(n\theta)$ .
- (c)  $\Phi(P_n) = E[-\pi, \pi] \cap TP_n$ .
- (d)  $\Phi$  is isometric.
- (e)  $E_n(f) = \tilde{E}_n(\Phi f)$  for all  $f \in C[-1, 1]$ .
- (f)  $\langle f, g \rangle_T = \langle \Phi f, \Phi g \rangle$  for all  $f, g \in C[-1, 1]$ .

*Proof.* For (a), since  $(\Phi f)(-x) = f(\cos(-x)) = f(\cos(x)) = (\Phi f)(x)$  and both  $f$  and  $\cos$  are continuous,  $f \circ \cos$  is also continuous, we conclude that  $\Phi f \in E[-\pi, \pi]$  and  $\Phi$  is well-defined. Now if  $\Phi f = 0$ , then  $\|f\|_\infty = \|\Phi f\|_\infty = 0$  and thus  $f = 0$ . Hence  $\Phi$  is injective. For the surjectivity, let  $g \in E[-\pi, \pi]$ .  $\Phi(g \circ \cos^{-1}) = g$  and  $g \circ \cos^{-1}$  is continuous. Thus  $\Phi$  is surjective.

(b) is immediate from the definition of  $\Phi$ .  $(\Phi T_n)(\theta) = \cos(n \cos^{-1}(\cos(\theta))) = \cos(n\theta)$ .

We now prove (c). From (a) we have  $\Phi$  is an isomorphism. Also, from (b) we have that  $\Phi(P_n) \subset TP_n$ . For any even trigonometric polynomial  $p \in TP_n$ ,  $p(x) = \sum_{k=0}^n a_k \cos(kx)$ . Then consider  $g = \sum_{k=0}^n a_k T_k$ . Then

$$(\Phi g)(x) = \sum_{k=0}^n a_k (T_k \circ \cos)(x) = \sum_{k=0}^n a_k \cos(k \cos^{-1}(\cos(x))) = \sum_{k=0}^n a_k \cos(kx) = p(x).$$

Hence  $TP_n \subset \Phi(P_n)$  and (c) is proven.

For (d),

$$\|\Phi f\|_\infty = \sup_{x \in [-\pi, \pi]} |f(\cos(x))| = \sup_{x \in [-1, 1]} |f(x)| = \|f\|_\infty.$$

(e) is an immediate consequence of (d).

Finally, for (f), by changing the variable  $x = \cos \theta$  and the fact that  $f(\cos(\theta))g(\cos(\theta))$  is even, we have

$$\langle f, g \rangle_T = \frac{1}{\pi} \int_{-1}^1 f(x)g(x) \frac{dx}{\sqrt{1-x^2}} = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\cos(\theta))g(\cos(\theta))d\theta = \langle \Phi f, \Phi g \rangle,$$

showing (f). ■

**Theorem 4.45**

If  $f \in C[-1, 1]$  admits a Chebyshev series

$$f(x) = \sum_{n=0}^{\infty} a_n T_n(x),$$

then

$$\tau_N f = \sum_{k=0}^N \left(1 - \frac{k}{N+1}\right) a_k T_k \rightarrow f$$

uniformly on  $[-1, 1]$ . If  $f$  is Lipschitz and  $P_N^C f$  is the truncated Chebyshev series to  $f$ , then

$$\|P_N^C f - f\|_\infty \lesssim (1 + \log N) E_N(f).$$

*Proof.* Consider the transformation  $\Phi : C[-1, 1] \rightarrow E[-\pi, \pi]$  defined by  $\Phi : f \rightarrow f \circ \cos$ . Then from [proposition 4.44](#) we have

$$(\Phi \tau_N f)(\theta) = \sum_{k=0}^N \left(1 - \frac{k}{N+1}\right) a_k \cos(k\theta) = \frac{1}{N+1} \sum_{j=0}^N \sum_{k=0}^j a_j \cos(j\theta) = \sigma_N(\Phi f)(\theta).$$

By the Fejer theorem,

$$\|\tau_N f - f\|_\infty = \|\Phi \tau_N f - \Phi f\|_\infty = \|\sigma_N(\Phi f) - \Phi f\|_\infty \rightarrow 0$$

as  $N \rightarrow \infty$ .

To see the second part, suppose that  $f$  is  $L$ -Lipschitz. Then

$$|\Phi f(\alpha) - \Phi f(\beta)| \leq L |\cos(\alpha) - \cos(\beta)| \leq L |\alpha - \beta|$$

since the derivative of  $\cos$  is bounded by 1. Thus  $\Phi f$  is  $L$ -Lipschitz and

$$\|P_N^C f - f\|_\infty = \|\Phi P_N^C f - \Phi f\|_\infty = \|\mathcal{S}_N(\Phi f) - \Phi f\|_\infty \lesssim (1 + \log N) \tilde{E}_N(\Phi f) = (1 + \log N) E_N(f)$$

by [theorem 4.40](#) and part (e) of [proposition 4.44](#). ■

**Theorem 4.46** (Jackson)

For  $f \in C[-1, 1]$ ,

$$E_N(f) \lesssim \omega\left(f, \frac{1}{N}\right).$$

*Proof.* Let  $\phi(\theta) = \sum_{k=0}^N c_k \cos(k\theta) \in TP_N$ , where  $c_k \in \mathbb{R}$  such that  $\phi \geq 0$ . For any  $2\pi$ -periodic  $f$ , define  $\Psi$  by

$$\Psi f(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\theta - t) \phi(t) dt = \frac{1}{2\pi} (f * \phi)(\theta).$$

Next we make the following observations. First,  $\Psi \mathbf{1} = \mathbf{1}$  where  $\mathbf{1}$  is the constant function  $\mathbf{1}(\theta) = 1$ . Second,  $\Psi$  is linear and positive. To see this, note that

$$\Psi(cf + g) = \frac{1}{2\pi} \int_{-\pi}^{\pi} (cf + g)(\theta - t) \phi(t) dt = c\Psi f + \Psi g$$

for any  $c \in \mathbb{R}$  and  $f, g \in C[-1, 1]$ . Also, if  $f \geq 0$ , then

$$\Psi f(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\theta - t) \phi(t) dt \geq 0.$$

Third,  $\Psi f \in TP_N$  for any  $f \in C[-1, 1]$ . Indeed,

$$\begin{aligned} \Psi f(\theta) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \phi(\theta - t) dt = \frac{1}{2\pi} \sum_{k=0}^N \int_{-\pi}^{\pi} f(t) c_k \cos(k(\theta - t)) dt \\ &= \frac{1}{2\pi} \sum_{k=0}^N B_k \cos(k\theta) + D_k \sin(k\theta) \in TP_N \end{aligned}$$

with

$$B_k = \frac{c_k}{2\pi} \int_{-\pi}^{\pi} f(t) \cos(kt) dt, \quad \text{and} \quad D_k = \frac{c_k}{2\pi} \int_{-\pi}^{\pi} f(t) \sin(kt) dt.$$

Now we have the last claim that

$$\|\Psi f - f\|_{\infty} \leq \omega(f, \frac{1}{N}) \left( 1 + \frac{N\pi}{2} \sqrt{2 - C} \right)$$

for some constant  $C > 0$ , which will be determined later. Since  $f$  is uniformly continuous,

$$|f(\theta - t) - f(\theta)| \leq \omega(f, |t|) \leq (1 + N|t|) \omega\left(f, \frac{1}{N}\right).$$

Using the first observation  $\Psi \mathbf{1} = \mathbf{1}$ , we have

$$|\Psi f(\theta) - f(\theta)| = \left| \frac{1}{2\pi} \int_{-\pi}^{\pi} (f(\theta - t) - f(\theta)) \phi(t) dt \right| \leq \frac{1}{2\pi} \omega\left(f, \frac{1}{N}\right) \int_{-\pi}^{\pi} (1 + N|t|) \phi(t) dt.$$

Also,

$$\begin{aligned} \frac{1}{2\pi} \int_{-\pi}^{\pi} (1 + N|t|) \phi(t) dt &= 1 + \frac{N}{2\pi} \int_{-\pi}^{\pi} |t| \phi(t) dt \\ &\leq 1 + N \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} |t|^2 \phi(t) dt \right)^{1/2} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi(t) dt \right)^{1/2} \\ &= 1 + N \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} |t|^2 \phi(t) dt \right)^{1/2} \end{aligned}$$

by the Cauchy-Schwarz inequality. Notice that

$$1 - \cos t = 2 \sin^2 \frac{t}{2} \geq 2 \frac{4}{\pi^2} \frac{t^2}{4} = \frac{2t^2}{\pi^2} \Rightarrow |t|^2 \leq \frac{\pi^2}{2} (1 - \cos t).$$



Then

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} (1 + N |t|) \phi(t) dt \leq 1 + N \left( \frac{1}{2\pi} \frac{\pi^2}{2} \int_{-\pi}^{\pi} (1 - \cos t) \phi(t) dt \right)^{1/2} = 1 + \frac{N\pi}{2} \sqrt{2 - C}.$$

Thus

$$\|\Psi f - f\|_{\infty} \leq \omega(f, \frac{1}{N}) \left( 1 + \frac{N\pi}{2} \sqrt{2 - C} \right).$$

Finally, we want to pin down our constant  $C$  so that  $\sqrt{2 - C}$  is minimized and  $\Psi \mathbf{1} = \mathbf{1}$ . We conjecture that  $\phi(\theta) = C_1 |p(\theta)|^2$ , where  $p(\theta) = \sum_{k=0}^N a_k e^{ik\theta}$ ,  $a_k = \sin\left(\frac{k+1}{N+2}\pi\right)$ . Compute that

$$\begin{aligned} C_1 |p(\theta)|^2 &= C_1 p(\theta) \overline{p(\theta)} = C_1 \sum_{k=0}^N a_k e^{ik\theta} \sum_{j=0}^N a_j e^{-ij\theta} \\ &= C_1 \sum_{k=0}^N \sum_{j=0}^N a_k a_j e^{i(k-j)\theta} = C_1 \sum_{k=0}^N a_k^2 + C_1 \sum_{s=1}^N \sum_{k=0}^{N-s} a_k a_{k+s} (e^{is\theta} + e^{-is\theta}) \\ &= C_1 \sum_{k=0}^N a_k^2 + 2C_1 \sum_{s=1}^N \sum_{k=0}^{N-s} a_k a_{k+s} \cos(s\theta). \end{aligned}$$

Take  $C = \left( \sum_{k=0}^N a_k^2 \right)^{-1}$ , then

$$\phi(\theta) = 1 + \sum_{s=1}^N 2C_1 b_s \cos(s\theta), \quad \text{where } b_s = \sum_{k=0}^{N-s} a_k a_{k+s}.$$

Now

$$\begin{aligned} 2b_1 &= \sum_{k=0}^{N-1} 2 \sin\left(\frac{k+1}{N+2}\pi\right) \sin\left(\frac{k+2}{N+2}\pi\right) = \sum_{k=1}^N 2 \sin\left(\frac{k}{N+2}\pi\right) \sin\left(\frac{k+1}{N+2}\pi\right) \\ &= \sum_{k=0}^{N-1} 2 \sin\left(\frac{k}{N+2}\pi\right) \sin\left(\frac{k+1}{N+2}\pi\right). \end{aligned}$$

Combining the first expression and the third one allows us to write

$$\begin{aligned} 2b_1 &= \sum_{k=0}^{N-1} \sin\left(\frac{k+1}{N+2}\pi\right) \left( \sin\left(\frac{k}{N+2}\pi\right) + \sin\left(\frac{k+2}{N+2}\pi\right) \right) \\ &= \sum_{k=0}^{N-1} \sin^2\left(\frac{k+1}{N+2}\pi\right) \cos \frac{2\pi}{N+2} = \cos\left(\frac{2\pi}{N+2}\right) \sum_{k=0}^{N-1} a_k^2 = C_1^{-1} \cos\left(\frac{2\pi}{N+2}\right). \end{aligned}$$

Now

$$C = 2C_1 b_1 = 2 \cos\left(\frac{2\pi}{N+2}\right) \Rightarrow 2 - C \lesssim \frac{1}{N^2}$$

by the Taylor expansion of  $\cos$ . It now follows from the last claim that

$$\|\Psi f - f\|_\infty \leq \omega(f, \frac{1}{N}) \left(1 + \frac{N\pi}{2} \sqrt{2-C}\right) \lesssim \omega(f, \frac{1}{N}).$$

The proof is complete. ■

### 4.3. Fourier Transform

#### Definition 4.47

For  $f \in \mathcal{L}^1(\mathbb{R})$ , its **Fourier transform** is defined as

$$\hat{f}(t) = \mathcal{F}f = \int_{\mathbb{R}} f(x) e^{-2\pi i t x} dx. \quad (1)$$

#### Remark

The Fourier series coefficients can be viewed as discrete Fourier transform  $f \mapsto \{a_n\}_{n \in \mathbb{Z}}$ , with

$$a_n = \int_{-1}^1 f(x) e^{-2\pi i n x} dx. \quad (2)$$

The inverse discrete Fourier transform is then given by

$$f(x) = \sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}. \quad (3)$$

#### Example

$$\hat{\chi}_{[a,b]}(t) = \int_a^b e^{-2\pi i t x} dx = \begin{cases} b-a & \text{if } t = 0, \\ \frac{-1}{2\pi i t} (e^{-2\pi i t b} - e^{-2\pi i t a}) & \text{if } t \neq 0. \end{cases}$$

#### Lemma 4.48 (Riemann-Lebesgue II)

Let  $f \in \mathcal{L}^1(\mathbb{R})$ . Then  $\hat{f}$  is uniformly continuous on  $\mathbb{R}$ , satisfying  $\|\hat{f}\|_\infty \leq \|f\|_1$ , and

$$\lim_{|t| \rightarrow \infty} \hat{f}(t) = 0.$$

*Proof.* We first prove the uniform continuity of  $\hat{f}$ . Let  $t_n \rightarrow t$ . Then since  $|e^{-2\pi i t_n x} f(x)| \leq |f(x)|$ , we may apply the Lebesgue dominated convergence theorem to obtain

$$\lim_{n \rightarrow \infty} \hat{f}(t_n) = \lim_{n \rightarrow \infty} \int_{\mathbb{R}} f(x) e^{-2\pi i t_n x} dx = \int_{\mathbb{R}} f(x) e^{-2\pi i t x} dx = \hat{f}(t).$$

Hence  $\hat{f}$  is uniformly continuous.

To see the second property, we have

$$|\hat{f}(t)| = \left| \int_{\mathbb{R}} f(x) e^{-2\pi i t x} dx \right| \leq \int_{\mathbb{R}} |f(x)| |e^{-2\pi i t x}| dx = \int_{\mathbb{R}} |f(x)| dx = \|f\|_1$$

for any  $t \in \mathbb{R}$  and thus  $\|\hat{f}\|_\infty \leq \|f\|_1$ .

Finally, if  $f = \chi_E$  where  $E = [a, b]$  is an interval, then

$$\hat{f}(t) = \begin{cases} b - a & \text{if } t = 0, \\ \frac{-1}{2\pi it} (e^{-2\pi itb} - e^{-2\pi ita}) & \text{if } t \neq 0. \end{cases}$$

Clearly  $\hat{f}(t) \rightarrow 0$  as  $|t| \rightarrow \infty$ . Since step functions are finite linear combinations of such characteristic functions, the result holds for step functions. For any integrable function, we can find a sequence of step functions  $f_n$  such that  $\|f_n - f\|_1 \rightarrow 0$ . Then

$$|\hat{f}(t) - \hat{f}_n(t)| = \left| \int_{\mathbb{R}} (f(x) - f_n(x)) e^{-2\pi itx} dx \right| \leq \int_{\mathbb{R}} |f(x) - f_n(x)| dx = \|f - f_n\|_1 \rightarrow 0$$

as  $n \rightarrow \infty$ . Since  $\hat{f}_n(t)$  is uniformly continuous and  $\hat{f}_n(t) \rightarrow 0$  as  $|t| \rightarrow \infty$ , we have  $\hat{f}(t) \rightarrow 0$  as well. ■

#### Proposition 4.49

Let  $\hat{f}$  be the Fourier transform of  $f$ .

- (a) If  $f \in \mathcal{L}^1(\mathbb{R})$  and  $g(x) = xf(x) \in \mathcal{L}^1(\mathbb{R})$  as well, then  $\hat{f} \in C^1(\mathbb{R})$  and  $\hat{f}'(t) = -2\pi i \hat{g}(t)$ .
- (b) If  $f \in \mathcal{L}^1(\mathbb{R}) \cap C^1(\mathbb{R})$  and  $f' \in \mathcal{L}^1(\mathbb{R})$ , then

$$(\widehat{f'})(t) = 2\pi it \hat{f}(t).$$

*Proof.* For (a),

$$\frac{1}{s} (\hat{f}(t+s) - \hat{f}(t)) = \frac{1}{s} \int_{\mathbb{R}} f(x) e^{-2\pi i(t+s)x} dx = \int_{\mathbb{R}} f(x) e^{-2\pi itx} \frac{e^{-2\pi isx} - 1}{s} dx.$$

Observe that

$$\left| f(x) e^{-2\pi itx} \frac{1}{s} (e^{-2\pi isx} - 1) \right| \leq |xf(x)| = |g(x)| \in \mathcal{L}^1(\mathbb{R}).$$

By the Lebesgue dominated convergence theorem,

$$\frac{1}{s} (\hat{f}(t+s) - \hat{f}(t)) = \int_{\mathbb{R}} f(x) e^{-2\pi itx} \frac{e^{-2\pi isx} - 1}{s} dx \rightarrow -2\pi i \int_{\mathbb{R}} g(x) e^{-2\pi itx} dx = -2\pi i \hat{g}(t).$$

For (b), using integration by parts,

$$(\widehat{f'})(t) = \int_{\mathbb{R}} f'(x) e^{-2\pi itx} dx = f(x) e^{-2\pi itx} \Big|_{-\infty}^{\infty} + 2\pi it \int_{\mathbb{R}} f(x) e^{-2\pi itx} dx = 2\pi it \hat{f}(t).$$

■

#### Proposition 4.50

Let  $f \in \mathcal{L}^1(\mathbb{R})$  and  $b, t \in \mathbb{R}$ . Then

(a) If  $g(x) = f(x - b)$ ,  $\hat{g}(t) = e^{-2\pi ibt} \hat{f}(t)$ .

(b) If  $g(x) = e^{2\pi ibx} f(x)$ ,  $\hat{g}(t) = \hat{f}(t - b)$ .

(c) If  $g(x) = f(bx)$ ,  $\hat{g}(t) = \frac{1}{|b|} \hat{f}\left(\frac{t}{b}\right)$ .

(d) If  $f, g \in \mathcal{L}^1(\mathbb{R})$ , then

$$\int \hat{f}(t)g(t)dt = \int f(t)\hat{g}(t)dt.$$

*Proof.* For (a), using a translation,

$$\hat{g}(t) = \int f(x - b)e^{-2\pi it(x-b)}dx = \int f(x)e^{-2\pi it(x-b)}dt = e^{-2\pi ibt} \int f(t)e^{-2\pi itx}dx = e^{-2\pi ibt} \hat{f}(t).$$

For (b), using a translation,

$$\hat{g}(t) = \int f(x)e^{2\pi ibx}e^{-2\pi itx}dx = \int f(x)e^{-2\pi it(x-b)}dx = \hat{f}(t - b).$$

For (c), using a dilation,

$$\hat{g}(t) = \int f(bx)e^{-2\pi itx}dx = \frac{1}{|b|} \int f(x)e^{-2\pi itx/b}dx = \frac{1}{|b|} \hat{f}\left(\frac{t}{b}\right).$$

For (d), using Fubini theorem,

$$\int \hat{f}(t)g(t)dt = \int \int f(x)g(t)e^{-2\pi itx}dxdt = \int \int f(x)g(t)e^{-2\pi itx}dt dx = \int f(x)\hat{g}(x)dx.$$

The use of Fubini theorem is justified as follows:

$$\int \int |f(x)g(t)e^{-2\pi itx}|dxdt = \int |g(t)|dt \int |f(x)|dx = \|f\|_1 \|g\|_1 < \infty,$$

since  $f, g \in \mathcal{L}^1(\mathbb{R})$ . ■

#### **Theorem 4.51** (Convolution Theorem)

(a) For  $p \in [1, \infty]$ , if  $f \in \mathcal{L}^1(\mathbb{R})$  and  $g \in \mathcal{L}^p(\mathbb{R})$ , then  $\|f * g\|_p \leq \|f\|_1 \|g\|_p$ .

(b) If  $f, g \in \mathcal{L}^1(\mathbb{R})$ , then  $\widehat{f * g} = \hat{f} \cdot \hat{g}$ .

*Proof.* We first prove (a). For the case  $p = \infty$ ,

$$|(f * g)(x)| = \int |f(y)g(x - y)|dy \leq \|f\|_1 \|g\|_\infty.$$

For the case  $p = 1$ , by Tonelli theorem,

$$\begin{aligned} \|f * g\|_1 &\leq \int \int |f(y)g(x - y)|dydx = \int \int |f(y)| |g(x - y)|dx dy \\ &= \|g\|_1 \int |f(y)|dy = \|f\|_1 \|g\|_1. \end{aligned}$$

For the general case where  $p \in (1, \infty)$ , with  $1/p + 1/p' = 1$ ,

$$\begin{aligned}\|f * g\|_p^p &= \int \left| \int f(x-y)g(y)dy \right|^p dx \leq \int \left( \int |f(x-y)g(y)| dy \right)^p dx \\ &\leq \int \left( \int |f(x-y)| dy \right)^{p/p'} \int |f(x-y)| |g(y)|^p dy dx \\ &= \|f\|_1^{p/p'} \|f * (g^p)\|_1 \leq \|f\|_1^{p/p'} \|g^p\|_1 \|f\|_1 = \|f\|_1^{p/p'} \|f\|_1 \|g\|_p^p.\end{aligned}$$

The second line uses the Hölder inequality and the inequality in the third line uses the result for  $p = 1$ . Now we obtain that

$$\|f * g\|_p = \|f\|_1 \|g\|_p$$

For (b), using Fubini theorem,

$$\begin{aligned}\widehat{f * g}(t) &= \int \int f(x-y)g(y)dy e^{-2\pi ixt} dx = \int \int f(x-y)g(y)e^{-2\pi ixt} dx dy \\ &= \int \int f(x-y)g(y)e^{-2\pi i(x-y)t} d(x-y) e^{-2\pi iyt} dy \\ &= \int g(y)\hat{f}(t)e^{-2\pi iyt} dy = \hat{f}(t)\hat{g}(t).\end{aligned}$$

We verify that  $(x, y) \mapsto |f(x-y)g(y)e^{-2\pi ixt}|$  is integrable. Indeed,

$$\begin{aligned}\int \int |f(x-y)g(y)e^{-2\pi ixt}| dy dx &= \int \int |f(x-y)| |g(y)| dx dy \\ &= \int |g(y)| dy \int |f(x-y)| dx = \|f\|_1 \|g\|_1 < \infty\end{aligned}$$

by Tonelli theorem. The proof is complete. ■

#### Definition 4.52

Given  $\epsilon > 0$ , the **Poisson kernel** is defined as

$$P_\epsilon(x) = \frac{1}{\pi} \frac{\epsilon}{x^2 + \epsilon^2}.$$

#### Proposition 4.53

Let  $P_\epsilon$  be the Poisson kernel. Then

(a)  $P_\epsilon(x) \geq 0$  for all  $x \in \mathbb{R}$  and  $\epsilon > 0$ .

(b) For any  $\epsilon > 0$ ,

$$\int P_\epsilon(x) dx = 1.$$

(c)  $\sup_\epsilon \|P_\epsilon\|_1 \leq M < \infty$  for some  $M > 0$ .

(d) For any given  $\eta > 0$ ,

$$\lim_{\epsilon \rightarrow 0} \int_{|x| > \eta} P_\epsilon(x) dx = 0.$$

*Proof.* (a) is trivial. For (b),

$$\int P_\epsilon(x) dx = \frac{1}{\pi} \epsilon^{-2} \epsilon^2 \pi = 1.$$

(c) follows immediately from (b). For (d), let  $\eta > 0$  be given. Then

$$\int_{|x|>\eta} P_\epsilon(x) dx = \frac{1}{\pi} \int_{|x|>\eta} \frac{\epsilon}{x^2 + \epsilon^2} dx = \frac{2\epsilon}{\pi} \int_\eta^\infty \frac{1}{x^2 + \epsilon^2} dx = \frac{2\epsilon}{\pi} \frac{1}{\epsilon} \left( \frac{\pi}{2} - \tan^{-1}\left(\frac{\eta}{\epsilon}\right) \right) \rightarrow 0$$

as  $\epsilon \rightarrow 0$ . ■

### Remark

The properties (b)-(d) are sometimes referred to as the **good kernel** property. (d) is used to approximate the dirac  $\delta$  function.

### Lemma 4.54

Let  $P_\epsilon$  be the Poisson kernel. Then

(a) If  $f$  is uniformly continuous and bounded on  $\mathbb{R}$ , then  $\|P_\epsilon * f - f\|_\infty \rightarrow 0$  as  $\epsilon \rightarrow 0$ .

(b) If  $f \in \mathcal{L}^p(\mathbb{R})$  where  $1 \leq p < \infty$ , then

$$\|P_\epsilon * f - f\|_p \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0.$$

*Proof.* For (a), we shall proceed with a similar approach in Fejer kernel and Dirichlet kernel. Write

$$\begin{aligned} |P_\epsilon * f(x) - f(x)| &= \left| \int P_\epsilon(x-y) f(y) dy - f(x) \right| = \left| \int P_\epsilon(x-y) (f(y) - f(x)) dy \right| \\ &\leq \int P_\epsilon(x-y) |f(y) - f(x)| dy. \end{aligned}$$

By the uniform continuity of  $f$ , for any  $\delta > 0$ , there exists  $\eta > 0$  such that on  $[x - \eta, x + \eta]$ ,  $|f(y) - f(x)| < \delta$ . Also, by (d) of [proposition 4.53](#), we can choose  $\epsilon$  small enough such that

$$\int_{|x-y|>\eta} P_\epsilon(x-y) dy < \delta.$$

Then we have

$$\begin{aligned} |P_\epsilon * f(x) - f(x)| &\leq \int_{|x-y|\leq\eta} P_\epsilon(x-y) |f(y) - f(x)| dy + \int_{|x-y|>\eta} P_\epsilon(x-y) |f(y) - f(x)| dy \\ &\leq \delta \int_{|x-y|\leq\eta} P_\epsilon(x-y) dy + 2 \|f\|_\infty \int_{|x-y|>\eta} P_\epsilon(x-y) dy \\ &\leq \delta + 2 \|f\|_\infty \delta = \delta(1 + 2 \|f\|_\infty) \end{aligned}$$

by the boundedness of  $f$ . Since  $\delta$  is arbitrary, we obtain that  $\|P_\epsilon * f - f\|_\infty \rightarrow 0$  as  $\epsilon \rightarrow 0$ .

For (b),

$$\begin{aligned}\|P_\epsilon * f - f\|_p^p &= \int \left| \int (f(x) - f(x-y)) P_\epsilon(y) dy \right|^p dx \\ &\leq \int \left( \int |(f(x) - f(x-y))| P_\epsilon(y) dy \right)^p dx \\ &\leq \int \int |f(x) - f(x-y)|^p P_\epsilon(y) dy dx\end{aligned}$$

by Jensen inequality with  $d\mu = P_\epsilon(y)dy$  and **proposition 4.53** (b). Next, by Fubini theorem, letting  $g(y) = \int |f(x) - f(x-y)|^p dx$ ,

$$\begin{aligned}\int \int |f(x) - f(x-y)|^p P_\epsilon(y) dy dx &= \int \int |f(x) - f(x-y)|^p P_\epsilon(y) dx dy \\ &= \int P_\epsilon(y) \int |f(x) - f(x-y)|^p dx dy \\ &= \int P_\epsilon(y) g(y) dy = (P_\epsilon * g)(0) \rightarrow 0\end{aligned}$$

as  $\epsilon \rightarrow 0$  by (a). Thus we conclude that

$$\|P_\epsilon * f - f\|_p \rightarrow 0$$

as  $\epsilon \rightarrow 0$  for any  $1 \leq p < \infty$ . ■

**Theorem 4.55** (Fourier Inversion Theorem)

Suppose  $f, \hat{f} \in \mathcal{L}^1(\mathbb{R})$ . Then

$$f(x) = \int \hat{f}(t) e^{2\pi i t x} dt$$

for almost every  $x \in \mathbb{R}$ .

*Proof.* Consider

$$I_\epsilon(x) = \int \hat{f}(t) e^{-2\pi\epsilon|t|} e^{2\pi i t x} dt.$$

Letting  $g_\epsilon(t; x) = e^{-2\pi\epsilon|t|} e^{2\pi i t x}$ , we have

$$I_\epsilon(x) = \int g_\epsilon(t; x) \hat{f}(t) dt = \int f(t) \hat{g}_\epsilon(t; x) dt$$

since  $g_\epsilon$  is clearly integrable and this follows from **proposition 4.50** (d). Compute that

$$\begin{aligned}\hat{g}_\epsilon(\xi; x) &= \int e^{-2\pi\epsilon|t|} e^{2\pi i t x} e^{-2\pi i \xi t} dt = \int_0^\infty e^{2\pi i t(x-\xi)-\epsilon t} dt + \int_{-\infty}^0 e^{2\pi i t(x-\xi)+\epsilon t} dt \\ &= \frac{-1}{2\pi(i(x-\xi)-\epsilon)} + \frac{1}{2\pi(i(x-\xi)+\epsilon)} = \frac{1}{\pi} \frac{\epsilon}{(x-\xi)^2 + \epsilon^2} = P_\epsilon(x-\xi).\end{aligned}$$

Thus

$$\int f(t) \hat{g}_\epsilon(t; x) dt = \int f(t) P_\epsilon(x - t) dt = (f * P_\epsilon)(x).$$

It follows that  $\|P_\epsilon * f - f\|_1 \rightarrow 0$  as  $\epsilon \rightarrow 0$  by [lemma 4.54](#) (b). It follows that by [theorem 2.25](#) there is a subsequence  $I_{\epsilon_k}(x) \rightarrow f(x)$  almost everywhere. On the other hand,

$$I_\epsilon(x) = \int \hat{f}(t) e^{-2\pi\epsilon|t|} e^{2\pi itx} dt \rightarrow \int \hat{f}(t) e^{2\pi itx} dt$$

as  $\epsilon \rightarrow 0$  by Lebesgue dominated convergence theorem since

$$\left| \hat{f}(t) e^{-2\pi\epsilon|t|} e^{2\pi itx} \right| \leq |\hat{f}(t)| \in \mathcal{L}^1(\mathbb{R}).$$

Thus

$$f(x) = \int \hat{f}(t) e^{2\pi itx} dt.$$

This completes the proof. ■

### Remark

We may also write

$$\hat{\hat{f}}(x) = f(-x).$$

### Definition 4.56

If  $f \in \mathcal{L}^2(\mathbb{R})$ , we define its fourier transform as

$$\hat{f}(t) = \lim_{N \rightarrow \infty} \int_{-N}^N f(x) e^{-2\pi itx} dx.$$

### Theorem 4.57 (Plancherel)

For  $f \in \mathcal{L}^2(\mathbb{R}) \cap \mathcal{L}^1(\mathbb{R})$ ,  $\|\hat{f}\|_2 = \|f\|_2$ .

*Proof.* Directly write

$$\begin{aligned} \|f\|_2^2 &= \int |f(x)|^2 dx = \int f(x) \overline{f(x)} dx = \int f(-x) \overline{f(-x)} dx \\ &= \int \hat{\hat{f}}(x) \overline{\hat{f}(-x)} dx = \int \hat{f}(t) \overline{\hat{f}(t)} dt = \int |\hat{f}(t)|^2 dt = \|\hat{f}\|_2^2. \end{aligned}$$

The second equality in the second line follows from [proposition 4.50](#) (d) and the following fact:

$$\widehat{\overline{f(-x)}}(t) = \int_{-\infty}^{\infty} \overline{f(-x)} e^{-2\pi itx} dx = - \int_{\infty}^{-\infty} \overline{f(u)} e^{2\pi itu} du = \int_{-\infty}^{\infty} \overline{f(u)} \cdot e^{-2\pi itu} du = \overline{\hat{f}(t)},$$

where we have used the change of variable  $u = -x$ . ■

### Definition 4.58



We denote the **fourier transform operator** as

$$\mathcal{F}f(t) = \int_{\mathbb{R}} f(x)e^{-2\pi itx} dx$$

for  $f \in \mathcal{L}^1(\mathbb{R})$ . If  $f \in \mathcal{L}^2(\mathbb{R})$ , we define

$$\mathcal{F}f(t) = \lim_{N \rightarrow \infty} \int_{-N}^N f(x)e^{-2\pi itx} dx$$

instead.

### Remark

From Plancherel theorem, it is immediate that  $\mathcal{F}$  is a bounded linear operator.

### Definition 4.59

The **Schwartz space**  $\mathcal{S}(\mathbb{R})$  is the space of all functions  $f \in C^\infty(\mathbb{R})$  such that

$$\sup_{x \in \mathbb{R}} |x^k D^m f(x)| < \infty$$

for all  $k, m \in \mathbb{N}$ , where  $D^m$  is the  $m$ -th differentiation operator.

### Proposition 4.60

Let  $\mathcal{S}(\mathbb{R})$  be the Schwartz space.

- (a)  $\mathcal{S}(\mathbb{R})$  is a vector space over  $\mathbb{R}$ .
- (b) If  $f \in \mathcal{S}(\mathbb{R})$ , then  $x^k f^{(m)}(x) \in \mathcal{S}(\mathbb{R})$  for all  $k, m \in \mathbb{N} \cup \{0\}$ .
- (c) If  $f \in \mathcal{S}(\mathbb{R})$ , then  $f \in \mathcal{L}^p(\mathbb{R})$  for all  $p \geq 1$ .

*Proof.* For (a), we check that  $\mathcal{S}(\mathbb{R})$  is closed under addition and scalar multiplication. Let  $f, g \in \mathcal{S}(\mathbb{R})$  and  $c \in \mathbb{R}$ . Then  $cf + g$  is also smooth and for  $k, l \in \mathbb{N} \cup \{0\}$ ,

$$\sup_{x \in \mathbb{R}} |x^k (cf + g)^{(l)}(x)| \leq |c| \sup_{x \in \mathbb{R}} |x^k f^{(l)}(x)| + \sup_{x \in \mathbb{R}} |x^k g^{(l)}(x)| < \infty$$

by the definition of Schwartz space. Then  $cf + g \in \mathcal{S}(\mathbb{R})$ , so  $\mathcal{S}(\mathbb{R})$  is a vector space over  $\mathbb{R}$ .

To prove (b), we only need to show the following two facts: first, for any  $f \in \mathcal{S}(\mathbb{R})$ ,  $xf(x) \in \mathcal{S}(\mathbb{R})$ ; second, for any  $f \in \mathcal{S}(\mathbb{R})$ ,  $f'(x) \in \mathcal{S}(\mathbb{R})$ . Suppose that  $f \in \mathcal{S}(\mathbb{R})$ . Then for any  $k, l \in \mathbb{N} \cup \{0\}$ ,

$$\sup_{x \in \mathbb{R}} |x^k (xf(x))^{(l)}| = \sup_{x \in \mathbb{R}} \left| x^k \left( \sum_{i=0}^l \binom{l}{i} x^{(i)} f^{(l-i)}(x) \right) \right| \leq \sup_{x \in \mathbb{R}} |x^{k+1} f^{(l)}(x)| + n \sup_{x \in \mathbb{R}} |x^k f^{(l-1)}(x)| < \infty$$

by the Leibniz formula. Also,

$$\sup_{x \in \mathbb{R}} |x^k (f'(x))^{(l)}| = \sup_{x \in \mathbb{R}} |x^k f^{(l+1)}(x)| < \infty.$$

Thus  $xf(x), f'(x) \in \mathcal{S}(\mathbb{R})$ . In general, the function of the form  $x^k f^{(l)}(x) \in \mathcal{S}(\mathbb{R})$  can be proved by using the above two facts finitely many times.

For (c), let  $E = [-1, 1]$ . By the smoothness of  $f$ , we know that there is some  $M$  such that  $\sup_{x \in E} |f(x)| \leq M$ . Also, from the definition of Schwartz space,  $\sup_{x \in \mathbb{R}} |x^2 f(x)| \leq C$  for some constant  $C$ . Then

$$\begin{aligned} \int |f(x)|^p dx &= \int_E |f(x)|^p dx + \int_{E^c} |f(x)|^p dx \\ &= 2M^p + \int_{E^c} \left| \frac{x^2 f(x)}{x^2} \right|^p dx \\ &\leq 2M^p + C^p \int_{E^c} \frac{1}{x^2} dx = 2M^p + 2C^p < \infty. \end{aligned}$$

Thus  $\|f\|_p < \infty$  for all  $p \geq 1$  and  $p \neq \infty$ . We check that  $f$  is bounded on  $\mathbb{R}$ . By the continuity of  $f$ , we have that  $f$  is always bounded on a compact set. Now if  $f$  does not vanish at infinity, then there is some  $\delta > 0$  and a sequence  $x_n$  such that  $|x_n| \rightarrow \infty$  and  $|f(x_n)| > \delta$ . Then  $\sup_{x \in \mathbb{R}} |xf(x)| \geq \delta \sup_{x \in \mathbb{R}} |x| = \infty$ , posing a contradiction. Thus  $f$  vanishes at infinity. We can find some compact interval  $E$  such that  $\sup_{x \in E} |f(x)| \geq \sup_{x \in E^c} |f(x)|$ . Then by the extreme value theorem,  $f$  is bounded on  $E$  and hence on  $\mathbb{R}$ . We conclude that  $f \in \mathcal{L}^p(\mathbb{R})$  for all  $p \geq 1$ . ■

#### Proposition 4.61

Let  $\mathcal{S}(\mathbb{R})$  be the Schwartz space. If  $f \in \mathcal{S}(\mathbb{R})$ , then  $\hat{f} \in \mathcal{S}(\mathbb{R})$ .

*Proof.* To see this, let  $f \in \mathcal{S}(\mathbb{R})$  be given. From [proposition 4.60](#) (b), we know that  $f$  and  $g(x) = xf(x) \in \mathcal{S}(\mathbb{R}) \subset \mathcal{L}^1(\mathbb{R})$ . Thus  $\hat{f} \in C^1(\mathbb{R})$  and  $\hat{f}'(t) = -2\pi i \hat{g}(t)$ . Since  $g \in \mathcal{S}(\mathbb{R}) \subset \mathcal{L}^1(\mathbb{R})$ , we can repeat the argument to obtain that  $\hat{f} \in C^2$  and  $\hat{f}''(t) = (-2\pi i)^2 \hat{G}(t)$ , where  $G(x) = x^2 f(x)$ . Apply the same argument repeatedly, we have that  $\hat{f} \in C^\infty(\mathbb{R})$  and  $\hat{f}^{(l)}(t) = (-2\pi i)^l \hat{h}(t)$ , where  $h(x) = x^l f(x)$  for all  $l \in \mathbb{N} \cup \{0\}$ . Also,

$$\sup_{x \in \mathbb{R}} |x^k \hat{f}^{(l)}(x)| = \sup_{x \in \mathbb{R}} |x^k (-2\pi i)^l \hat{h}(x)| \leq (2\pi)^l \sup_{x \in \mathbb{R}} |x^k \hat{h}(x)| < \infty.$$

The last inequality follows from the fact that  $h \in \mathcal{S}(\mathbb{R}) \subset \mathcal{L}^1(\mathbb{R})$  and the Riemann-Lebesgue lemma guarantees that  $\hat{h}$  vanishes at infinity. We conclude that  $f \in \mathcal{S}(\mathbb{R})$  implies  $\hat{f} \in \mathcal{S}(\mathbb{R})$ . ■

#### Proposition 4.62

$\mathcal{S}(\mathbb{R})$  is dense in  $\mathcal{L}^p(\mathbb{R})$  for  $1 \leq p < \infty$ .

*Proof.* Since continuous functions with compact support are dense in  $\mathcal{L}^p(\mathbb{R})$ , it suffices to show that  $\mathcal{S}(\mathbb{R})$  is dense in the space of continuous functions with compact support. Without loss of generality, we can assume that  $f$  is supported on  $[-a, a]$  for some  $a > 0$ . By the Weierstrass theorem, we can find a polynomial  $q$  such that  $\|f - q\|_\infty < \epsilon/2$ . Consider the

function

$$\phi_n(t) = \begin{cases} e^{-\frac{1}{n(t^2-a^2)}} & \text{if } |t| < a \\ 0 & \text{if } |t| \geq a. \end{cases}$$

Note that  $\phi_n \rightarrow \chi_{(-a,a)}$  pointwisely as  $n \rightarrow \infty$  and bounded by 1. We verify that  $\phi_n \in \mathcal{S}(\mathbb{R})$  for all  $n \in \mathbb{N}$ . Indeed, for any  $k, l \in \mathbb{N} \cup \{0\}$ , since  $D^l \phi_n$  will result in

$$t^k D^l \phi_n(t) = r(t; n, k, l) e^{-\frac{1}{n(t^2-a^2)}}$$

on  $[-a, a]$  for some rational function  $r(t; n, k, l)$  having singularities only at  $t = \pm a$ , we have that

$$\sup_{t \in \mathbb{R}} |t^k D^l \phi_n(t)| < \infty.$$

Hence,  $\phi_n \in \mathcal{S}(\mathbb{R})$  for all  $n \in \mathbb{N}$ .

Now it follows from **proposition 4.60** (b) that  $q\phi_n \in \mathcal{S}(\mathbb{R})$  by extending the polynomial  $q$  on  $\mathbb{R}$ . Then

$$\begin{aligned} \int |f - q\phi_n|^p d\mu &= \int_{-a}^a |f - q\phi_n|^p d\mu \leq 2^{p-1} \left( \int_{-a}^a |f - q|^p d\mu + \int_{-a}^a |q - q\phi_n|^p d\mu \right) \\ &\leq 2^{p-1} \left( 2a\epsilon^p + \int_{-a}^a |q - q\phi_n|^p d\mu \right) \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$  by the Lebesgue dominated convergence theorem using  $|q - q\phi_n|^p \rightarrow 0$  pointwisely a.e. and  $|q - q\phi_n|^p \leq 2^p |q|^p$  is integrable. The last inequality comes from the convexity  $(x/2 + y/2)^p \leq x^p + y^p$  for  $x, y \geq 0$  and  $p \geq 1$ . We conclude that  $\mathcal{S}(\mathbb{R})$  is dense in  $\mathcal{L}^p(\mathbb{R})$  for  $1 \leq p < \infty$ . ■

### Definition 4.63

A linear operator  $T : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  is said to be **unitary** if

- (a)  $T$  is invertible.
- (b)  $\|Tf\|_2 = \|f\|_1$  for all  $f \in \mathcal{H}_1$ .

### Proposition 4.64

Let  $\mathcal{F}$  be the Fourier transform operator on  $\mathcal{L}^2(\mathbb{R})$ .

- (a)  $\mathcal{F}$  is unitary on  $\mathcal{L}^2(\mathbb{R})$ .
- (b)  $\mathcal{F}^4 = I$ .

*Proof.* (a) is directly from the Plancherel theorem. For (b), using **proposition 4.61**,

$$\mathcal{F}^4 f(t) = \mathcal{F}^2 f(-t) = f(t),$$

by the Fourier inversion theorem for Schwartz functions. Since  $\mathcal{F}$  is unitary, it is also a bounded linear operator, and hence a continuous operator. It now follows from **proposi-**

tion 4.62 that for any  $f \in \mathcal{L}^2(\mathbb{R})$ , there is a sequence  $f_n \in \mathcal{S}(\mathbb{R})$  such that

$$\|f_n - f\|_2 \rightarrow 0.$$

Then

$$\mathcal{F}^4 f(t) = \mathcal{F}^4 \lim_{n \rightarrow \infty} f_n(t) = \lim_{n \rightarrow \infty} \mathcal{F}^4 f_n(t) = \lim_{n \rightarrow \infty} f_n(t) = f(t)$$

for any  $f \in \mathcal{L}^2(\mathbb{R})$  by the continuity of  $\mathcal{F}$ . ■

### Example

We can use the Fourier transform to solve some PDEs. Consider the Laplace equation

$$\begin{cases} \nabla^2 \cdot u = 0 & \text{for } u : \mathbb{R}^2 \rightarrow \mathbb{R}, x \in \mathbb{R}, y > 0, \\ u(x, 0) = f(x) & \text{for } x \in \mathbb{R}. \end{cases}$$

Apply the Fourier transform in  $x$  direction, the original PDE becomes

$$\begin{cases} 4\pi^2 t^2 \hat{u}(t, y) + \hat{u}_{yy}(t, y) = 0 & \text{for } t \in \mathbb{R}, y > 0, \\ \hat{u}(t, 0) = \hat{f}(t) & \text{for } t \in \mathbb{R}. \end{cases}$$

Fix  $t$ , conjecture that  $\hat{u}(t, y) = A(t)e^{-2\pi|t|y} + B(t)e^{2\pi|t|y}$ . Then we have

$$\begin{cases} \hat{u}(t, y) = A(t)e^{-2\pi|t|y}, \\ A(t) = \hat{f}(t). \end{cases}$$

Since

$$\hat{u}(t, y) = \hat{f}(t)e^{-2\pi|t|y} = \hat{f}(t)\hat{P}_y(t) = \widehat{f * P_y}(t),$$

we obtain

$$\begin{cases} u(x, y) = f * P_y(x), \\ \lim_{y \rightarrow 0} u(x, y) = f(x). \end{cases}$$