

Formal Verification of Central Limit Theorem in HOL Theorem Prover

Thi Cam Tu Phan

A thesis submitted for the degree of

Master of Computing (Advanced)

The Australian National University

School of Computing



Australian
National
University

May 2025

© Copyright by Thi Cam Tu Phan, 2025

All Rights Reserved

Abstract

Contents

Contents	3
1 Introduction	5
1.1 Motivation	6
2 Background and Related Work	7
3 Preliminaries	9
4 Central Limit Theorem	11
5 Future Work	13
6 Conclusion	15

Introduction

Formal verification has become an essential approach in modern mathematics and computer science; it provides a strong framework for proving mathematical proofs and algorithms. Traditional techniques of formal verification include the following: Model Checking ?, Testing ?, and Theorem Proving ?.

Model Checking (MC) is a well-valued static formal verification method since, when applied correctly, it provides complete assurance that something satisfies certain specifications. Model checking is a completely automated procedure and is considered easier compared to other techniques such as theorem proving. However, because of the state explosion problem ?, application of model checking in large systems is quite a challenge due to the immense state space size of the systems analyzed. Also, the success of model checking strongly depends on the quality of the analyzed models. In contrast, testing, despite being highly valued for revealing defects in systems, is inherently limited, since it can only indicate the presence of errors, but lacks the ability to verify the correctness of a system.

Theorem proving is another widely used formal analysis technique. It does not suffer from the limitations of state space size as model checking does, which allows the analysis of larger and more complex systems. Furthermore, theorem provers use very expressive logics, like first-order or higher-order logics, which enable the study of a wider range of systems without the restrictions that often come with modeling. The most prominent provers are HOL4 ?, HOL Light ?, Coq ?, and Isabelle/HOL ?, all broadly used in the community.

Formalization plays a vital role in probability theory to ensure that basic results are rigorous mathematically and computable in automated reasoning systems. This project formalizes the Central Limit Theorem using the HOL4 proof assistant, hence enriching the domain of probability in formalized mathematics.

1.1 Motivation

The Central Limit Theorem is one of the most fundamental results in the Probability Theory. This theory finds broad applications in statistics, data science, finance, and

engineering. The theorem underlines that under certain conditions, the sum of many independent random variables will, as the number goes large, converge in distribution to a normal distribution irrespective of the individual distributions of those variables ?. This property justifies many real-world applications and models.

Although HOL4 already contains formal proofs of some fundamental probabilistic results, such as the Law of Large Numbers, a formalized proof of the Central Limit Theorem is still absent. Completing a proof would realize several important milestones, including the following:

- **Enriching the HOL4 Theory Library:**

- The gap in the library would be filled by adding some advanced results from probability theory.
- A more general foundation would be laid for future processes of formal verification.

- **Supporting Future Applications:**

- Strengthening the framework for developers and researchers to model or verify real-world problems involving probabilistic reasoning.

- **Providing Analytical Tools:**

- Enable efficient analysis of problems dependent on the normal distribution and its characteristics.

Formalizing the Central Limit Theorem also aligns with ongoing efforts to bridge traditional mathematical insights with computational tools. As a consequence, it paves the way for applications in areas such as artificial intelligence, machine learning, and quantitative modeling, where probabilistic reasoning is increasingly critical.

This project will contribute to a stronger, more versatile foundation for probabilistic formalization, empowering researchers and developers with better tools to tackle complex, real-world challenges.

Background and Related Work

The history of the Central Limit Theorem, hereinafter referred to as CLT, is quite a fascinating journey through the gradual development of probability theory. Over the years, various versions of CLT have been formulated, reflecting its adaptability and foundational importance. Each such variation addresses certain mathematical challenges and applications, therefore enriching the broader understanding of convergence in probability. In this chapter, we explore some theorem-proving tools and their methodologies to formalize the CLT, emphasizing ...

2.1 Background

Let us brief the few evolvement of CLT, mainly noticing its critical features and variants developed further. The first steps towards CLT took place during the 18th century when Abraham de Moivre showed that, under summing many independent, identically distributed random variables-large number of rolled dice or coin flips-results in the normal distribution-a distribution with the shape of a bell ?. De Moivre laid down the very background for some approximations in the Binomial Distribution.

In 1810, Pierre-Simon Laplace expanded de Moivre's insights by proving that the sum of independent random variables converges to a normal distribution under broader conditions ?. His work established the CLT as a universal principle for analyzing aggregate phenomena, such as population averages and measurement errors.

In the 19th century, mathematicians such as Pafnuty Chebyshev formalized the conditions of the theorem, including those of variance and expectation, so that the theorem became more precise and mathematically sound ?. At the beginning of the 20th century, Lyapunov generalized CLT by introducing specific criteria-Lyapunov's condition-which explained when the theorem was applicable ?. William Feller ? later refined it to address discrete random variables.

Developments in the modern era since World War II have broadened the scope of the CLT beyond normal distributions to stable distributions and applications involving stochastic processes and high-dimensional data. These extensions illustrate how the theorem can

be adapted to a wide range of probabilistic and statistical contexts.

Today, the CLT comes in a variety of variants each suited to particular situations:

- **Classical CLT:** For sums of independent, identically distributed random variables possessing finite variance and mean.
- **Generalized CLT:** When variables are weakly dependent or not identically distributed.
- **Triangular Arrays:** For sums of random variables arranged in arrays where the conditions vary across rows.
- **Local and Integral Versions:** While the local CLT concerns pointwise probabilities, the integral version deals with cumulative distributions.

From about 1810 to 1935, most of the efforts were devoted to proving the CLT for sums of independent random variables, with more recent generalizations involving only weakly dependent variables. Modern formulations neatly distinguish between normed sums, triangular arrays, and local versus integral theorems.

Chapter 3

Preliminaries

Preliminaries 123

Chapter 4

Central Limit Theorem

asldfjkldasjflkdfjldsaf

Chapter 5

Future Work

Future Work

Chapter 6

Conclusion

asdfsdfasdf

asdasasfdfdf