

# Cosystoles and Cheeger Constants of the Simplex

Kai Renken

October 29, 2020

## Table of contents

- 1 Introduction
- 2 On general cosystoles and Cheeger constants
- 3 The first Cheeger constant of a simplex
- 4 Alternative generalizations of the classical Cheeger constant

# An intuitive approach

## An intuitive approach

The classical Cheeger constant intuitively measures the stability of a connected graph.

## An intuitive approach

The classical Cheeger constant intuitively measures the stability of a connected graph.




Figure: A "weakly" connected graph

## An intuitive approach

The classical Cheeger constant intuitively measures the stability of a connected graph.




Figure: A "weakly" connected graph

Deleting the one edge in the middle will give a disconnected graph, consisting of two connected components, the smallest of them consisting of 5 vertices.

## An intuitive approach

The classical Cheeger constant intuitively measures the stability of a connected graph.




Figure: A "weakly" connected graph

Deleting the one edge in the middle will give a disconnected graph, consisting of two connected components, the smallest of them consisting of 5 vertices. The Cheeger constant of this graph is  $\frac{1}{5}$ .

# The definition of the classical Cheeger constant



## The definition of the classical Cheeger constant

### Definition

Let  $G = (V, E)$  be a (simple) graph. Then the **Cheeger constant** of  $G$  is defined by

$$h(G) = \min \left\{ \frac{|\delta(A)|}{|A|} : A \subset V, 1 \leq |A| \leq \frac{|V|}{2} \right\},$$

with  $\delta(A) := \{e = (v, w) \in E : v \in A, w \in V \setminus A\}$ .

## The definition of the classical Cheeger constant

### Definition

Let  $G = (V, E)$  be a (simple) graph. Then the **Cheeger constant** of  $G$  is defined by

$$h(G) = \min \left\{ \frac{|\delta(A)|}{|A|} : A \subset V, 1 \leq |A| \leq \frac{|V|}{2} \right\},$$

with  $\delta(A) := \{e = (v, w) \in E : v \in A, w \in V \setminus A\}$ .




Figure: A "weakly" connected graph

# The classical Cheeger constant of the simplex

## The classical Cheeger constant of the simplex

Let  $K_n$  be the complete graph on  $n$  vertices (which is the 1-skeleton of the standard simplex on  $n$ -vertices).

## The classical Cheeger constant of the simplex

Let  $K_n$  be the complete graph on  $n$  vertices (which is the 1-skeleton of the standard simplex on  $n$ -vertices).

Then we have:

$$h(K_n) = \left\lceil \frac{n}{2} \right\rceil$$

as follows:

## The classical Cheeger constant of the simplex

Let  $K_n$  be the complete graph on  $n$  vertices (which is the 1-skeleton of the standard simplex on  $n$ -vertices).

Then we have:

$$h(K_n) = \left\lceil \frac{n}{2} \right\rceil$$

as follows:

For any subset  $A \subset [n] := \{1, \dots, n\}$  we have:

## The classical Cheeger constant of the simplex

Let  $K_n$  be the complete graph on  $n$  vertices (which is the 1-skeleton of the standard simplex on  $n$ -vertices).

Then we have:

$$h(K_n) = \left\lceil \frac{n}{2} \right\rceil$$

as follows:

For any subset  $A \subset [n] := \{1, \dots, n\}$  we have:

$$\frac{|\delta(A)|}{|A|} = \frac{|A|(n - |A|)}{|A|} = n - |A|,$$

## The classical Cheeger constant of the simplex

Let  $K_n$  be the complete graph on  $n$  vertices (which is the 1-skeleton of the standard simplex on  $n$ -vertices).

Then we have:

$$h(K_n) = \left\lceil \frac{n}{2} \right\rceil$$

as follows:

For any subset  $A \subset [n] := \{1, \dots, n\}$  we have:

$$\frac{|\delta(A)|}{|A|} = \frac{|A|(n - |A|)}{|A|} = n - |A|,$$

so by  $|A| \leq \frac{n}{2}$  we get:



## The classical Cheeger constant of the simplex

Let  $K_n$  be the complete graph on  $n$  vertices (which is the 1-skeleton of the standard simplex on  $n$ -vertices).

Then we have:

$$h(K_n) = \left\lceil \frac{n}{2} \right\rceil$$

as follows:

For any subset  $A \subset [n] := \{1, \dots, n\}$  we have:

$$\frac{|\delta(A)|}{|A|} = \frac{|A|(n - |A|)}{|A|} = n - |A|,$$

so by  $|A| \leq \frac{n}{2}$  we get:

$$h(K_n) = n - \left\lfloor \frac{n}{2} \right\rfloor = \left\lceil \frac{n}{2} \right\rceil$$

# Cosystoles and the cosystolic norm

## Cosystoles and the cosystolic norm

From now we consider all cochain groups with  $\mathbb{Z}_2$ -coefficients.

## Cosystoles and the cosystolic norm

From now we consider all cochain groups with  $\mathbb{Z}_2$ -coefficients.

### Definition

Let  $X$  be an (abstract) simplicial complex. Then the **norm** of a cochain  $\varphi \in C^k(X)$  is defined by:

$$\|\varphi\| := |\text{supp}(\varphi)|$$

## Cosystoles and the cosystolic norm

From now we consider all cochain groups with  $\mathbb{Z}_2$ -coefficients.

### Definition

Let  $X$  be an (abstract) simplicial complex. Then the **norm** of a cochain  $\varphi \in C^k(X)$  is defined by:

$$\|\varphi\| := |\text{supp}(\varphi)|$$

The **cosystolic norm** of a cochain  $\varphi \in C^k(X)$  (for  $k \geq 1$ ) is defined by:

$$\|\varphi\|_{\text{csy}} := \min \left\{ \|\delta^{k-1}(\phi) + \varphi\| : \phi \in C^{k-1}(X) \right\}$$

## Cosystoles and the cosystolic norm

From now we consider all cochain groups with  $\mathbb{Z}_2$ -coefficients.

### Definition

Let  $X$  be an (abstract) simplicial complex. Then the **norm** of a cochain  $\varphi \in C^k(X)$  is defined by:

$$\|\varphi\| := |\text{supp}(\varphi)|$$

The **cosystolic norm** of a cochain  $\varphi \in C^k(X)$  (for  $k \geq 1$ ) is defined by:

$$\|\varphi\|_{\text{csy}} := \min \left\{ \|\delta^{k-1}(\phi) + \varphi\| : \phi \in C^{k-1}(X) \right\}$$

A cochain  $\varphi \in C^k(X)$  is called a  **$k$ -cosystole**, if it satisfies  $\|\varphi\|_{\text{csy}} = \|\varphi\|$ .

## Coboundary expansion and the $k$ -th Cheeger constant

### Definition

For a cochain  $\varphi \in C^k(X) \setminus \text{Im}(\delta^{k-1})$  the quotient

$$\|\varphi\|_{\text{exp}} := \frac{\|\delta^k(\varphi)\|}{\|\varphi\|_{\text{csy}}}$$

is called the **coboundary expansion** of  $\varphi$  and

## Coboundary expansion and the $k$ -th Cheeger constant

### Definition

For a cochain  $\varphi \in C^k(X) \setminus \text{Im}(\delta^{k-1})$  the quotient

$$\|\varphi\|_{\text{exp}} := \frac{\|\delta^k(\varphi)\|}{\|\varphi\|_{\text{csy}}}$$

is called the **coboundary expansion** of  $\varphi$  and

$$h_k(X) := \min_{\substack{\varphi \in C^k(X) \\ \varphi \notin \text{Im}(\delta^{k-1})}} \|\varphi\|_{\text{exp}}$$

is called the  $k$ -th **Cheeger constant** of  $X$ .



## Coboundary expansion and the $k$ -th Cheeger constant

### Definition

For a cochain  $\varphi \in C^k(X) \setminus \text{Im}(\delta^{k-1})$  the quotient

$$\|\varphi\|_{\text{exp}} := \frac{\|\delta^k(\varphi)\|}{\|\varphi\|_{\text{csy}}}$$

is called the **coboundary expansion** of  $\varphi$  and

$$h_k(X) := \min_{\substack{\varphi \in C^k(X) \\ \varphi \notin \text{Im}(\delta^{k-1})}} \|\varphi\|_{\text{exp}}$$

is called the  $k$ -th **Cheeger constant** of  $X$ .

A cosystole  $\varphi \in C^k(X) \setminus \text{Im}(\delta^{k-1})$  satisfying  $\|\varphi\|_{\text{exp}} = h_k(X)$  is called a **Cheeger cosystole**.

## Relations to the classical Cheeger constant

## Relations to the classical Cheeger constant

The classical Cheeger constant of a graph can be considered as the 0-th Cheeger constant by defining the cosystolic norm of a 0-cochain  $\varphi \in C^0(X)$  as

$$\|\varphi\|_{csy} := \min\{|\text{supp}(\varphi)|, |X^{(0)}| - |\text{supp}(\varphi)|\}$$

## Relations to the classical Cheeger constant

The classical Cheeger constant of a graph can be considered as the 0-th Cheeger constant by defining the cosystolic norm of a 0-cochain  $\varphi \in C^0(X)$  as

$$\|\varphi\|_{csy} := \min\{|\text{supp}(\varphi)|, |X^{(0)}| - |\text{supp}(\varphi)|\}$$

The classical Cheeger constant of a graph equals 0 iff the graph is disconnected.

## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

Let  $\varphi \in C^k(X)$ , then we have:

## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

Let  $\varphi \in C^k(X)$ , then we have:

$$\|\varphi\|_{\text{csy}} > 0 \Leftrightarrow \varphi \notin \text{Im}(\delta^{k-1})$$

## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

Let  $\varphi \in C^k(X)$ , then we have:

$$\|\varphi\|_{\text{csy}} > 0 \Leftrightarrow \varphi \notin \text{Im}(\delta^{k-1})$$

and



## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

Let  $\varphi \in C^k(X)$ , then we have:

$$\|\varphi\|_{\text{csy}} > 0 \Leftrightarrow \varphi \notin \text{Im}(\delta^{k-1})$$

and

$$\delta^k(\varphi) = 0 \Leftrightarrow \varphi \in \ker(\delta^k),$$

## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

Let  $\varphi \in C^k(X)$ , then we have:

$$\|\varphi\|_{\text{csy}} > 0 \Leftrightarrow \varphi \notin \text{Im}(\delta^{k-1})$$

and

$$\delta^k(\varphi) = 0 \Leftrightarrow \varphi \in \ker(\delta^k),$$

so the existence of a cochain  $\varphi \in C^k(X)$  satisfying  $\|\varphi\|_{\text{csy}} > 0$  and  $\delta^k(\varphi) = 0$  is equivalent to

$$\text{Im}(\delta^{k-1}) \subsetneq \ker(\delta^k),$$

## Relations to the classical Cheeger constant

The  $k$ -th Cheeger constant of a simplicial complex  $X$  equals 0 iff the  $k$ -th cohomology group  $H^k(X)$  is non-trivial, as follows:

Let  $\varphi \in C^k(X)$ , then we have:

$$\|\varphi\|_{\text{csy}} > 0 \Leftrightarrow \varphi \notin \text{Im}(\delta^{k-1})$$

and

$$\delta^k(\varphi) = 0 \Leftrightarrow \varphi \in \ker(\delta^k),$$

so the existence of a cochain  $\varphi \in C^k(X)$  satisfying  $\|\varphi\|_{\text{csy}} > 0$  and  $\delta^k(\varphi) = 0$  is equivalent to

$$\text{Im}(\delta^{k-1}) \subsetneq \ker(\delta^k),$$

which is equivalent to

$$H^k(X) \neq \{0\}$$

## Example: The 1-cosystoles of $\Delta^{[6]}$

## Example: The 1-cosystoles of $\Delta^{[6]}$

Let  $\Delta^{[n]}$  always denote the standard simplex on  $n$  vertices.

## Example: The 1-cosystoles of $\Delta^{[6]}$

Let  $\Delta^{[n]}$  always denote the standard simplex on  $n$  vertices.




Figure: The supports of all 1-cosystoles of  $\Delta^{[6]}$  (up to isomorphism)

## Some values for higher dimensional Cheeger constants

## Some values for higher dimensional Cheeger constants

### Theorem (Wallach, Meshulam)

Let  $\Delta^{[n]}$  be the standard simplex on  $n$  vertices and  $1 \leq k \leq n-2$ , then we have:

$$\frac{n}{k+2} \leq h_k(\Delta^{[n]}) \leq \left\lceil \frac{n}{k+2} \right\rceil$$

If  $n$  is divisible by  $k+2$ , then we have:

$$h_k(\Delta^{[n]}) = \frac{n}{k+2}$$



## Some values for higher dimensional Cheeger constants

### Theorem (Wallach, Meshulam)

Let  $\Delta^{[n]}$  be the standard simplex on  $n$  vertices and  $1 \leq k \leq n-2$ , then we have:

$$\frac{n}{k+2} \leq h_k(\Delta^{[n]}) \leq \left\lceil \frac{n}{k+2} \right\rceil$$

If  $n$  is divisible by  $k+2$ , then we have:

$$h_k(\Delta^{[n]}) = \frac{n}{k+2}$$

### Theorem (Kozlov)

Let  $n > 2$  not be a power of 2, then we have:

$$h_1(\Delta^{[n]}) = \frac{n}{3}$$

# Hitting sets and hitting numbers

## Hitting sets and hitting numbers

### Definition

Let  $V$  be some set and  $\mathcal{F} \subseteq 2^V$  a family of subsets of  $V$ . A subset  $P \subseteq V$  is called a **hitting set** of  $\mathcal{F}$  if we have  $P \cap F \neq \emptyset$  for all  $F \in \mathcal{F}$ . The **hitting number** of  $\mathcal{F}$  is defined by

$$\tau(\mathcal{F}) := \min\{|P| : P \subseteq V \text{ is a hitting set of } \mathcal{F}\}$$

## Hitting sets and hitting numbers

### Definition

Let  $V$  be some set and  $\mathcal{F} \subseteq 2^V$  a family of subsets of  $V$ . A subset  $P \subseteq V$  is called a **hitting set** of  $\mathcal{F}$  if we have  $P \cap F \neq \emptyset$  for all  $F \in \mathcal{F}$ . The **hitting number** of  $\mathcal{F}$  is defined by

$$\tau(\mathcal{F}) := \min\{|P| : P \subseteq V \text{ is a hitting set of } \mathcal{F}\}$$

If  $\mathcal{F}$  is a family of chains / cochains, the hitting sets and the hitting number of  $\mathcal{F}$  are defined as the hitting sets and the hitting number of  $\{\text{supp}(\varphi) : \varphi \in \mathcal{F}\}$ .

## Hitting sets and hitting numbers

### Definition

Let  $V$  be some set and  $\mathcal{F} \subseteq 2^V$  a family of subsets of  $V$ . A subset  $P \subseteq V$  is called a **hitting set** of  $\mathcal{F}$  if we have  $P \cap F \neq \emptyset$  for all  $F \in \mathcal{F}$ . The **hitting number** of  $\mathcal{F}$  is defined by

$$\tau(\mathcal{F}) := \min\{|P| : P \subseteq V \text{ is a hitting set of } \mathcal{F}\}$$

If  $\mathcal{F}$  is a family of chains / cochains, the hitting sets and the hitting number of  $\mathcal{F}$  are defined as the hitting sets and the hitting number of  $\{\text{supp}(\varphi) : \varphi \in \mathcal{F}\}$ .

### Example

Let  $V := \{1, 2, 3, 4, 5\}$  and  $\mathcal{F} := \{\{1, 2\}, \{2, 3, 4\}, \{1, 5\}, \{2, 4, 5\}\}$ , then we have  $\tau(\mathcal{F}) = 2$ .

# The cycle detection theorem

## The cycle detection theorem

### Theorem (Kozlov)

*Let  $X$  be a simplicial complex,  $k \geq 1$ , and  $\varphi \in C^k(X)$ . Let now  $\mathcal{F} = \{\alpha_1, \dots, \alpha_t\}$  be a family of  $k$ -cycles in  $C_k(X)$ , such that  $\langle \varphi, \alpha_i \rangle = 1$  for all  $1 \leq i \leq t$ , then we have:*

$$\|\varphi\|_{\text{csy}} \geq \tau(\mathcal{F})$$

## The cycle detection theorem

Proof.



## The cycle detection theorem

Proof.

Let  $\psi \in C^{k-1}(X)$ , then for any  $1 \leq i \leq t$  we have:

## The cycle detection theorem

Proof.

Let  $\psi \in C^{k-1}(X)$ , then for any  $1 \leq i \leq t$  we have:

$$\begin{aligned}\langle \varphi + \delta^{k-1}(\psi), \alpha_i \rangle &= \langle \varphi, \alpha_i \rangle + \langle \delta^{k-1}(\psi), \alpha_i \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, \partial_{k-1}(\alpha_i) \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, 0 \rangle \\ &= \langle \varphi, \alpha_i \rangle = 1\end{aligned}$$

## The cycle detection theorem

Proof.

Let  $\psi \in C^{k-1}(X)$ , then for any  $1 \leq i \leq t$  we have:

$$\begin{aligned}\langle \varphi + \delta^{k-1}(\psi), \alpha_i \rangle &= \langle \varphi, \alpha_i \rangle + \langle \delta^{k-1}(\psi), \alpha_i \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, \partial_{k-1}(\alpha_i) \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, 0 \rangle \\ &= \langle \varphi, \alpha_i \rangle = 1\end{aligned}$$

This means that we have  $\text{supp}(\varphi + \delta^{k-1}(\psi)) \cap \text{supp}(\alpha_i) \neq \emptyset$  for all  $1 \leq i \leq t$ , so  $\text{supp}(\varphi + \delta^{k-1}(\psi))$  is a hitting set of  $\mathcal{F}$  and we get:

## The cycle detection theorem

Proof.

Let  $\psi \in C^{k-1}(X)$ , then for any  $1 \leq i \leq t$  we have:

$$\begin{aligned}\langle \varphi + \delta^{k-1}(\psi), \alpha_i \rangle &= \langle \varphi, \alpha_i \rangle + \langle \delta^{k-1}(\psi), \alpha_i \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, \partial_{k-1}(\alpha_i) \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, 0 \rangle \\ &= \langle \varphi, \alpha_i \rangle = 1\end{aligned}$$

This means that we have  $\text{supp}(\varphi + \delta^{k-1}(\psi)) \cap \text{supp}(\alpha_i) \neq \emptyset$  for all  $1 \leq i \leq t$ , so  $\text{supp}(\varphi + \delta^{k-1}(\psi))$  is a hitting set of  $\mathcal{F}$  and we get:

$$\|\varphi + \delta^{k-1}(\psi)\| = |\text{supp}(\varphi + \delta^{k-1}(\psi))| \geq \tau(\mathcal{F})$$

## The cycle detection theorem

Proof.

Let  $\psi \in C^{k-1}(X)$ , then for any  $1 \leq i \leq t$  we have:

$$\begin{aligned}\langle \varphi + \delta^{k-1}(\psi), \alpha_i \rangle &= \langle \varphi, \alpha_i \rangle + \langle \delta^{k-1}(\psi), \alpha_i \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, \partial_{k-1}(\alpha_i) \rangle \\ &= \langle \varphi, \alpha_i \rangle + \langle \psi, 0 \rangle \\ &= \langle \varphi, \alpha_i \rangle = 1\end{aligned}$$

This means that we have  $\text{supp}(\varphi + \delta^{k-1}(\psi)) \cap \text{supp}(\alpha_i) \neq \emptyset$  for all  $1 \leq i \leq t$ , so  $\text{supp}(\varphi + \delta^{k-1}(\psi))$  is a hitting set of  $\mathcal{F}$  and we get:

$$\|\varphi + \delta^{k-1}(\psi)\| = |\text{supp}(\varphi + \delta^{k-1}(\psi))| \geq \tau(\mathcal{F})$$

Since  $\psi$  was chosen arbitrarily we are done. □

# The cycle detection theorem

## The cycle detection theorem

### Corollary (Kozlov)

*Let  $\varphi \in C^k(X)$  and  $\mathcal{F} = \{\alpha_1, \dots, \alpha_{\|\varphi\|}\} \subset C_k(X)$  be a family of  $k$ -cycles, such that their supports are pairwise disjoint, then  $\varphi$  is a cosystole.*

# The cycle detection theorem



## The cycle detection theorem

Question: Are all cosystoles detectable using the cycle detection theorem?

## The cycle detection theorem

Question: Are all cosystoles detectable using the cycle detection theorem?

Answer: We can not prove it by now, but we also do not know any counterexample.

## The cycle detection theorem

Question: Are all cosystoles detectable using the cycle detection theorem?

Answer: We can not prove it by now, but we also do not know any counterexample.

Question: Are all cosystoles detectable using disjoint cycles?

## The cycle detection theorem

Question: Are all cosystoles detectable using the cycle detection theorem?

Answer: We can not prove it by now, but we also do not know any counterexample.

Question: Are all cosystoles detectable using disjoint cycles?

Answer: No. The following 1-cosystole is not detectable this way:




Figure: The support of a 1-cosystole, which can not be determined using disjoint cycles

## The cycle detection theorem

Question: Are all cosystoles detectable using the cycle detection theorem?

Answer: We can not prove it by now, but we also do not know any counterexample.

Question: Are all cosystoles detectable using disjoint cycles?

Answer: No. The following 1-cosystole is not detectable this way:




Figure: The support of a 1-cosystole, which can not be determined using disjoint cycles

Conjecture: For every proper  $n$  and  $k$  there is a Cheeger cosystole in  $C^k(\Delta^{[n]})$  which is detectable using disjoint cycles.

# Maximal cosystoles

# Maximal cosystoles

## Definition

Let  $X$  be a simplicial complex and  $1 \leq k \leq \dim(X)$ , then

$$C_{\max}(X, k) := \max \left\{ \|\varphi\|_{\text{csy}} : \varphi \in C^k(X) \right\}$$

is the largest norm a  $k$ -cosystole in  $X$  can attain.

# The largest cosystoles of the simplex



# The largest cosystoles of the simplex

Theorem (Renken)

$$C_{\max}(\Delta^{[n]}, 1) = \binom{\lceil \frac{n}{2} \rceil}{2} + \binom{\lfloor \frac{n}{2} \rfloor}{2}$$

## The case when $n$ is a power of 2

## The case when $n$ is a power of 2

### Conjecture

*Let  $n$  be a power of 2. Then we have:*

$$h_1(\Delta^{[n]}) > \frac{n}{3}$$

## The case when $n$ is a power of 2

### Conjecture

*Let  $n$  be a power of 2. Then we have:*

$$h_1(\Delta^{[n]}) > \frac{n}{3}$$

### Theorem (Kozlov)

$$h_1(\Delta^{[8]}) > \frac{8}{3}$$

## The case when $n$ is a power of 2

### Conjecture

*Let  $n$  be a power of 2. Then we have:*

$$h_1(\Delta^{[n]}) > \frac{n}{3}$$

### Theorem (Kozlov)

$$h_1(\Delta^{[8]}) > \frac{8}{3}$$

### Theorem (Renken)

$$h_1(\Delta^{[16]}) > \frac{16}{3}$$

# Disjoint cycle expansion

## Disjoint cycle expansion

### Definition

Let  $X$  be a simplicial complex,  $\mathcal{F} \subset C_k(X)$  a family of cycles, such that their supports are pairwise disjoint and

$$P(\mathcal{F}) := \{\varphi \in C^k(X) : |\text{supp}(\varphi) \cap \text{supp}(F)| = 1 \text{ for all } F \in \mathcal{F} \text{ and } \text{supp}(\varphi) \subset \bigcup_{F \in \mathcal{F}} \text{supp}(F)\}$$

## Disjoint cycle expansion

### Definition

Let  $X$  be a simplicial complex,  $\mathcal{F} \subset C_k(X)$  a family of cycles, such that their supports are pairwise disjoint and

$$P(\mathcal{F}) := \{\varphi \in C^k(X) : |\text{supp}(\varphi) \cap \text{supp}(F)| = 1 \text{ for all } F \in \mathcal{F} \text{ and } \text{supp}(\varphi) \subset \bigcup_{F \in \mathcal{F}} \text{supp}(F)\}$$

Then we define

$$\gamma_{\mathcal{F}} := \frac{\min_{\varphi \in P(\mathcal{F})} \|\delta^k(\varphi)\|}{|\mathcal{F}|}$$



## Disjoint cycle expansion

### Definition

Let  $X$  be a simplicial complex,  $\mathcal{F} \subset C_k(X)$  a family of cycles, such that their supports are pairwise disjoint and

$$P(\mathcal{F}) := \{\varphi \in C^k(X) : |\text{supp}(\varphi) \cap \text{supp}(F)| = 1 \text{ for all } F \in \mathcal{F} \text{ and } \text{supp}(\varphi) \subset \bigcup_{F \in \mathcal{F}} \text{supp}(F)\}$$

Then we define

$$\gamma_{\mathcal{F}} := \frac{\min_{\varphi \in P(\mathcal{F})} \|\delta^k(\varphi)\|}{|\mathcal{F}|}$$

and we call

$$\gamma_k(X) := \min_{\mathcal{F} \in \mathfrak{C}} \gamma_{\mathcal{F}}$$

the  $k$ -th **disjoint cycle expansion** of  $X$  with

$$\mathfrak{C} := \{\mathcal{F} \subset C_k(X) : F \text{ is a cycle and } \text{supp}(F) \cap \text{supp}(F') = \emptyset \text{ for all } F, F' \in \mathcal{F} (F \neq F')\}$$

# Hitting expansion

## Hitting expansion

### Definition

Let  $X$  be a simplicial complex,  $\mathcal{F} \subset C_k(X)$  a family of cycles and

$$P'(\mathcal{F}) := \{\varphi \in C^k(X) : |\text{supp}(\varphi) \cap \text{supp}(F)| \text{ is odd for all } F \in \mathcal{F}\}$$

## Hitting expansion

### Definition

Let  $X$  be a simplicial complex,  $\mathcal{F} \subset C_k(X)$  a family of cycles and

$$P'(\mathcal{F}) := \{\varphi \in C^k(X) : |\text{supp}(\varphi) \cap \text{supp}(F)| \text{ is odd for all } F \in \mathcal{F}\}$$

Then we define

$$\rho_{\mathcal{F}} := \frac{\min_{\varphi \in P'(\mathcal{F})} \|\delta^k(\varphi)\|}{\tau(\mathcal{F})}$$

## Hitting expansion

### Definition

Let  $X$  be a simplicial complex,  $\mathcal{F} \subset C_k(X)$  a family of cycles and

$$P'(\mathcal{F}) := \{\varphi \in C^k(X) : |\text{supp}(\varphi) \cap \text{supp}(F)| \text{ is odd for all } F \in \mathcal{F}\}$$

Then we define

$$\rho_{\mathcal{F}} := \frac{\min_{\varphi \in P'(\mathcal{F})} \|\delta^k(\varphi)\|}{\tau(\mathcal{F})}$$

and we call

$$\rho_k(X) := \min_{\mathcal{F} \in \mathfrak{C}} \rho_{\mathcal{F}}$$

the  $k$ -th **hitting expansion** of  $X$ , with

$$\mathfrak{C} := \{\mathcal{F} \subset C_k(X) : F \text{ is a cycle, for all } F \in \mathcal{F}\}$$

## Relations to the Cheeger constant

## Relations to the Cheeger constant

### Theorem (Kozlov)

*Let  $X$  be a simplicial complex and  $k \geq 1$ , then we have:*

$$h_k(X) \leq \rho_k(X) \leq \gamma_k(X)$$

## Relations to the Cheeger constant

### Theorem (Kozlov)

*Let  $X$  be a simplicial complex and  $k \geq 1$ , then we have:*

$$h_k(X) \leq \rho_k(X) \leq \gamma_k(X)$$

### Theorem (Kozlov)

*Let  $k + 2$  divide  $n$ , then we have:*

$$\gamma_k(\Delta^{[n]}) = \rho_k(\Delta^{[n]}) = \frac{n}{k+2}$$



## Relations to the Cheeger constant

### Theorem (Kozlov)

*Let  $X$  be a simplicial complex and  $k \geq 1$ , then we have:*

$$h_k(X) \leq \rho_k(X) \leq \gamma_k(X)$$

### Theorem (Kozlov)

*Let  $k + 2$  divide  $n$ , then we have:*

$$\gamma_k(\Delta^{[n]}) = \rho_k(\Delta^{[n]}) = \frac{n}{k+2}$$

### Theorem (Renken)

*Let  $n$  not be a power of 2, then we have:*

$$\gamma_1(\Delta^{[n]}) = \rho_1(\Delta^{[n]}) = \frac{n}{3}$$

## Bibliography



Dmitry N. Kozlov, *The first Cheeger constant of a simplex*, Graphs and Combinatorics (2017) 33: 1543.  
<https://doi.org/10.1007/s00373-017-1853-9>



N. Linial, R. Meshulam, *Homological connectivity of random 2-complexes*, Combinatorica 26, 2006, no. 4, 475-487



M. Gromov, *Singularities, expanders and topology of maps. Part 2. From combinatorics to topology via algebraic isoperimetry*, Geom. Funct. Anal. 20, (2010), no. 2, 416-526.



M. Wallach and R. Meshulam, *Homological connectivity of random  $k$ -dimensional complexes*, Random Structures Algorithms 34, 2009, no. 3, 408-417



Dmitry N. Kozlov and Roy Meshulam, *Quantitative aspects of acyclicity*, arXiv:1802.03210 [math.CO], 2018