

A cohomological perspective for high order problems (I)

Kaibo Hu

University of Minnesota

Chinese Academy of Sciences

December 01, 2020

- 1 Motivation
- 2 de Rham complexes
- 3 Finite element sequences

1 Motivation

2 de Rham complexes

3 Finite element sequences

High order problems

Examples of high order problems

- plate (thin structures):

$$\Delta^2 u = f,$$

- model problem for electromagnetic and generalized continuum:


$$-\operatorname{curl} \operatorname{div}(\operatorname{grad} \operatorname{curl}) u = f,$$

$$\operatorname{div} u = 0.$$

$\operatorname{div} \operatorname{grad} = \Delta = \operatorname{curl} \operatorname{curl} - \operatorname{grad} \operatorname{div}$, formally $-\operatorname{curl} \operatorname{div}(\operatorname{grad} \operatorname{curl}) = \operatorname{curl}^4$.

Challenges for finite element discretization:

- conformity,
- inf-sup conditions (for vector/tensor fields and multi-fields).



Fluid mechanics

Stokes equations : strongly diffusive, stationary, incompressible

$$\begin{cases} -\Delta \mathbf{u} + \nabla p &= \mathbf{f}, \\ \nabla \cdot \mathbf{u} &= 0, \end{cases}$$

Boundary condition: $\mathbf{u} = 0$ on $\partial\Omega$.

Variational form : find $\mathbf{u} \in [H_0^1(\Omega)]^n$, $p \in L_0^2(\Omega)$ s.t.,

$$\begin{cases} (\nabla \mathbf{u}, \nabla \mathbf{v}) - (p, \nabla \cdot \mathbf{v}) &= (\mathbf{f}, \mathbf{v}), \quad \forall \mathbf{v} \in [H_0^1(\Omega)]^n, \\ (\nabla \cdot \mathbf{u}, q) &= 0, \quad \forall q \in L_0^2(\Omega). \end{cases}$$

Discrete variational form : find $\mathbf{u}_h \in \mathbf{V}_h$, $p \in Q_h$ s.t.,

$$\begin{cases} (\nabla \mathbf{u}_h, \nabla \mathbf{v}_h) - (p_h, \nabla \cdot \mathbf{v}_h) &= (\mathbf{f}, \mathbf{v}_h), \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \\ (\nabla \cdot \mathbf{u}_h, q_h) &= 0, \quad \forall q_h \in Q_h. \end{cases}$$

How to choose \mathbf{V}_h and Q_h ?

- **numerical stability** (inf-sup condition): for any $q_h \in Q_h$, there exists $\mathbf{v}_h \in \mathbf{V}_h$, $\|\mathbf{v}_h\| = 1$, s.t. $(\nabla \cdot \mathbf{v}_h, q_h) \geq \|q_h\|_{L^2}$; $\nabla \cdot \mathbf{V}_h$ should be “larger” than Q_h .
- precise **divergence-free constraint**:

$$(\nabla \cdot \mathbf{v}_h, q_h) = 0, \forall q_h \in Q_h \implies \nabla \cdot \mathbf{v}_h = 0 \quad Q_h \text{ should be “larger” than } \nabla \cdot \mathbf{V}_h.$$

mass conservation, important for numerics, John et al. *SIAM Review* 2017

- balance: $\nabla \cdot \mathbf{V}_h = Q_h$,
- constructing finite elements satisfying $\nabla \cdot \mathbf{V}_h = Q_h$ turns out to be very challenging.

Scott-Vogelius, stable Stokes pairs etc.

1 Motivation

2 de Rham complexes

3 Finite element sequences

de Rham complex

de Rham complex (3D version)


$$0 \longrightarrow C^\infty(\Omega) \xrightarrow{\text{grad}} C^\infty(\Omega; \mathbb{R}^3) \xrightarrow{\text{curl}} C^\infty(\Omega; \mathbb{R}^3) \xrightarrow{\text{div}} C^\infty(\Omega) \longrightarrow 0.$$

$$d^0 := \text{grad}, \quad d^1 := \text{curl}, \quad d^2 := \text{div}.$$

- complex property: $d^k \circ d^{k-1} = 0$, $\Rightarrow \mathcal{R}(d^{k-1}) \subset \mathcal{N}(d^k)$,
 $\text{curl} \circ \text{grad} = 0 \Rightarrow \mathcal{R}(\text{grad}) \subset \mathcal{N}(\text{curl})$, $\text{div} \circ \text{curl} = 0 \Rightarrow \mathcal{R}(\text{curl}) \subset \mathcal{N}(\text{div})$
- cohomology: $\mathcal{H}^k := \mathcal{N}(d^k) / \mathcal{R}(d^{k-1})$,
 $\mathcal{H}^0 := \mathcal{N}(\text{grad})$, $\mathcal{H}^1 := \mathcal{N}(\text{curl}) / \mathcal{R}(\text{grad})$, $\mathcal{H}^2 := \mathcal{N}(\text{div}) / \mathcal{R}(\text{curl})$
- exactness (contractible domains): $\mathcal{N}(d^k) = \mathcal{R}(d^{k-1})$, i.e., $d^k u = 0 \Rightarrow u = d^{k-1} v$
 $\text{curl } u = 0 \Rightarrow u = \text{grad } \phi$, $\text{div } v = 0 \Rightarrow v = \text{curl } \psi$.

de Rham complex and topology:

dimension of \mathcal{H}^k = number of “k-dimensional holes” (c.f. de Rham theorem)



Examples where $\dim \mathcal{H}^1 = 1$ and $\dim \mathcal{H}^2 = 1$, respectively.

Left: curl-free field which is not grad, Right: div-free field which is not curl.

(figure from *Finite element exterior calculus*, D.N.Arnold, SIAM 2008.)

From complexes to PDEs

Formal adjoint of operators:

$$\operatorname{grad}^* = -\operatorname{div}, \quad \operatorname{curl}^* = \operatorname{curl}, \quad \operatorname{div}^* = -\operatorname{grad}.$$

$$\int_{\Omega} \operatorname{grad} u \cdot v = - \int_{\Omega} u \operatorname{div} v + \text{bound. term}, \quad \int_{\Omega} \operatorname{curl} u \cdot v = \int_{\Omega} u \cdot \operatorname{curl} v + \text{bound. term}$$

$$(\operatorname{grad} u, v) = (u, -\operatorname{div} v), \quad (\operatorname{curl} u, v) = (u, \operatorname{curl} v)$$

Formal adjoint of de Rham complex:

$$0 \longleftarrow C^{\infty}(\Omega) \xleftarrow{-\operatorname{div}} C^{\infty}(\Omega; \mathbb{R}^3) \xleftarrow{\operatorname{curl}} C^{\infty}(\Omega; \mathbb{R}^3) \xleftarrow{-\operatorname{grad}} C^{\infty}(\Omega) \longleftarrow 0.$$

$$d_2^* := -\operatorname{div}, \quad d_1^* := \operatorname{curl}, \quad d_0^* := -\operatorname{grad}.$$

connections to PDEs: Hodge-Laplacian problems.

$$(d^{k-1}d_{k-1}^* + d_k^*d^k)u = f.$$

connections to PDEs: Hodge-Laplacian problems.

$$(d^{k-1}d_{k-1}^* + d_k^*d^k)u = f.$$

$$0 \begin{array}{c} \longrightarrow \\ \longleftarrow \end{array} \mathcal{C}^\infty(\Omega) \begin{array}{c} \xrightarrow{\text{grad}} \\ \xleftarrow{-\text{div}} \end{array} \mathcal{C}^\infty(\Omega; \mathbb{R}^3) \qquad \mathcal{C}^\infty(\Omega; \mathbb{R}^3) \qquad \mathcal{C}^\infty(\Omega) \qquad 0.$$

Hodge-Laplacian problem:

$$-\operatorname{div} \operatorname{grad} u = f.$$

Poisson equation.

connections to PDEs: Hodge-Laplacian problems.

$$(d^{k-1}d_{k-1}^* + d_k^*d^k)u = f.$$

$$0 \quad C^\infty(\Omega) \begin{array}{c} \xrightarrow{\text{grad}} \\ \xleftarrow{-\text{div}} \end{array} C^\infty(\Omega; \mathbb{R}^3) \begin{array}{c} \xrightarrow{\text{curl}} \\ \xleftarrow{\text{curl}} \end{array} C^\infty(\Omega; \mathbb{R}^3) \quad C^\infty(\Omega) \quad 0.$$

Hodge-Laplacian problem:

$$-\text{grad div } v + \text{curl curl } v = f.$$

Maxwell equations.

connections to PDEs: Hodge-Laplacian problems.

$$(d^{k-1}d_{k-1}^* + d_k^*d^k)u = f.$$

$$0 \quad C^\infty(\Omega) \quad C^\infty(\Omega; \mathbb{R}^3) \begin{array}{c} \xrightarrow{\text{curl}} \\ \xleftarrow{\text{curl}} \end{array} C^\infty(\Omega; \mathbb{R}^3) \begin{array}{c} \xrightarrow{\text{div}} \\ \xleftarrow{-\text{grad}} \end{array} C^\infty(\Omega) \quad 0.$$

Hodge-Laplacian problem:

$$\text{curl curl } v - \text{grad div } v = f.$$

Maxwell equations.

connections to PDEs: Hodge-Laplacian problems.

$$(d^{k-1}d_{k-1}^* + d_k^*d^k)u = f.$$

$$0 \quad C^\infty(\Omega) \quad C^\infty(\Omega; \mathbb{R}^3) \quad C^\infty(\Omega; \mathbb{R}^3) \begin{array}{c} \xrightarrow{\text{div}} \\ \xleftarrow{-\text{grad}} \end{array} C^\infty(\Omega) \rightleftharpoons 0.$$

Hodge-Laplacian problem:


$$-\operatorname{div} \operatorname{grad} u = f.$$

Poisson equation.

Sobolev version and discretization

$$0 \longrightarrow H(\text{grad}) \xrightarrow{\text{grad}} H(\text{curl}) \xrightarrow{\text{curl}} H(\text{div}) \xrightarrow{\text{div}} L^2 \longrightarrow 0$$

$$H(d) := \{u \in L^2 : du \in L^2\}.$$



$$0 \longrightarrow \mathcal{P}_1 \xrightarrow{\text{grad}} [\mathcal{P}_0]^3 + [\mathcal{P}_0]^3 \times x \xrightarrow{\text{curl}} [\mathcal{P}_0]^3 + \mathcal{P}_0 \otimes x \xrightarrow{\text{div}} \mathcal{P}_0 \longrightarrow 0.$$

Raviart-Thomas (1977), Nédélec (1980) in numerical analysis, Bossavit (1988), Hiptmair (1999) for differential forms, Whitney (1957) for studying topology.

Smoother de Rham complexes

- connecting high order problems to Stokes problem

$$0 \longrightarrow H^1 \xrightarrow{\text{grad}} H(\text{grad curl}) \xrightarrow{\text{curl}} [H^1]^3 \xrightarrow{\text{div}} L^2 \longrightarrow 0,$$

$$0 \longrightarrow H^2 \xrightarrow{\text{grad}} H^1(\text{curl}) \xrightarrow{\text{curl}} [H^1]^3 \xrightarrow{\text{div}} L^2 \longrightarrow 0,$$

where $H^1(\text{curl}) := \{u \in [H^1]^3 : \text{curl } u \in [H^1]^3\}$.

- exactness (cohomology): c.f., Costabel, McIntosh 2010, for *any real number* s ,

$$0 \longrightarrow H^s \xrightarrow{\text{grad}} H^{s-1} \otimes \mathbb{R}^3 \xrightarrow{\text{curl}} H^{s-2} \otimes \mathbb{R}^3 \xrightarrow{\text{div}} H^{s-3} \longrightarrow 0$$


1 Motivation

2 de Rham complexes

3 Finite element sequences

Conservative Stokes discretization: $[H^1]^3 - L^2$

- puzzle of Scott-Vogelius $([C^0\mathcal{P}_r]^n - C^{-1}\mathcal{P}_{r-1})$: 2D stable for $r \geq 4$, no “singular vertices”; 3D open.




- a cohomological perspective

$$0 \longrightarrow H^1 \xrightarrow{\text{grad}} H(\text{grad curl}) \xrightarrow{\text{curl}} [H^1]^3 \xrightarrow{\text{div}} L^2 \longrightarrow 0.$$

$$0 \longrightarrow C^0 \text{ spline} \xrightarrow{\text{grad}} * \xrightarrow{\text{curl}} \mathbf{V}_h \xrightarrow{\text{div}} Q_h \longrightarrow 0.$$

(Incomplete) review on finite element Stokes complexes (on simplicial meshes)

- first FE Stokes complex: Falk-Neilan 2013,





- 3D tetrahedral element: Neilan 2015 (starting with \mathcal{P}_9-C^1), Q.Zhang-Z.Zhang 2020 (H^1 complex, different 1-forms),
- construction on macroelements: Christiansen-Hu 2018 (low order complex), Fu-Guzmán-Neilan 2018 (Alfeld split, any dim), Guzmán-Lischke-Neilan, 2020 (Powell-Sabin, Worsey-Farin split),
- nonconforming elements: Mardal-Tai-Winther 2002, Tai-Winther 2006, Huang 2020 (fewer dofs),
- virtual elements: Zhao-Zhang-Mao-Chen 2019 (nonconforming), Beirão da Veiga-Dassi-Vacca 2020 (conforming 3D).


Stabilize Scott-Vogelius elements

inf-sup condition: velocity space large enough, to “control” pressure


- piecewise constant pressure \leftarrow face dofs of velocity, $\int_T \operatorname{div} u = \int_{\partial T} u \cdot n$,
- higher order pressure modes \leftarrow interior dofs of velocity.



- Bernadi-Raugel bubble: $b_{BR} := \{b_F n_F\}$, scalar face bubble b_F , but $\operatorname{div} b_{BR} \notin \mathcal{P}_0$!
- modified Bernadi-Raugel bubble: $b_{mBR} := \{b_F n_F + \sum_{j=1}^k \lambda_F^j w_{k-j}\}$, $\operatorname{div} b_{mBR} \subset \mathcal{P}_0$.



tetrahedra construction with canonical dofs: Hu-Q.Zhang-Z.Zhang 2020,
arXiv:2008.03793



- construct 1-form by Poincaré integrals p^2 :

$$V_h^1 := \text{grad } V_h^1 \oplus p^2 V_h^2.$$

basic idea of p^2 : integral of functions

- canonical dofs: Whitney dofs + vertex evaluation.

Poincaré operators

Question:

- how to find out explicit potential? $u = D\phi$, knowing u , find out ϕ .
- how to prove the Poincaré lemma (local exactness of de Rham sequence)?

Example: $D = \text{grad}$, $\phi = \phi(x_0) + \int_{\gamma(y)} u \, dy$



General d^k (curl, div etc.):

- Poincaré operators (differential geometry books; Hiptmair 1999)
 $\mathfrak{p}^k : C^\infty \Lambda^k \mapsto C^\infty \Lambda^{k-1}$, satisfying

$$d^{k-1}\mathfrak{p}^k + \mathfrak{p}^{k+1}d^k = \text{id}_{C^\infty \Lambda^k},$$

- exactness: $du = 0 \Rightarrow u = (d\mathfrak{p} + \mathfrak{p}d)u = d(\mathfrak{p}u)$.

$$\cdots \rightleftarrows V^{i-1} \begin{array}{c} \xrightarrow{d^{i-1}} \\ \xleftarrow{\mathfrak{p}^i} \end{array} V^i \begin{array}{c} \xrightarrow{d^i} \\ \xleftarrow{\mathfrak{p}^{i+1}} \end{array} V^{i+1} \rightleftarrows \cdots$$

Corollary: $V^i = dV^{i-1} \oplus \mathfrak{p}V^{i+1}$.

- explicit forms of \mathfrak{p}^k for de Rham complexes: using Cartan's magic formula
- 3D vector proxy (with $W = 0$):

$$\mathfrak{p}_1 u = \int_0^1 u_{tx} \cdot x dt, \quad \mathfrak{p}_2 v = \int_0^1 t v_{tx} \times x dt, \quad \mathfrak{p}_3 w = \int_0^1 t^2 w_{tx} x dt.$$

applications in numerical analysis: constructing polynomial exact sequences. e.g.,

$$0 \longrightarrow \mathcal{P}_1 \xrightarrow{\text{grad}} [\mathcal{P}_0]^3 + [\mathcal{P}_0]^3 \times x \xrightarrow{\text{curl}} [\mathcal{P}_0]^3 + \mathcal{P}_0 \otimes x \xrightarrow{\text{div}} \mathcal{P}_0 \longrightarrow 0.$$

$$0 \longrightarrow \mathcal{P}_r \xrightarrow{\text{grad}} [\mathcal{P}_{r-1}]^3 \xrightarrow{\text{curl}} [\mathcal{P}_{r-2}]^3 \xrightarrow{\text{div}} \mathcal{P}_{r-3} \longrightarrow 0.$$

Finite element exterior calculus (FEEC): cohomological approach for numerical PDEs (Arnold, Falk, Winther 2006 Acta. Num., Arnold, Falk, Winther 2010 Bulletin of AMS, Arnold 2018 SIAM book).

Stokes case

$$V_h^1 := \text{grad } V_h^1 \oplus \mathfrak{p}^2 V_h^2.$$

interior point chosen as base point.

Take home messages:

- complex is important,
- enriching classical Scott-Vogelius pair with face and interior bubbles,
- application of Poincaré operators.

Overview of part 2: spurious solutions and grad curl-complexes

$$0 \longrightarrow H^q \xrightarrow{\text{grad}} H^{q-1} \otimes \mathbb{V} \xrightarrow{\text{grad curl}} H^{q-3} \otimes \mathbb{T} \xrightarrow{\text{curl}} H^{q-4} \otimes \mathbb{M} \xrightarrow{\text{div}} H^{q-5} \otimes \mathbb{V} \longrightarrow 0.$$

References

- *A family of finite element Stokes complexes in three dimensions*, Kaibo Hu, Qian Zhang, Zhimin Zhang; 2020, arXiv:2008.03793.
- *Simple curl-curl-conforming finite elements in two dimensions*; Kaibo Hu, Qian Zhang, Zhimin Zhang; SIAM Scientific Computing 2020 (accepted).
- *Generalized finite element systems for smooth differential forms and Stokes problem*, Snorre H. Christiansen and Kaibo Hu; Numerische Mathematik 2018.